

# Discontinuous Distribution in Bees

Nature

130, 58-59

DOI: [10.1038/130058b0](https://doi.org/10.1038/130058b0)

Citation Report

| # | ARTICLE   | IF  | CITATIONS |
|---|---|-----|-----------|
| 1 | Metachronal rhythms and gill movements of the nymph of <i>caenis horaria</i> (Ephemeroptera) in relation to water flow. Proceedings of the Royal Society of London Series B, Biological Sciences, 1934, 115, 30-48. | 3.0 | 11        |
| 2 | A CLIMATIC PATTERN BETWEEN LATITUDES 40° AND 70° N. AND ITS PROBABLE INFLUENCE ON BIOLOGICAL DISTRIBUTIONS. Proceedings of the Linnean Society of London, 1960, 171, 89-121.  | 0.8 | 2         |
| 3 | Insect Pollination of High Arctic Flowers. Journal of Ecology, 1972, 60, 831.   | 4.0 | 210       |
| 4 | Benthic macroinvertebrate distributions in the riffle-pool communities of two east Texas streams. Hydrobiologia, 1986, 135, 61-70.  | 2.0 | 44        |
| 5 | Strong zonation of benthic communities across a tidal freshwater height gradient. Freshwater Biology, 2019, 64, 1284-1294.  | 2.4 | 4         |
| 6 | Revision of the bee genus <i>Capicola</i> (Hymenoptera: Apoidea: Melittidae) distributed in the Southwest of Africa. European Journal of Entomology, 2007, 104, 311-340.  | 1.2 | 16        |