

CITATION REPORT

List of articles citing

Determination of fatty acid ethyl esters (FAEE) and ethyl glucuronide (EtG) in hair: a promising way for retrospective detection of alcohol abuse during pregnancy?

DOI: 10.1097/ftd.ob013e318167d602

Therapeutic Drug Monitoring, 2008, 30, 255-63.

Source: <https://exaly.com/paper-pdf/45010156/citation-report.pdf>

Version: 2024-04-24

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
132	Ethyl glucuronide determination: head hair versus non-head hair. <i>Alcohol and Alcoholism</i> , 2009 , 44, 62-6	3.5	65
131	Liquid chromatography with tandem mass spectrometric detection for the measurement of ethyl glucuronide and ethyl sulfate in meconium: new biomarkers of gestational ethanol exposure?. <i>Therapeutic Drug Monitoring</i> , 2008 , 30, 725-32	3.2	35
130	Ethylglucuronide determination in urine and hair from alcohol withdrawal patients. <i>Journal of Analytical Toxicology</i> , 2009 , 33, 155-61	2.9	27
129	The effect of hair pigment on the incorporation of fatty acid ethyl esters (FAEE). <i>Alcohol and Alcoholism</i> , 2009 , 44, 287-92	3.5	25
128	ChemInform Abstract: Determination of Fatty Acid Ethyl Esters (FAEE) and Ethyl Glucuronide (ETG) in Hair: A Promising Way for Retrospective Detection of Alcohol Abuse During Pregnancy?. 2009 , 40, no		
127	Development and validation of a gas chromatography-negative chemical ionization tandem mass spectrometry method for the determination of ethyl glucuronide in hair and its application to forensic toxicology. 2009 , 877, 2337-43		71
126	Determination of ethyl-glucuronide in hair for heavy drinking detection using liquid chromatography-tandem mass spectrometry following solid-phase extraction. 2009 , 394, 1895-901		39
125	Ethyl glucuronide in hair. A sensitive and specific marker of chronic heavy drinking. 2009 , 104, 915-20		78
124	Ethyl glucuronide in hair compared with traditional alcohol biomarkers--a pilot study of heavy drinkers referred to an alcohol detoxification unit. 2009 , 33, 812-6		43
123	Comparison of ethyl glucuronide in hair with carbohydrate-deficient transferrin in serum as markers of chronic high levels of alcohol consumption. <i>Forensic Science International</i> , 2009 , 188, 140-3	2.6	40
122	Hair analysis of fatty acid ethyl esters in the detection of excessive drinking in the context of fetal alcohol spectrum disorders. <i>Therapeutic Drug Monitoring</i> , 2009 , 31, 261-6	3.2	32
121	Marqueurs biologiques de l'alcoolisme. 2010 , 5, 1-9		2
120	Population Baseline of Meconium Ethyl Glucuronide and Ethyl Sulfate Concentrations in Newborns of Nondrinking Women in 2 Mediterranean Cohorts. <i>Therapeutic Drug Monitoring</i> , 2010 , 32, 359-63	3.2	32
119	Agreement between the fatty acid ethyl ester hair test for alcohol and social workers' reports. <i>Therapeutic Drug Monitoring</i> , 2010 , 32, 294-9	3.2	15
118	A fully validated high-performance liquid chromatography-tandem mass spectrometry method for the determination of ethyl glucuronide in hair for the proof of strict alcohol abstinence. 2010 , 396, 2441-7		40
117	Quantification of fatty acid ethyl esters (FAEE) and ethyl glucuronide (EtG) in meconium from newborns for detection of alcohol abuse in a maternal health evaluation study. 2010 , 396, 2469-77		79
116	Analytical methods for abused drugs in hair and their applications. 2010 , 397, 1039-67		49

115	Fatty acid ethyl ester concentrations in hair and self-reported alcohol consumption in 644 cases from different origin. <i>Forensic Science International</i> , 2010 , 196, 111-7	2.6	55
114	Determination of ethyl glucuronide in hair samples of Chinese people by protein precipitation (PPT) and large volume injection-gas chromatography-tandem mass spectrometry (LVI-GC/MS/MS). 2010 , 878, 3161-6		27
113	Combined use of fatty acid ethyl esters and ethyl glucuronide in hair for diagnosis of alcohol abuse: interpretation and advantages. <i>Forensic Science International</i> , 2010 , 196, 101-10	2.6	125
112	Correlation between drugs of abuse and alcohol by hair analysis: parents at risk for having children with fetal alcohol spectrum disorder. 2010 , 44, 615-21		17
111	Influence of ethanol dose and pigmentation on the incorporation of ethyl glucuronide into rat hair. 2010 , 44, 507-14		39
110	Effect of bleaching on ethyl glucuronide in hair: an in vitro experiment. <i>Forensic Science International</i> , 2010 , 198, 23-7	2.6	73
109	Ethyl glucuronide and ethyl sulfate in meconium and hair-potential biomarkers of intrauterine exposure to ethanol. <i>Forensic Science International</i> , 2010 , 196, 74-7	2.6	70
108	Ethyl glucuronide determination in meconium and hair by hydrophilic interaction liquid chromatography-tandem mass spectrometry. <i>Forensic Science International</i> , 2010 , 196, 121-7	2.6	45
107	Alcohol biomarkers in applied settings: recent advances and future research opportunities. 2010 , 34, 955-67		113
106	Determination of Bioactive Components in 300°C Pyrolyzate of Extract from Oil-Tea Cake by Pyrolysis- GC/MS. 2010 ,		
105	Headspace microextraction: recent bioanalytical applications and issues. 2010 , 2, 123-41		6
104	Molecular targets of alcohol action: Translational research for pharmacotherapy development and screening. 2011 , 98, 293-347		14
103	Hair: a complementary source of bioanalytical information in forensic toxicology. 2011 , 3, 67-79		52
102	Handbook of hair in health and disease. 2011 ,		5
101	Changes in transferrin glycosylation during pregnancy may lead to false-positive carbohydrate-deficient transferrin (CDT) results in testing for riskful alcohol consumption. 2011 , 412, 129-33		48
100	Determinaci3n de drogas de abuso en pelo. 2011 , 37, 59-66		1
99	Phosphatidylethanol in blood (B-PEth): a marker for alcohol use and abuse. <i>Drug Testing and Analysis</i> , 2011 , 3, 195-200	3.5	94
98	Ethyl-glucuronide and ethyl-sulfate in placental and fetal tissues by liquid chromatography coupled with tandem mass spectrometry. 2011 , 418, 30-6		29

97	Alkoholkonsummarker. <i>Rechtsmedizin</i> , 2011 , 21, 69-79	0.6	6
96	Comparison of ethyl glucuronide (EtG) and fatty acid ethyl esters (FAEEs) concentrations in hair for testing abstinence. 2011 , 400, 175-81		48
95	Impact of hair-care products on FAEE hair concentrations in substance abuse monitoring. 2011 , 400, 183-8		53
94	Chemometric evaluation of nine alcohol biomarkers in a large population of clinically-classified subjects: pre-eminence of ethyl glucuronide concentration in hair for confirmatory classification. 2011 , 401, 2153-64		43
93	Determination of fatty acid ethyl esters in hair by GC-MS and application in a population of cocaine users. 2011 , 54, 1192-5		16
92	Gas chromatography/mass spectrometry based hair steroid profiling may reveal pathogenesis in hair follicles of the scalp. 2011 , 25, 1184-92		26
91	Is there a better way to monitor abstinence among substance abusers awaiting transplantation?. 2012 , 17, 180-7		2
90	Workplace alcohol testing program by combined use of ethyl glucuronide and fatty acid ethyl esters in hair. <i>Alcohol and Alcoholism</i> , 2012 , 47, 127-32	3.5	38
89	Metabolomics via Biomedical Mass Spectrometry: From Sampling to Clinical Applications. 205-224		
88	[Ethylglucuronide assays in urine and hair]. 2012 , 70, 629-34		1
87	Preliminary investigations on ethyl glucuronide and ethyl sulfate cutoffs for detecting alcohol consumption on the basis of an ingestion experiment and on data from withdrawal treatment. 2012 , 126, 757-64		31
86	Determination of maternal-fetal biomarkers of prenatal exposure to ethanol: a review. 2012 , 69, 209-22		89
85	Comparison of ethyl glucuronide in hair with self-reported alcohol consumption. <i>Alcohol and Alcoholism</i> , 2012 , 47, 267-72	3.5	37
84	Toxicology testing in alternative specimen matrices. 2012 , 32, 467-92		19
83	Alcohol biomarkers. 2012 , 32, 391-406		35
82	A preliminary investigation of the AUDIT and DUDIT in comparison to biomarkers for alcohol and drug use among HIV-infected clinic attendees in Cape Town, South Africa. 2012 , 15, 346-51		21
81	Ethyl glucuronide in human hair after daily consumption of 16 or 32 g of ethanol for 3 months. <i>Forensic Science International</i> , 2012 , 215, 51-5	2.6	75
80	Can ethyl glucuronide in hair be determined only in 3 cm hair strands?. <i>Forensic Science International</i> , 2012 , 218, 3-9	2.6	21

79	Practical experiences in application of hair fatty acid ethyl esters and ethyl glucuronide for detection of chronic alcohol abuse in forensic cases. <i>Forensic Science International</i> , 2012 , 218, 82-91	2.6	76
78	Interpretation problems in a forensic case of abstinence determination using alcohol markers in hair. <i>Forensic Science International</i> , 2012 , 217, e4-7	2.6	16
77	The influence of ethanol containing cosmetics on ethyl glucuronide concentration in hair. <i>Forensic Science International</i> , 2012 , 218, 123-5	2.6	51
76	Diagnostic performance of ethyl glucuronide in hair for the investigation of alcohol drinking behavior: a comparison with traditional biomarkers. 2012 , 126, 243-50		63
75	A comparison of the performance of quality controls prepared from spiked, fortified and authentic hair for ethyl glucuronide analysis. <i>Forensic Science International</i> , 2013 , 232, 60-6	2.6	9
74	Fatty acid ethyl esters in hair as alcohol markers: estimating a reliable cut-off point by evaluation of 1,057 autopsy cases. 2013 , 9, 184-93		7
73	Determination of ethyl glucuronide levels in hair for the assessment of alcohol abstinence. <i>Forensic Science International</i> , 2013 , 232, 229-36	2.6	33
72	Application of mass spectrometry to hair analysis for forensic toxicological investigations. 2013 , 32, 312-32		51
71	Non-oxidative ethanol metabolites as a measure of alcohol intake. 2013 , 415, 322-9		51
70	Sensitivity and specificity of urinary ethyl glucuronide and ethyl sulfate in liver disease patients. 2013 , 37, 150-5		42
69	Higher levels of hair ethyl glucuronide in patients with decreased kidney function. 2013 , 37 Suppl 1, E14-6		23
68	Hair ethyl glucuronide is highly sensitive and specific for detecting moderate-to-heavy drinking in patients with liver disease. <i>Alcohol and Alcoholism</i> , 2013 , 48, 83-7	3.5	37
67	[Direct metabolites of ethanol as biological markers of alcohol use: basic aspects and applications]. 2013 , 81, 493-502		12
66	Diagnostic value of concentration profiles of glucocorticosteroids and endocannabinoids in hair. <i>Therapeutic Drug Monitoring</i> , 2013 , 35, 600-7	3.2	35
65	Coloring, bleaching, and perming: influence on EtG content in hair. <i>Therapeutic Drug Monitoring</i> , 2013 , 35, 527-9	3.2	56
64	A comparison of the different animal models of fetal alcohol spectrum disorders and their use in studying complex behaviors. 2014 , 2, 93		121
63	Study protocol: Asking QUESIONS about Alcohol in pregnancy (AQUA): a longitudinal cohort study of fetal effects of low to moderate alcohol exposure. 2014 , 14, 302		20
62	Fatty acid ethyl esters in hair: correlation with self-reported ethanol intake in 160 subjects and influence of estroprogestin therapy. <i>Drug Testing and Analysis</i> , 2014 , 6, 930-5	3.5	10

61	Proteomic approaches and identification of novel therapeutic targets for alcoholism. 2014 , 39, 104-30		28
60	Meconium indicators of maternal alcohol abuse during pregnancy and association with patient characteristics. 2014 , 2014, 702848		21
59	Influence of thermal hair straightening on ethyl glucuronide content in hair. <i>Drug Testing and Analysis</i> , 2014 , 6 Suppl 1, 74-7	3.5	32
58	Examination of sex differences in fatty acid ethyl ester and ethyl glucuronide hair analysis. <i>Drug Testing and Analysis</i> , 2014 , 6 Suppl 1, 30-6	3.5	11
57	Determination of ethyl glucuronide in hair improves evaluation of long-term alcohol abstinence in liver transplant candidates. 2014 , 34, 469-76		39
56	Sensitivity and specificity of EtG in hair as a marker of chronic excessive drinking: pooled analysis of raw data and meta-analysis of diagnostic accuracy studies. <i>Therapeutic Drug Monitoring</i> , 2014 , 36, 560-73 ^{3,2}		29
55	[What ethanol metabolites as biological markers tell us about alcohol use]. 2014 , 164, 25-33		4
54	An evaluation of washing and extraction techniques in the analysis of ethyl glucuronide and fatty acid ethyl esters from hair samples. 2014 , 953-954, 115-9		11
53	A SPME-GC/MS procedure for the determination of fatty acid ethyl esters in hair for confirmation of abstinence test results. 2014 , 52, 955-60		9
52	Fetal and perinatal exposure to drugs and chemicals: novel biomarkers of risk. 2014 , 54, 295-315		16
51	Validation of a novel method to identify in utero ethanol exposure: simultaneous meconium extraction of fatty acid ethyl esters, ethyl glucuronide, and ethyl sulfate followed by LC-MS/MS quantification. 2014 , 406, 1945-55		29
50	Alcohol-medical drug interactions. 2014 , 125, 543-59		13
49	Hair ethyl glucuronide levels as a marker for alcohol use and abuse: a review of the current state of the art. 2014 , 134, 1-11		97
48	Analysis of ethyl glucuronide and ethyl sulfate using aqueous normal-phase chromatography with mass spectrometry. 2015 , 38, 1515-20		5
47	Ethanol metabolites: their role in the assessment of alcohol intake. 2015 , 39, 2060-72		78
46	Interpretation of group-level factors from a large population dataset in the determination of ethyl glucuronide in hair. <i>Drug Testing and Analysis</i> , 2015 , 7, 407-13	3.5	12
45	Direct Alcohol Biomarkers Ethyl Glucuronide, Ethyl Sulfate, Fatty Acid Ethyl Esters, and Phosphatidylethanol. 2015 , 181-220		3
44	Maternal and neonatal hair and breast milk in the assessment of perinatal exposure to drugs of abuse. 2015 , 7, 1273-97		20

43	Alcohol Biomarkers in Hair. 2015 , 71-139		6
42	Determination of direct alcohol markers: a review. 2015 , 407, 4907-25		55
41	Biomarkers of alcohol misuse: recent advances and future prospects. 2016 , 11, 78-89		18
40	Biomarker-Based Approaches for Assessing Alcohol Use Disorders. 2016 , 13, 166		61
39	Integration of Clinical Examination, Self-Report, and Hair Ethyl Glucuronide Analysis for Evaluation of Patients With Alcoholic Liver Disease Prior to Liver Transplantation. 2016 , 26, 40-6		5
38	Nonoxidative ethanol metabolism in humans-from biomarkers to bioactive lipids. 2016 , 68, 916-923		29
37	A novel, simultaneous extraction of FAEE and EtG from meconium and analysis by LC-MS/MS. 2016 , 408, 2587-94		8
36	Alternative sampling strategies for the assessment of alcohol intake of living persons. 2016 , 49, 1078-91		29
35	Levels of Hair Ethyl Glucuronide in Patients with Decreased Kidney Function: Possibility of Misclassification of Social Drinkers. 2016 , 40, 451-6		12
34	Segmental hair analysis to assess effectiveness of single-session motivational intervention to stop ethanol use during pregnancy. 2016 , 158, 45-51		8
33	Fetal exposure to ethanol: relationship between ethyl glucuronide in maternal hair during pregnancy and ethyl glucuronide in neonatal meconium. 2016 , 54, 427-35		27
32	Evaluation of direct and indirect ethanol biomarkers using a likelihood ratio approach to identify chronic alcohol abusers for forensic purposes. <i>Forensic Science International</i> , 2017 , 271, 13-22	2.6	18
31	Distribution pattern of ethyl glucuronide and caffeine concentrations over the scalp of a single person in a forensic context. <i>Drug Testing and Analysis</i> , 2017 , 9, 1594-1603	3.5	14
30	Substance Use in Pregnancy: Managing the High-Risk Patient. <i>Journal of Obstetrics and Gynaecology Canada</i> , 2017 , 39, 837-838	1.3	
29	Consommation de substances psychoactives pendant la grossesse : Prise en charge de la patiente à risque élevé <i>Journal of Obstetrics and Gynaecology Canada</i> , 2017 , 39, 839-840	1.3	
28	Commentary on current changes of the SoHT 2016 consensus on alcohol markers in hair and further background information. <i>Forensic Science International</i> , 2017 , 278, 326-333	2.6	23
27	Quantification of Phosphatidylethanol in Whole Blood as a Proxy for Chronic Alcohol Consumption, Using Ultra Performance Convergence Chromatography Tandem Mass Spectrometry. <i>Therapeutic Drug Monitoring</i> , 2018 , 40, 268-275	3.2	7
26	Influence of alcohol containing and alcohol free cosmetics on FAEE concentrations in hair. A performance evaluation of ethyl palmitate as sole marker, versus the sum of four FAEEs. <i>Forensic Science International</i> , 2018 , 283, 29-34	2.6	6

25	Development and validation of a Partial Least Squares-Discriminant Analysis (PLS-DA) model based on the determination of ethyl glucuronide (EtG) and fatty acid ethyl esters (FAEEs) in hair for the diagnosis of chronic alcohol abuse. <i>Forensic Science International</i> , 2018 , 282, 221-230	2.6	11
24	Ethyl glucuronide hair testing: A review. <i>Forensic Science International</i> , 2019 , 300, 106-119	2.6	23
23	Biomarkers of Alcohol Misuse. 2019 , 557-565		0
22	Aktuelles zur forensisch-toxikologischen Haaranalytik. <i>Rechtsmedizin</i> , 2019 , 29, 137-153	0.6	
21	Alcohol Biomarkers: Clinical Issues and Analytical Methods. 2019 , 25-42		3
20	Ethyl Glucuronide as a Long-term Alcohol Biomarker in Fingernail and Hair. Matrix Comparison and Evaluation of Gender Bias. <i>Alcohol and Alcoholism</i> , 2019 , 54, 402-407	3.5	11
19	Letter to the Editor-Hair Ethanol Glucuronide (EtG) and Washout Effects: Can and Should Pre-Analytical Washout Effects be Assessed During the Method Validation Phase?. <i>Journal of Forensic Sciences</i> , 2019 , 64, 323-324	1.8	
18	The Clinical Presentations. <i>SpringerBriefs in Well-being and Quality of Life Research</i> , 2019 , 13-41	0.1	
17	Improved measurement of ethyl glucuronide concentrations in hair using UPLCMS/MS for the evaluation of chronic ethanol consumption. <i>Forensic Science International</i> , 2020 , 306, 110071	2.6	2
16	A Systematic Review of Solid-Phase Microextraction Applications in the Forensic Context. <i>Journal of Analytical Toxicology</i> , 2020 , 44, 268-297	2.9	4
15	Determination of ethyl glucuronide in human hair samples: Decontamination vs extraction. <i>Drug Testing and Analysis</i> , 2020 , 12, 948-956	3.5	3
14	Validation of hair ethyl glucuronide using transdermal monitoring and self-reported alcohol use in women of childbearing potential. <i>Neuropsychopharmacology Reports</i> , 2021 , 41, 144-151	2.2	2
13	Les marqueurs de l'Alcoolisme chronique. Focus sur les approches immuno-chimiques. <i>Toxicologie Analytique Et Clinique</i> , 2009 , 21, 21-25	0.4	12
12	The fatty acid ethyl esters (FAEE) hair test: emerging technology for the diagnosis of fetal alcohol spectrum disorders (FASD). <i>Toxicologie Analytique Et Clinique</i> , 2009 , 21, 61-65	0.4	5
11	Testing for alcohol use in hair: is ethyl glucuronide (EtG) stable in hair?. <i>Toxicologie Analytique Et Clinique</i> , 2009 , 21, 67-71	0.4	20
10	Gas chromatography tandem mass spectrometry for biomarkers of alcohol abuse in human hair. <i>Therapeutic Drug Monitoring</i> , 2010 , 32, 216-23	3.2	15
9	The Diagnosis of Alcoholism Through the Identification of Biochemical Markers in Hair. 2009 , 151-169		
8	Alcohols and Aldehydes. 2010 , 291-306		

7	Slate and Trait Markers of Alcohol Abuse. 2012 , 47-92		
6	Biological State Marker for Alcohol Consumption. 2015 , 261-292		
5	The role of (bio)markers in hair analysis. 2015 , 120-134		
4	Fetal alcohol spectrum disorders: research challenges and opportunities. <i>Alcohol Research</i> , 2011 , 34, 4-14		26
3	Focus on: biomarkers of fetal alcohol exposure and fetal alcohol effects. <i>Alcohol Research</i> , 2011 , 34, 56-63		29
2	Alcohol and Head and Neck Cancer: Updates on the Role of Oxidative Stress, Genetic, Epigenetics, Oral Microbiota, Antioxidants, and Alkylating Agents.. <i>Antioxidants</i> , 2022 , 11,	7.1	3
1	Old and New Biomarkers of Alcohol Abuse: Narrative Review. 2023 , 12, 2124		0