

CITATION REPORT

List of articles citing

Revised equations for estimated GFR from serum creatinine in Japan

DOI: 10.1053/j.ajkd.2008.12.034

American Journal of Kidney Diseases, 2009, 53, 982-92.

Source: <https://exaly.com/paper-pdf/46024405/citation-report.pdf>

Version: 2024-04-25

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
2291	Challenges of Interpreting Results From Analyses That Use Extensive Claims Data. 2009 , 151, 895		
2290	A scoring system to predict renal outcome in IgA nephropathy: a nationwide 10-year prospective cohort study. 2009 , 24, 3068-74		164
2289	Determinants of decline in glomerular filtration rate in nonproteinuric subjects with or without diabetes and hypertension. 2009 , 4, 1432-40		59
2288	Retraction. Combination treatment of angiotensin-II receptor blocker and angiotensin-converting-enzyme inhibitor in non-diabetic renal disease (COOPERATE): a randomised controlled trial. 2009 , 151, 897		1
2287	Impacts of changes in obesity parameters for the prediction of blood pressure change in Japanese individuals. 2009 , 32, 421-7		9
2286	Prognostic significance of home blood pressure control on renal and cardiovascular outcomes in elderly patients with chronic kidney disease. 2009 , 32, 1123-9		20
2285	Association of gene polymorphisms with chronic kidney disease in Japanese individuals. 2009 , 24, 539-47		13
2284	Impact of reduced renal function on prognosis in Japanese patients with coronary artery disease: a prospective cohort of Shinken Database 2007. 2009 , 32, 920-6		13
2283	Reduction in urinary excretion of neutrophil gelatinase-associated lipocalin by angiotensin receptor blockers in hypertensive patients. 2009 , 24, 2608-9; author reply 2609-10		15
2282	Angiotensin-II receptor antagonist combined with calcium channel blocker or diuretic for essential hypertension. 2009 , 32, 962-8		51
2281	Reduction of circulating soluble fms-like tyrosine kinase-1 plays a significant role in renal dysfunction-associated aggravation of atherosclerosis. 2009 , 120, 2470-7		44
2280	Higher levels of urinary albumin excretion within the normal range predict faster decline in glomerular filtration rate in diabetic patients. 2009 , 32, 1518-20		51
2279	Contrast medium volume to estimated glomerular filtration rate ratio as a predictor of contrast-induced nephropathy developing after elective percutaneous coronary intervention. 2009 , 54, 214-20		26
2278	Plasma granzyme B as a predicting factor of coronary artery disease--clinical significance in patients with chronic renal failure. 2009 , 54, 409-15		13
2277	Antiproteinuric and blood pressure-lowering effects of a fixed-dose combination of losartan and hydrochlorothiazide in hypertensive patients with stage 3 chronic kidney disease. 2009 , 29, 1061-72		8
2276	Comparison of blood glucose values on admission for acute myocardial infarction in patients with versus without diabetes mellitus. 2009 , 104, 769-74		43
2275	GFR estimation in Japan and China: what accounts for the difference?. <i>American Journal of Kidney Diseases</i> , 2009 , 53, 932-5	7.4	73

2274	Effect of a carbonaceous oral adsorbent on the progression of CKD: a multicenter, randomized, controlled trial. <i>American Journal of Kidney Diseases</i> , 2009 , 54, 459-67	7.4	124
2273	Negative effects of anemia on quality of life and its improvement by complete correction of anemia by administration of recombinant human erythropoietin in posttransplant patients. 2009 , 13, 355-360		13
2272	Prevalence of anemia according to stage of chronic kidney disease in a large screening cohort of Japanese. 2009 , 13, 614-20		29
2271	Prevalence of chronic kidney disease in the Japanese general population. 2009 , 13, 621-30		362
2270	Evidence-based practice guideline for the treatment of CKD. 2009 , 13, 537-66		95
2269	Low levels of plasma soluble receptor for advanced glycation end products are associated with severe leukoaraiosis in acute stroke patients. 2009 , 287, 41-4		27
2268	Association of urinary albumin excretion with insulin resistance in Japanese subjects: impact of gender difference on insulin resistance. 2009 , 48, 1621-7		5
2267	Significance of urinary liver-fatty acid-binding protein in cardiac catheterization in patients with coronary artery disease. 2009 , 48, 1731-7		24
2266	Association of metabolic syndrome with urinary albumin excretion, low-grade inflammation, and low glomerular filtration rate among non-diabetic Japanese subjects. 2009 , 48, 1855-62		4
2265	Did Akhenaten have the Antley-Bixler syndrome?. 2009 , 151, 892; author reply 892		2
2264	The CKD-EPI equation and MDRD study equation find similar prevalence of chronic kidney disease in Asian populations. 2009 , 151, 892-3; author reply 893		20
2263	Challenges of interpreting results from analyses that use extensive claims data. 2009 , 151, 895; author reply 895		
2262	A pain in the neck: probiotics for ulcerative colitis. 2009 , 151, 895-7		10
2261	Relevance of molecular markers in prostate cancer. 2009 , 151, 894; author reply 894-5		
2260	Efficacy and safety of tacrolimus for lupus nephritis: a placebo-controlled double-blind multicenter study. 2009 , 19, 606-615		86
2259	Anemia is an independent predictor of long-term adverse outcomes in patients hospitalized with heart failure in Japan. A report from the Japanese Cardiac Registry of Heart Failure in Cardiology (JCARE-CARD). 2009 , 73, 1901-8		67
2258	Chronic kidney disease as an independent risk for long-term adverse outcomes in patients hospitalized with heart failure in Japan. Report from the Japanese Cardiac Registry of Heart Failure in Cardiology (JCARE-CARD). 2009 , 73, 1442-7		70
2257	Chronic kidney disease as an independent risk factor for new-onset atrial fibrillation in hypertensive patients. 2010 , 28, 1738-44		87

2256	Advances in glomerular filtration rate-estimating equations. 2010 , 19, 298-307	47
2255	Current Opinion in Nephrology and Hypertension. Current world literature. 2010 , 19, 315-32	1
2254	Estimated glomerular filtration rate is superior to measured creatinine clearance for predicting postoperative renal dysfunction in patients undergoing pancreatoduodenectomy. 2010 , 39, 20-5	9
2253	Crucial role of kidney function in resistance to antihypertensive therapy in patients with diabetes mellitus. 2010 , 28, 2323-8	10
2252	Relation between the estimated glomerular filtration rate and pulse wave velocity in Japanese. 2010 , 49, 1315-20	7
2251	Influence of low-grade inflammation on plasma B-type natriuretic peptide levels. 2010 , 49, 2659-68	10
2250	4. Evidence-based Clinical Practice Guidelines for CKD (2009).. 2010 , 99, 2984-2989	
2249	3. Up-to-date Clinical Practice for CKD.. 2010 , 99, 559-563	
2248	[Determination of glomerular filtration rate (GFR) in patients with chronic kidney diseases]. 2010 , 99, 172-6	
2247	Effects of irbesartan on 24-h blood pressure changes and urinary albumin levels in Japanese outpatients. 2010 , 1, 669-673	
2246	Genetic risk for myocardial infarction in Japanese individuals with or without chronic kidney disease. 2010 , 25, 743-9	3
2245	Correlation between estimated glomerular filtration rate (eGFR) and apparent diffusion coefficient (ADC) values of the kidneys. 2010 , 9, 59-64	33
2244	Association of genetic variants with chronic kidney disease in Japanese individuals with or without hypertension or diabetes mellitus. 2010 , 1, 137-145	8
2243	Relationship between renal dysfunction and severity of coronary artery disease in Japanese patients. 2010 , 74, 786-91	30
2242	Plasma B-type natriuretic peptide level and cardiovascular events in chronic kidney disease in a community-based population. 2010 , 74, 792-7	24
2241	Characteristics of revascularization treatment for arteriosclerosis obliterans in patients with and without hemodialysis. 2010 , 74, 2426-33	11
2240	Plasma cystatin C concentration reflects the severity of coronary artery disease in patients without chronic kidney disease. 2010 , 74, 2441-7	22
2239	Impact of left ventricular hypertrophy on the time-course of renal function in hypertensive patients [a subanalysis of the CASE-J trial] 2010 , 74, 2132-8	3

2238	Preoperative renal dysfunction and mortality after off-pump coronary artery bypass grafting in Japanese. 2010 , 74, 1866-72	16
2237	The administration of an active vitamin D(3) analogue reduced the serum concentrations of 1-84 and truncated parathyroid hormone in pseudohypoparathyroidism type Ib patients. 2010 , 57, 609-14	4
2236	Associations of metabolic factors, especially serum retinol-binding protein 4 (RBP4), with blood pressure in Japanese--the Tanno and Sobetsu study. 2010 , 57, 811-7	22
2235	A Cross-Sectional Study on the Effects of Long Term Very Low Protein Diets in Patients with Chronic Kidney Disease. 2010 , 7, 7-13	5
2234	[Prevalence of kidney disease in HIV-infected patients in Japan--A single center study]. 2010 , 84, 28-32	2
2233	The levels of somatostatin receptors in causative tumors of oncogenic osteomalacia are insufficient for their agonist to normalize serum phosphate levels. 2010 , 86, 455-62	12
2232	Utility of the Japanese GFR estimation equation for evaluating potential donor kidney function. 2010 , 14, 63-7	22
2231	Very low but stable glomerular filtration rate after living kidney donation: is the concept of "chronic kidney disease" applicable to kidney donors?. 2010 , 14, 356-62	15
2230	Different clinical outcomes for cardiovascular events and mortality in chronic kidney disease according to underlying renal disease: the Gonryo study. 2010 , 14, 333-9	44
2229	Measurement of health-related quality of life in patients with chronic kidney disease in Japan with EuroQol (EQ-5D). 2010 , 14, 340-8	41
2228	Incidence of symptomatic stroke and cancer in chronic kidney disease patients treated with epoetins. 2010 , 14, 445-52	14
2227	Discrepancy between cystatin C and creatinine points leading to a diagnosis of postrenal acute kidney injury and its reversibility: three case reports. 2010 , 14, 608-13	7
2226	Chronic Kidney Disease Japan Cohort study: baseline characteristics and factors associated with causative diseases and renal function. 2010 , 14, 558-70	85
2225	Predictive value of serum cystatin C, β_2 -microglobulin, and urinary liver-type fatty acid-binding protein on the development of contrast-induced nephropathy. 2010 , 25, 85-90	20
2224	Leptin mediates the relationship between abdominal obesity and microalbuminuria in type 2 diabetic patients. 2010 , 1, 42-48	2
2223	Investigation of clinical factors associated with development of chronic kidney disease and eGFR decline in Japanese patients with type 2 diabetes mellitus. 2010 , 1, 65-71	
2222	Iodine-123-metaiodobenzylguanidine imaging can predict future cardiac events in heart failure patients with preserved ejection fraction. 2010 , 24, 679-86	36
2221	ASCI 2010 contrast media guideline for cardiac imaging: a report of the Asian Society of Cardiovascular Imaging cardiac computed tomography and cardiac magnetic resonance imaging guideline working group. 2010 , 26, 203-12	10

2220	Association of kidney function with coronary atherosclerosis and calcification in autopsy samples from Japanese elders: the Hisayama study. <i>American Journal of Kidney Diseases</i> , 2010 , 55, 21-30	7.4	137
2219	Estimating GFR using the CKD Epidemiology Collaboration (CKD-EPI) creatinine equation: more accurate GFR estimates, lower CKD prevalence estimates, and better risk predictions. <i>American Journal of Kidney Diseases</i> , 2010 , 55, 622-7	7.4	551
2218	Modification of the CKD epidemiology collaboration (CKD-EPI) equation for Japanese: accuracy and use for population estimates. <i>American Journal of Kidney Diseases</i> , 2010 , 56, 32-8	7.4	287
2217	A case of Churg-Strauss syndrome with necrotizing crescentic glomerulonephritis accompanied by acute coronary syndrome due to vasospasm. <i>American Journal of Kidney Diseases</i> , 2010 , 56, e5-9	7.4	7
2216	Silent brain infarction and rapid decline of kidney function in patients with CKD: a prospective cohort study. <i>American Journal of Kidney Diseases</i> , 2010 , 56, 468-76	7.4	23
2215	Difference in risk factors between acute coronary syndrome and stable angina pectoris in the Japanese: smoking as a crucial risk factor of acute coronary syndrome. 2010 , 55, 345-53		25
2214	Low-density lipoprotein cholesterol to high-density lipoprotein cholesterol ratio as a useful marker for early-stage carotid atherosclerosis. 2010 , 59, 653-7		25
2213	Annual incidence of atrial fibrillation and related factors in adults. 2010 , 106, 1129-33		38
2212	Relation of smoking status to serum levels of N-terminal pro-brain natriuretic peptide in middle-aged men without overt cardiovascular disease. 2010 , 106, 1456-60		22
2211	Efficacy of bilateral internal thoracic artery grafting in patients with chronic kidney disease. 2010 , 89, 1106-11		20
2210	Differential expression of glycogenes in tonsillar B lymphocytes in association with proteinuria and renal dysfunction in IgA nephropathy. 2010 , 136, 447-55		35
2209	Persistent glomerular hematuria in living kidney donors confers a risk of progressive kidney disease in donors after heminephrectomy. 2010 , 10, 1597-604		33
2208	Morning hypertension assessed by home monitoring is a strong predictor of concentric left ventricular hypertrophy in patients with untreated hypertension. 2010 , 12, 776-83		20
2207	Metabolic impact on serum levels of complement component 3 in Japanese patients. 2010 , 24, 113-8		15
2206	Dietary salt reduction in rural patients with albuminuria using family and community support: the Mima study. 2010 , 9, 6		2
2205	Serum C-reactive protein and thioredoxin levels in subjects with mildly reduced glomerular filtration rate. 2010 , 11, 7		19
2204	The estimated GFR, but not the stage of diabetic nephropathy graded by the urinary albumin excretion, is associated with the carotid intima-media thickness in patients with type 2 diabetes mellitus: a cross-sectional study. 2010 , 9, 18		51
2203	Angiogenesis and chronic kidney disease. 2010 , 3, 13		51

2202	Comparison of sustained hemodiafiltration with continuous venovenous hemodiafiltration for the treatment of critically ill patients with acute kidney injury. 2010 , 34, 331-8	47
2201	Prognostic study of cardiac and renal events in Japanese patients with chronic kidney disease and cardiovascular risk using myocardial perfusion SPECT: J-ACCESS 3 study design. 2010 , 14, 379-85	10
2200	Evaluation of blood purification and bortezomib plus dexamethasone therapy for the treatment of acute renal failure due to myeloma cast nephropathy. 2010 , 14, 451-6	9
2199	2009 Japanese Society for Dialysis Therapy guidelines for peritoneal dialysis. 2010 , 14, 489-504	49
2198	Prospective evaluation of pharmacokinetically guided dosing of carboplatin in Japanese patients with cancer. 2010 , 101, 2601-5	16
2197	Relationship between chronic kidney disease and silent cerebral infarction in patients with Type 2 diabetes. 2010 , 27, 538-43	13
2196	Comparison of the prevalence of chronic kidney disease in Japanese patients with Type 1 and Type 2 diabetes. 2010 , 27, 1017-23	27
2195	Renal outcomes in chronic kidney disease. 2010 , 15 Suppl 2, 27-30	14
2194	Current status of estimated glomerular filtration rate (eGFR) equations for Asians and an approach to create a common eGFR equation. 2010 , 15 Suppl 2, 45-8	22
2193	Poster Abstracts. 2010 , 15, 62-144	2
2192	Nestin is a novel marker for renal tubulointerstitial injury in immunoglobulin A nephropathy. 2010 , 15, 568-74	19
2191	Serum fetuin-A is an independent marker of insulin resistance in Japanese men. 2010 , 17, 925-33	55
2190	[Significance of serum creatinine measurement in the specific screening test system for detection of chronic kidney disease]. 2010 , 101, 25-8	1
2189	Efficacy of combination therapy with telmisartan plus amlodipine in patients with poorly controlled hypertension. 2010 , 3, 342-6	5
2188	Effects of pitavastatin (LIVALO tablet) on the estimated glomerular filtration rate (eGFR) in hypercholesterolemic patients with chronic kidney disease. Sub-analysis of the LIVALO Effectiveness and Safety (LIVES) Study. 2010 , 17, 601-9	48
2187	Impact of statin therapy on renal function and long-term prognosis in acute coronary syndrome patients with chronic kidney disease. 2010 , 51, 312-8	8
2186	Ethnic coefficients for glomerular filtration rate estimation by the Modification of Diet in Renal Disease study equations in the Korean population. 2010 , 25, 1616-25	88
2185	Brachial-ankle pulse wave velocity as an index of central arterial stiffness. 2010 , 17, 658-65	116

2184	Glomerular density in renal biopsy specimens predicts the long-term prognosis of IgA nephropathy. 2010 , 5, 39-44	51
2183	Impact of long-acting calcium channel blockers on the prognosis of patients with coronary artery disease with and without chronic kidney disease: a comparison of three drugs. 2010 , 38, 253-65	2
2182	Association between urinary type IV collagen level and deterioration of renal function in type 2 diabetic patients without overt proteinuria. 2010 , 33, 1805-10	52
2181	Cerebral microvascular disease predicts renal failure in type 2 diabetes. 2010 , 21, 520-6	40
2180	The relation of coffee consumption to serum uric Acid in Japanese men and women aged 49-76 years. 2010 , 2010,	40
2179	Involvement of the basilar artery in diabetes mellitus: an MRI study of brainstem infarctions. 2010 , 64, 230-5	11
2178	Estimating the contribution of genes to variation in renal drug clearance by active secretion using multiple data from clinical phase I studies. 2010 , 50, 109-14	
2177	Comparison between the antiproteinuric effects of the calcium channel blockers benidipine and cilnidipine in combination with angiotensin receptor blockers in hypertensive patients with chronic kidney disease. 2010 , 19, 1027-37	11
2176	Impact of age and overt proteinuria on outcomes of stage 3 to 5 chronic kidney disease in a referred cohort. 2010 , 5, 1558-65	48
2175	High frequencies of diabetic micro- and macroangiopathies in patients with type 2 diabetes mellitus with decreased estimated glomerular filtration rate and normoalbuminuria. 2010 , 25, 1161-7	36
2174	Prognostic factors for progression of early- and late-stage calcific aortic valve disease in Japanese: the Japanese Aortic Stenosis Study (JASS) Retrospective Analysis. 2010 , 33, 269-74	26
2173	Association of serum cystatin C with pulse wave velocity, but not pressure wave reflection, in subjects with normal renal function or mild chronic kidney disease. 2010 , 23, 967-73	12
2172	Chronic kidney disease in long-term survivors of myeloablative allogeneic haematopoietic cell transplantation: prevalence and risk factors. 2010 , 25, 278-82	67
2171	Prehypertension subtype with elevated C-reactive protein: risk of ischemic stroke in a general Japanese population. 2010 , 23, 1108-13	23
2170	Impact of the clinical conditions at dialysis initiation on mortality in incident haemodialysis patients: a national cohort study in Taiwan. 2010 , 25, 2616-24	121
2169	Changes in waist circumference and body mass index in relation to changes in serum uric acid in Japanese individuals. 2010 , 37, 410-6	44
2168	A comparative assessment of the RIFLE, AKIN and conventional criteria for acute kidney injury after hematopoietic SCT. 2010 , 45, 1427-34	52
2167	Assessment and significance of abdominal aortic calcification in chronic kidney disease. 2010 , 25, 1888-95	37

2166	Stage of chronic kidney disease is an outcome-predicting factor of angioplasty for atheromatous renal artery stenosis. 2010 , 33, 1206-10	8
2165	Impact of renal function on cardiovascular events in elderly hypertensive patients treated with efonidipine. 2010 , 33, 1211-20	32
2164	Marked cerebral atrophy is correlated with kidney dysfunction in nondisabled adults. 2010 , 33, 1232-7	30
2163	Association of albuminuria and reduced estimated glomerular filtration rate with incident stroke and coronary artery disease in patients with type 2 diabetes. 2010 , 33, 1298-304	35
2162	The novel and independent association between single-point SNP of NPHP4 gene and renal function in non-diabetic Japanese population: the Takahata study. 2010 , 55, 791-5	7
2161	Mild renal dysfunction is a risk factor for a decrease in bone mineral density and vertebral fractures in Japanese postmenopausal women. 2010 , 95, 4635-42	44
2160	Brachial-ankle pulse wave velocity as a risk stratification index for the short-term prognosis of type 2 diabetic patients with coronary artery disease. 2010 , 33, 1018-24	69
2159	Rationale and design of the Eplerenone combination Versus conventional Agents to Lower blood pressure on Urinary Antialbuminuric Treatment Effect (EVALUATE) trial: a double-blinded randomized placebo-controlled trial to evaluate the antialbuminuric effects of an aldosterone blocker in hypertensive patients with albuminuria. 2010 , 33, 616-21	23
2158	Relationship between chronic kidney disease and sleep blood pressure in patients with sleep apnea syndrome. 2010 , 33, 1278-82	9
2157	High level of plasma endothelin-1 predicts development of hypertension in normotensive subjects. 2010 , 23, 1103-7	11
2156	Expression of angiotensin II type 1 receptor-interacting molecule in normal human kidney and IgA nephropathy. 2010 , 299, F720-31	21
2155	Chronic kidney disease and end-stage renal disease in the elderly population: current prevalence, future projections, and clinical significance. 2010 , 17, 293-301	153
2154	Prevalence and risk factors for renal artery stenosis and chronic kidney disease in Japanese patients with peripheral arterial disease. 2010 , 33, 911-5	16
2153	C-reactive protein is associated with cigarette smoking-induced hyperfiltration and proteinuria in an apparently healthy population. 2010 , 33, 1129-36	22
2152	Metabolomic profiling of uremic solutes in CKD patients. 2010 , 33, 944-52	95
2151	Continuous low-dose infusion of human atrial natriuretic peptide in patients with left ventricular dysfunction undergoing coronary artery bypass grafting: the NU-HIT (Nihon University working group study of low-dose Human ANP Infusion Therapy during cardiac surgery) for left ventricular dysfunction. 2010 , 55, 1844-51	61
2150	The influence of the obesity paradox and chronic kidney disease on long-term survival in a Japanese cohort with peripheral arterial disease. 2010 , 52, 110-7	63
2149	Silent cerebral infarction is associated with the development and progression of nephropathy in patients with type 2 diabetes. 2010 , 33, 1000-3	7

2148	Long-term effect of metformin on blood glucose control in non-obese patients with type 2 diabetes mellitus. 2010 , 7, 83	24
2147	Diabetic Nephropathy Remission and Regression Team Trial in Japan (DNETT-Japan): Rationale and study design. 2010 , 87, 228-32	5
2146	New insights in glomerular filtration rate formulas and chronic kidney disease classification. 2010 , 411, 1341-7	13
2145	Is estimated donor glomerular filtration rate before death a better predictor of decreased-donor kidney function?. 2010 , 42, 3938-40	1
2144	Serum levels and urinary excretion of salusin-alpha in renal insufficiency. 2010 , 162, 129-32	21
2143	Association between high-sensitivity cardiac troponin T levels and the predicted cardiovascular risk in middle-aged men without overt cardiovascular disease. 2010 , 159, 972-8	82
2142	A prediction model for the risk of incident chronic kidney disease. 2010 , 123, 836-846.e2	64
2141	Insufficient control of morning home blood pressure in Japanese patients with hypertension associated with diabetes mellitus. 2010 , 1, 266-72	1
2140	Mild anemia is frequent and associated with micro- and macroangiopathies in patients with type 2 diabetes mellitus. 2010 , 1, 273-8	13
2139	Renal dysfunction impairs circadian variation of endothelial function in patients with essential hypertension. 2010 , 4, 265-71	4
2138	Pentraxin 3 in unstable angina and non-ST-segment elevation myocardial infarction. 2010 , 210, 220-5	75
2137	Central versus peripheral arterial stiffness in association with coronary, cerebral and peripheral arterial disease. 2010 , 211, 480-5	72
2136	Arterial stiffness and declines in individuals with normal renal function/early chronic kidney disease. 2010 , 212, 345-50	81
2135	Chronic kidney disease is associated with increased carotid artery stiffness without morphological changes in participants of health check-up programs. 2010 , 213, 306-10	9
2134	Early recognition and prevention of chronic kidney disease. 2010 , 375, 1296-309	368
2133	Usefulness of assessing masked and white-coat hypertension by ambulatory blood pressure monitoring for determining prevalent risk of chronic kidney disease: the Ohasama study. 2010 , 33, 1192-8	34
2132	Cystatin C is a sensitive marker for detecting a reduced glomerular filtration rate when assessing chronic kidney disease in patients with rheumatoid arthritis and secondary amyloidosis. 2010 , 39, 33-7	14
2131	Implication of CKD-EPI equation to estimate glomerular filtration rate in Chinese patients with chronic kidney disease. 2011 , 33, 859-65	16

2130	Kidney function and histological damage in autopsy subjects with myocardial infarction. 2011 , 33, 847-52	7
2129	Usefulness of the resistive index in renal Doppler ultrasonography as an indicator of vascular damage in patients with risks of atherosclerosis. 2011 , 26, 3256-62	45
2128	Indexing of renal function parameters by body surface area: intelligence or folly?. 2011 , 119, c289-92	34
2127	Beneficial effect of switching from a combination of angiotensin II receptor blockers other than losartan and thiazides to a fixed dose of losartan/hydrochlorothiazide on uric acid metabolism in hypertensive patients. 2011 , 33, 565-70	7
2126	Benefits of staple food restriction for Japanese obese patients with chronic kidney disease: a pilot study. 2011 , 21, 340-6	
2125	Results of low-dose human atrial natriuretic peptide infusion in nondialysis patients with chronic kidney disease undergoing coronary artery bypass grafting: the NU-HIT (Nihon University working group study of low-dose HANP Infusion Therapy during cardiac surgery) trial for CKD. 2011 , 58, 897-903	81
2124	Additive antioxidative effects of azelnidipine on angiotensin receptor blocker olmesartan treatment for type 2 diabetic patients with albuminuria. 2011 , 34, 935-41	25
2123	Olmesartan improves endothelial function in hypertensive patients: link with extracellular superoxide dismutase. 2011 , 34, 686-92	28
2122	Reappraisal of proteinuria and estimated GFR to predict progression to ESRD or death for hospitalized chronic kidney disease patients. 2011 , 33, 31-9	6
2121	Decreased plasma B-type natriuretic peptide levels in obesity are not explained by altered left ventricular hemodynamics. 2011 , 5, e267-360	3
2120	High level of plasma adiponectin in acute stroke patients is associated with stroke mortality. 2011 , 304, 102-6	25
2119	Increases in urinary albumin and beta2-microglobulin are independently associated with blood pressure in the Japanese general population: the Takahata Study. 2011 , 34, 831-5	15
2118	Glomerular filtration rate and 10-year mortality in a 70-year-old community-dwelling Japanese population. 2011 , 23, 223-30	
2117	Urinary type IV collagen in nondiabetic kidney disease. 2011 , 117, c160-6	10
2116	Glyoxalase I retards renal senescence. 2011 , 179, 2810-21	36
2115	Reference intervals for serum creatinine with enzymatic assay and evaluation of four equations to estimate glomerular filtration rate in a healthy Chinese adult population. 2011 , 412, 1793-7	13
2114	Association between serum soluble TNF receptors and renal dysfunction in type 2 diabetic patients without proteinuria. 2011 , 92, 174-80	23
2113	MR imaging of the biliary tract with Gd-EOB-DTPA: effect of liver function on signal intensity. 2011 , 77, 325-9	62

2112	Relations of blood inflammatory marker levels with cerebral microbleeds. 2011 , 42, 3202-6	71
2111	The association between post-load low insulin level and high mortality rates in patients with no previous diagnosis of diabetes mellitus. 2011 , 148, 59-63	
2110	Effect of edaravone on the estimated glomerular filtration rate in patients with acute ischemic stroke and chronic kidney disease. 2011 , 20, 111-6	6
2109	Cerebral microbleeds and dilative remodeling of the basilar artery: a magnetic resonance imaging study. 2011 , 20, 429-35	6
2108	I. Stroke in general. 2011 , 20, S7-30	21
2107	Reduced immune response to influenza A (H1N1) 2009 monovalent vaccine in HIV-infected Japanese subjects. 2011 , 29, 5694-8	15
2106	Severe obstructive sleep apnea increases cystatin C in clinically latent renal dysfunction. 2011 , 105, 643-9	29
2105	Glucose intolerance in renal transplant recipients is associated with increased urinary albumin excretion. 2011 , 24, 241-5	7
2104	Relationship between renal parenchymal volume and single kidney glomerular filtration rate before and after unilateral nephrectomy. 2011 , 77, 1404-8	30
2103	Renal hypothermia using ice-cold saline for retroperitoneal laparoscopic partial nephrectomy: evaluation of split renal function with technetium-99m-dimercaptosuccinic acid renal scintigraphy. 2011 , 77, 814-8	8
2102	Evaluation of the Chronic Kidney Disease Epidemiology Collaboration equation for estimating the glomerular filtration rate in multiple ethnicities. 2011 , 79, 555-62	288
2101	Comparison of effects of azelnidipine and trichlormethiazide in combination with olmesartan on blood pressure and metabolic parameters in hypertensive type 2 diabetic patients. 2011 , 2, 490-6	2
2100	Clinical evaluation of chronic nephrotoxicity of long-term cyclosporine A treatment in adult patients with steroid-dependent nephrotic syndrome. 2011 , 16, 319-25	5
2099	Validation study of oxford classification of IgA nephropathy: the significance of extracapillary proliferation. 2011 , 6, 2806-13	107
2098	Serum 1,5-anhydro-D-glucitol levels predict first-ever cardiovascular disease: an 11-year population-based cohort study in Japan, the Suita study. 2011 , 216, 477-83	53
2097	Sex-related differences in Japanese patients with peripheral arterial disease. 2011 , 219, 846-50	25
2096	Qualitative score of systemic arteriosclerosis by vascular ultrasonography as a predictor of coronary artery disease in type 2 diabetes. 2011 , 219, 623-9	14
2095	Association of a polymorphism of BTN2A1 with chronic kidney disease in individuals with or without hypertension or diabetes mellitus. 2011 , 2, 325-331	3

2094	Serum C-reactive protein levels and death and cardiovascular events in mild to moderate chronic kidney disease. 2011 , 52, 180-4	7
2093	Elevated cardiac troponin T predicts adverse outcomes in hypertensive patients. 2011 , 52, 164-9	21
2092	Effects of lipid-lowering therapy with rosuvastatin on kidney function and oxidative stress in patients with diabetic nephropathy. 2011 , 18, 1018-28	74
2091	Impact of mild to moderate renal dysfunction on left ventricular relaxation function and prognosis in ambulatory patients with nonischemic dilated cardiomyopathy. 2011 , 52, 366-71	5
2090	Utility of short-term variability of repolarization as a marker for monitoring a safe exercise training program in patients with cardiac diseases. 2011 , 52, 304-7	5
2089	Statin treatment decreased serum asymmetric dimethylarginine (ADMA) levels in ischemic stroke patients. 2011 , 18, 131-7	24
2088	Flow-mediated dilation is associated with microalbuminuria independent of cardiovascular risk factors in type 2 diabetes - interrelations with arterial thickness and stiffness. 2011 , 18, 744-52	24
2087	Prognostic factors affecting clinical outcomes after coronary artery bypass surgery: analysis of patients with chronic kidney disease after 5.9 years of follow-up. 2011 , 78, 156-65	6
2086	Effects of Six Functional SNPs on the Urinary 8-Isoprostane Level in a General Japanese Population; Shimane COHRE Study. 2011 , 30, 291-298	2
2085	Evidence for the immunosuppressive potential of calcineurin inhibitor-sparing regimens in liver transplant recipients with impaired renal function. 2011 , 2011, 483728	5
2084	Coffee consumption and cystatin-C-based estimated glomerular filtration rates in healthy young adults: results of a clinical trial. 2011 , 2011, 146865	19
2083	Gender-based outcomes among patients with diabetes mellitus after percutaneous coronary intervention in the drug-eluting stent era. 2011 , 52, 348-52	19
2082	Renal dysfunction in chronic hepatitis B patients treated with adefovir dipivoxil. 2011 , 52, 102-111	4
2081	Hyperuricemia is independently associated with coronary heart disease and renal dysfunction in patients with type 2 diabetes mellitus. 2011 , 6, e27817	70
2080	Factors increasing quantitative interstitial fibrosis from 0 hr to 1 year in living kidney transplant patients receiving tacrolimus. 2011 , 91, 78-85	20
2079	Evaluation of new acute kidney injury biomarkers in a mixed intensive care unit. 2011 , 39, 2464-9	134
2078	Chronic kidney disease in urolithiasis patients following successful extracorporeal shockwave lithotripsy. 2012 , 5, 3-6	2
2077	Effects of hyperacute blood pressure and heart rate on stroke outcomes after intravenous tissue plasminogen activator. 2011 , 29, 1980-7	21

2076	Effects of chronic kidney disease and post-angiographic acute kidney injury on long-term prognosis after coronary artery angiography. 2011 , 26, 1838-46	11
2075	Elevated serum uric acid predicts chronic kidney disease. 2011 , 342, 461-6	19
2074	Renal function and cardiovascular events during antihypertensive treatment in elderly patients. 2011 , 7, 59-63	
2073	Prevalence and prognostic implications of pre-diabetic state in patients with heart failure. 2011 , 75, 2833-9	39
2072	Effects of combination therapy with warfarin and bucolome for anticoagulation in patients with atrial fibrillation. 2011 , 75, 201-3	5
2071	Association of low glomerular filtration rate with the incidence of stroke in patients following complete coronary revascularization. 2011 , 75, 2372-8	5
2070	Quality of life as an independent predictor for cardiac events and death in patients with heart failure. 2011 , 75, 1661-9	61
2069	Elevated cystatin C levels predict the incidence of vasospastic angina. 2011 , 75, 2439-44	12
2068	Efficacy analysis of the lipid-lowering and renoprotective effects of rosuvastatin in patients with chronic kidney disease. 2011 , 58, 663-74	17
2067	Postprandial serum C-peptide to plasma glucose ratio as a predictor of subsequent insulin treatment in patients with type 2 diabetes. 2011 , 58, 315-22	62
2066	Influence of proteinuria on glycated albumin values in diabetic patients with chronic kidney disease. 2011 , 50, 23-9	58
2065	Association between metabolic syndrome and carotid atherosclerosis: relevance of combined criteria including the serum adiponectin level for the general population. 2011 , 50, 381-7	6
2064	Determinants of serum high molecular weight (HMW) adiponectin levels in patients with coronary artery disease: associations with cardio-renal-anemia syndrome. 2011 , 50, 2953-60	13
2063	Establishment of a method to detect microalbuminuria by measuring the total urinary protein-to-creatinine ratio in diabetic patients. 2011 , 225, 195-202	6
2062	Relationship between blood pressure control status and lifestyle in hypertensive outpatients. 2011 , 50, 2107-12	16
2061	Triple therapy with single daily dose of cyclosporine in Japanese patients with idiopathic membranous nephropathy. 2011 , 50, 1905-10	3
2060	Is mild renal dysfunction a risk factor for carotid atherosclerosis in apparently healthy adults?. 2011 , 50, 2285-9	7
2059	[108th Scientific Meeting of the Japanese Society of Internal Medicine: symposium: 3. The interaction between kidney and other organs; What should physicians know about it? (2) cardiovascular system and kidney]. 2011 , 100, 2537-43	

2058	Association between chronic kidney disease and small residual urine volumes in patients with benign prostatic hyperplasia. 2011 , 16, 335-9		11
2057	Management of asymptomatic hyperuricaemia in patients with chronic kidney disease by Japanese nephrologists: a questionnaire survey. 2011 , 16, 518-21		18
2056	Inaccuracies in estimated glomerular filtration rate in one Australian renal centre. 2011 , 16, 486-94		3
2055	Urinary angiotensin-converting enzyme 2 in patients with CKD. 2011 , 16, 567-72		43
2054	Association of a genetic variant of BTN2A1 with chronic kidney disease in Japanese individuals. 2011 , 16, 642-8		7
2053	Postoperative renal function after partial nephrectomy for renal cell carcinoma in patients with pre-existing chronic kidney disease: a comparison with radical nephrectomy. 2011 , 18, 472-6		21
2052	Mercaptoacetyltriglycine-3 renogram is not superior to estimated glomerular filtration rate measurement for the prediction of long-term renal function after nephrectomy. 2011 , 18, 570-4		2
2051	Renal parenchymal volume increases after contralateral nephrectomy: assessment using three-dimensional ultrasonography. 2011 , 18, 857-60		12
2050	Evidence for severe atherosclerotic changes in chronic hemodialysis patients: comparative autopsy study against cardiovascular disease patients without chronic kidney disease. 2011 , 15, 51-7		4
2049	Guideline-practice gap in the management of predialysis chronic kidney disease mineral bone disorder in Japan. 2011 , 15 Suppl 1, 2-8		12
2048	High serum concentrations of pentosidine, an advanced glycation end product, are associated with low normal value of ankle-brachial index in apparently healthy men. 2011 , 60, 649-54		8
2047	Nonalcoholic steatohepatitis and increased risk of chronic kidney disease. 2011 , 60, 735-9		108
2046	Chronic kidney disease and heart failure--Bidirectional close link and common therapeutic goal. 2011 , 57, 8-17		72
2045	Long-term outcomes of women with coronary artery disease following complete coronary revascularization. 2011 , 58, 158-64		5
2044	High-sensitivity cardiac troponin T in essential hypertension. 2011 , 58, 226-31		36
2043	GFR estimating equations in a multiethnic Asian population. <i>American Journal of Kidney Diseases</i> , 2011 , 58, 56-63	7.4	176
2042	Comparative evaluation of the Cockcroft-Gault Equation and the Modification of Diet in Renal Disease (MDRD) study equation for drug dosing: an opinion of the Nephrology Practice and Research Network of the American College of Clinical Pharmacy. 2011 , 31, 1130-44		69
2041	An early and simple predictor of severe left main and/or three-vessel disease in patients with non-ST-segment elevation acute coronary syndrome. 2011 , 107, 495-500		54

2040	Chronic kidney disease and CHADS(2) score independently predict cardiovascular events and mortality in patients with nonvalvular atrial fibrillation. 2011 , 107, 912-6	78
2039	Effect of left ventricular reverse remodeling on long-term prognosis after therapy with angiotensin-converting enzyme inhibitors or angiotensin II receptor blockers and β blockers in patients with idiopathic dilated cardiomyopathy. 2011 , 107, 1065-70	50
2038	Prevention of contrast-induced nephropathy by bolus injection of sodium bicarbonate in patients with chronic kidney disease undergoing emergent coronary procedures. 2011 , 107, 1163-7	35
2037	Management of non-ST-segment elevation acute myocardial infarction in patients with chronic kidney disease (from the Korea Acute Myocardial Infarction Registry). 2011 , 108, 206-13	21
2036	Relation of plasma levels of adiponectin to left ventricular diastolic dysfunction in patients undergoing cardiac catheterization for coronary artery disease. 2011 , 108, 1081-5	12
2035	Sex-specific threshold levels of plasma B-type natriuretic peptide for prediction of cardiovascular event risk in a Japanese population initially free of cardiovascular disease. 2011 , 108, 1564-9	11
2034	High white blood cell count and low estimated glomerular filtration rate are independently associated with serum level of monocyte chemoattractant protein-1 in a general population. 2011 , 34, 189-94	13
2033	Beneficial cardiometabolic actions of telmisartan plus amlodipine therapy in elderly patients with poorly controlled hypertension. 2011 , 34, 261-5	6
2032	Manifestations of transthyretin-related familial amyloidotic polyneuropathy: long-term follow-up of Japanese patients after liver transplantation. 2011 , 41, 1211-8	37
2031	A simple model for predicting incidence of chronic kidney disease in HIV-infected patients. 2011 , 15, 242-7	17
2030	Kidney transplantation recovers the reduction level of serum sulfatide in ESRD patients via processes correlated to oxidative stress and platelet count. 2011 , 28, 125-35	13
2029	Design and rationale of Japanese evaluation between Formula of Azelnidipine and amlodipine add on olmesartan to Get antialbuminuric effect study (J-FLAG) : evaluation of the antialbuminuric effects between calcium channel blocker with sympatholytic action in hypertensive patients with diabetes and albuminuria. 2011 , 25, 311-7	1
2028	Evaluation of contributing factors to restless legs syndrome in migraine patients. 2011 , 258, 2026-35	41
2027	Elevated serum levels of interleukin-18 in patients with overt diabetic nephropathy: effects of miglitol. 2011 , 15, 58-63	10
2026	Effects of renal dysfunction on cardiovascular events in diabetic patients with hypertension: challenge-DM Study subgroup analysis. 2011 , 15, 64-72	1
2025	Methylprednisolone pulse therapy combined with mizoribine following tonsillectomy for immunoglobulin A nephropathy: clinical remission rate, steroid sparing effect, and maintenance of renal function. 2011 , 15, 73-8	14
2024	Doubling of serum creatinine: is it appropriate as the endpoint for CKD? Proposal of a new surrogate endpoint based on the reciprocal of serum creatinine. 2011 , 15, 100-7	2
2023	Resistive index predicts renal prognosis in chronic kidney disease: results of a 4-year follow-up. 2011 , 15, 114-20	31

2022	The association between renal tubular damage and rapid renal deterioration in the Japanese population: the Takahata study. 2011 , 15, 235-41	29
2021	A case of slowly progressive scleroderma kidney. 2011 , 15, 430-433	0
2020	Risk of macrovascular disease stratified by stage of chronic kidney disease in type 2 diabetic patients: critical level of the estimated glomerular filtration rate and the significance of hyperuricemia. 2011 , 15, 391-397	19
2019	The relationship between renal volume and renal function in autosomal dominant polycystic kidney disease. 2011 , 15, 539-45	28
2018	Role of chronic kidney disease in cardiovascular disease: are we different from others?. 2011 , 15, 450-5	11
2017	Kidney function, albuminuria and cardiovascular risk factors in post-operative living kidney donors: a single-center, cross-sectional study. 2011 , 15, 514-21	11
2016	Estimated glomerular filtration rate in peritoneal dialysis practice. 2011 , 15, 611-2	1
2015	Clinical outcomes in patients with chronic kidney disease: a 5-year retrospective cohort study at a University Hospital in Japan. 2011 , 15, 831-40	8
2014	Safety and efficacy of administering the maximal dose of candesartan in renal transplant recipients. 2011 , 15, 907-15	3
2013	Significance of past history of renal failure for the detection of high-risk individuals for cardiovascular and end-stage renal disease: analysis of data from a nationwide health checkup. 2011 , 15, 841-7	2
2012	Pioglitazone reduces urinary albumin excretion in renin-angiotensin system inhibitor-treated type 2 diabetic patients with hypertension and microalbuminuria: the APRIME study. 2011 , 15, 848-53	15
2011	Effects of nicorandil on the reduction of BNP levels in patients with chronic kidney disease. 2011 , 15, 854-60	4
2010	Performance of serum cystatin C versus serum creatinine as a marker of glomerular filtration rate as measured by inulin renal clearance. 2011 , 15, 868-76	35
2009	Reduced renal function is associated with combined increases in ventricular-systolic stiffness and arterial load in patients undergoing cardiac catheterization for coronary artery disease. 2011 , 26, 10-6	10
2008	Impact of chronic kidney disease on the severity of initially diagnosed coronary artery disease and the patient prognosis in the Japanese population. 2011 , 26, 370-8	18
2007	Elevated B-type natriuretic peptide level as a marker of subsequent thromboembolic events in patients with atrial fibrillation. 2011 , 26, 530-5	20
2006	Low transition rate from normo- and low microalbuminuria to proteinuria in Japanese type 2 diabetic individuals: the Japan Diabetes Complications Study (JDACS). 2011 , 54, 1025-31	48
2005	The relation between estimated glomerular filtration rate and proteinuria in Okayama Prefecture, Japan. 2011 , 16, 191-5	3

2004	Association of the use of proton pump inhibitors with adverse cardiovascular and bleeding outcomes after percutaneous coronary intervention in the Japanese real world clinical practice. 2011 , 26, 222-33	9
2003	Long-term safety and efficacy of sirolimus-eluting stents versus bare-metal stents in real world clinical practice in Japan. 2011 , 26, 234-45	93
2002	Association between asymptomatic hyperuricemia and new-onset chronic kidney disease in Japanese male workers: a long-term retrospective cohort study. 2011 , 12, 31	37
2001	Low serum amylase in association with metabolic syndrome and diabetes: A community-based study. 2011 , 10, 34	57
2000	Impact of chronic kidney disease on the presence and severity of aortic stenosis in patients at high risk for coronary artery disease. 2011 , 9, 31	15
1999	Efficacy of adsorbent in delaying dialysis initiation among chronic kidney disease patients. 2011 , 40, 212-216	5
1998	Polymorphism of proinflammatory cytokine genes and albuminuria in the Japanese general population: the Takahata study. 2011 , 26, 3902-7	3
1997	Skin autofluorescence is associated with renal function and cardiovascular diseases in pre-dialysis chronic kidney disease patients. 2011 , 26, 214-20	32
1996	Assessing creatinine clearance from modification of diet in renal disease study equations in the ADEMEX cohort: limitations and potential applications. 2011 , 6, 598-604	3
1995	Effects of multiple factorial intervention on ambulatory BP profile and renal function in hypertensive type 2 diabetic patients with overt nephropathy - a pilot study. 2011 , 33, 255-63	11
1994	Urinary protein as a marker for systolic blood pressure reduction in patients with type 2 diabetes mellitus participating in an in-hospital diabetes education program. 2011 , 39, 70-5	
1993	Risks for glomerular filtration rate decline in association with progression of albuminuria in type 2 diabetes. 2011 , 26, 2924-30	30
1992	Serum level of triglycerides is a potent risk factor comparable to LDL cholesterol for coronary heart disease in Japanese patients with type 2 diabetes: subanalysis of the Japan Diabetes Complications Study (JDACS). 2011 , 96, 3448-56	135
1991	The effect of low-dose corticosteroids on short- and long-term renal outcome in patients with cholesterol crystal embolism. 2011 , 33, 298-306	21
1990	High prevalence of obstructive sleep apnea and its association with renal function among nondialysis chronic kidney disease patients in Japan: a cross-sectional study. 2011 , 6, 995-1000	92
1989	Prolonged carotid sinus reflex is a risk factor for contrast-induced nephropathy following carotid artery stenting. 2011 , 32, 441-5	11
1988	Nine-year incidence and risk factors for retinal vein occlusion in a general Japanese population: the Hisayama Study. 2011 , 52, 5905-9	51
1987	Urinary albumin excretion during angiotensin II receptor blockade: comparison of combination treatment with a diuretic or a calcium-channel blocker. 2011 , 24, 466-73	20

1986	Cigarette smoking and the association with glomerular hyperfiltration and proteinuria in healthy middle-aged men. 2011 , 6, 2462-9	59
1985	Baseline incidence and severity of renal insufficiency evaluated by estimated glomerular filtration rates in patients scheduled for contrast-enhanced CT. 2011 , 52, 581-6	12
1984	Brachial-ankle pulse wave velocity and rate of renal function decline and mortality in chronic kidney disease. 2011 , 6, 724-32	83
1983	Uric acid levels predict future development of chronic kidney disease. 2011 , 33, 352-7	74
1982	Urinary cystatin C as a potential risk marker for cardiovascular disease and chronic kidney disease in patients with obesity and metabolic syndrome. 2011 , 6, 265-73	29
1981	Low-dose aspirin therapy in patients with type 2 diabetes and reduced glomerular filtration rate: subanalysis from the JPAD trial. 2011 , 34, 280-5	29
1980	Predictive factors associated with the period of time before initiation of hemodialysis in CKD stages 4 and 5. 2011 , 117, c341-7	12
1979	Reduced estimated glomerular filtration rate is associated with stroke outcome after intravenous rt-PA: the Stroke Acute Management with Urgent Risk-Factor Assessment and Improvement (SAMURAI) rt-PA registry. 2011 , 31, 123-9	84
1978	Clinical characteristics of kidney disease in Japanese HIV-infected patients. 2011 , 118, c285-91	34
1977	Additional Treatment with Fenofibrate for Patients Treated with Pitavastatin Under Ordinary Medical Practice for Hypertriglyceridemia in Japan (APPROACH-J Study). 2011 , 2, 57-66	5
1976	Increased ACE and decreased ACE2 expression in kidneys from patients with IgA nephropathy. 2011 , 117, c57-66	31
1975	A predictive clinical grading system for immunoglobulin A nephropathy by combining proteinuria and estimated glomerular filtration rate. 2011 , 118, c292-300	10
1974	Renal function after catheter ablation of atrial fibrillation. 2011 , 124, 2380-7	46
1973	Clinical significance of urinary liver-type fatty acid-binding protein in diabetic nephropathy of type 2 diabetic patients. 2011 , 34, 691-6	96
1972	Combination of angiotensin II receptor antagonist with calcium channel blocker or diuretic as antihypertensive therapy for patients with chronic kidney disease. 2011 , 33, 366-72	11
1971	Combined effects of coffee consumption and serum γ -glutamyltransferase on serum C-reactive protein in middle-aged and elderly Japanese men and women. 2011 , 49, 1661-7	13
1970	Clinical impact of angiographic restenosis after bare-metal stent implantation on long-term outcomes in patients with coronary artery disease. 2011 , 75, 2566-72	10
1969	C-reactive protein is a predictor for developing proteinuria in a screened cohort. 2011 , 117, c51-6	3

1968	Abdominal dynamic CT in patients with renal dysfunction: contrast agent dose reduction with low tube voltage and high tube current-time product settings at 256-detector row CT. 2011 , 261, 467-76	89
1967	Association of serum leptin levels with progression of diabetic kidney disease in patients with type 2 diabetes. 2011 , 34, 2557-9	12
1966	Serum high-sensitivity cardiac troponin T is a significant biomarker of left-ventricular diastolic dysfunction in subjects with non-diabetic chronic kidney disease. 2011 , 1, 166-77	18
1965	Elevated white blood cell count worsens proteinuria but not estimated glomerular filtration rate: the Kansai Healthcare Study. 2011 , 34, 324-9	3
1964	Glomerular density-associated changes in clinicopathological features of minimal change nephrotic syndrome in adults. 2011 , 34, 542-8	17
1963	Risk of cardiovascular disease in kidney donors as a chronic kidney disease cohort. 2012 , 5, 7-11	2
1962	Decreased plasma level of vitamin C in chronic kidney disease: comparison between diabetic and non-diabetic patients. 2011 , 26, 1252-7	30
1961	The need for robust validation for MDRD-based glomerular filtration rate estimation in various CKD populations. 2011 , 26, 2780-5	70
1960	Asymptomatic coronary heart disease in patients with type 2 diabetes with vascular complications: a cross-sectional study. 2011 , 1, e000139	10
1959	Peritoneal macrophage infiltration is correlated with baseline peritoneal solute transport rate in peritoneal dialysis patients. 2011 , 26, 2322-32	28
1958	Long-term impact of mild chronic kidney disease in patients with acute coronary syndrome undergoing percutaneous coronary interventions. 2011 , 26, 2906-11	18
1957	Low glomerular density is a risk factor for progression in idiopathic membranous nephropathy. 2011 , 26, 3555-60	25
1956	Arterial stiffness is associated with incident albuminuria and decreased glomerular filtration rate in type 2 diabetic patients. 2011 , 34, 2570-5	64
1955	Angiotensin II receptor blocker and long-acting calcium channel blocker combination therapy decreases urinary albumin excretion while maintaining glomerular filtration rate. 2011 , 34, 1121-6	5
1954	Benidipine reduces albuminuria and plasma aldosterone in mild-to-moderate stage chronic kidney disease with albuminuria. 2011 , 34, 268-73	39
1953	Influence of adrenomedullin 2/intermedin gene polymorphism on blood pressure, renal function and silent cerebrovascular lesions in Japanese: the Ohasama study. 2011 , 34, 1327-32	10
1952	Prognostic significance of glomerular filtration rate estimated by the Japanese equation among patients who underwent complete coronary revascularization. 2011 , 34, 378-83	6
1951	Estimating glomerular filtration rate in Asian subjects: where do we stand?. 2011 , 80, 439-40	18

1950	Blood pressure components and risk for chronic kidney disease in middle-aged Japanese men: The Kansai Healthcare Study. 2011 , 34, 536-41	7
1949	High-normal blood pressure is associated with microalbuminuria in the general population: the Watari study. 2011 , 34, 1135-40	15
1948	Increased risk of cardiovascular events and mortality among non-diabetic chronic kidney disease patients with hypertensive nephropathy: the Gonryo study. 2011 , 34, 1106-10	42
1947	Aldosterone-to-renin ratio and home blood pressure in subjects with higher and lower sodium intake: the Ohasama study. 2011 , 34, 361-6	16
1946	Chronic kidney disease and subclinical lacunar infarction are independently associated with frontal lobe dysfunction in community-dwelling elderly subjects: the Sefuri brain MRI study. 2011 , 34, 1023-8	11
1945	Association between human metapneumovirus seroprevalence and hypertension in elderly subjects in a long-term care facility. 2011 , 34, 474-8	6
1944	Consequence of masked hypertension in treated hypertensive outpatients: 1-year follow-up study. 2011 , 33, 270-4	4
1943	Efficacy and safety of pitavastatin in Japanese patients with hypercholesterolemia: LIVES study and subanalysis. 2011 , 9, 555-62	29
1942	Malnutrition and inflammation determine prognosis of patients with CRS type 4. 2011 , 31, 276-80	3
1941	The clinician and estimation of glomerular filtration rate by creatinine-based formulas: current limitations and quo vadis. 2011 , 6, 937-50	73
1940	Relationship between renal function and the risk of recurrent atrial fibrillation following catheter ablation. 2011 , 97, 137-42	30
1939	Midlife and late-life blood pressure and dementia in Japanese elderly: the Hisayama study. 2011 , 58, 22-8	165
1938	Comparison of sustained hemodiafiltration with acetate-free dialysate and continuous venovenous hemodiafiltration for the treatment of critically ill patients with acute kidney injury. 2011 , 2011, 432094	19
1937	Arterial-cardiac destiffening following long-term antihypertensive treatment. 2011 , 24, 1080-6	19
1936	Dose-dense therapy is of benefit in primary treatment of ovarian cancer? In favor. 2011 , 22 Suppl 8, viii29-viii32	
1935	Design and rationale of the study of assessment for kidney function by urinary microalbumin in randomized (SAKURA) trial. 2011 , 33, 455-62	2
1934	Switching therapy from variable-dose multiple pill to fixed-dose single-pill combinations of angiotensin II receptor blockers and thiazides for hypertension. 2011 , 33, 309-15	7
1933	N-terminal pro-brain natriuretic peptide and risk of cardiovascular events in a Japanese community: the Hisayama study. 2011 , 31, 2997-3003	38

1932	Creatinine and HMH (5-hydroxy-1-methylhydantoin, NZ-419) as intrinsic hydroxyl radical scavengers. 2011 , 5, 162-75	10
1931	Plasma 25-hydroxyvitamin d is independently associated with hemoglobin concentration in male subjects with type 2 diabetes mellitus. 2011 , 2011, 362981	17
1930	Plasma aldosterone levels and development of insulin resistance: prospective study in a general population. 2011 , 58, 1043-8	45
1929	Effect of an Oral Adsorbent, AST-120, on Dialysis Initiation and Survival in Patients with Chronic Kidney Disease. 2012 , 2012, 376128	26
1928	High sodium intake is associated with masked hypertension in Japanese patients with type 2 diabetes and treated hypertension. 2012 , 25, 1170-4	15
1927	Association between prehypertension and chronic kidney disease in the Japanese general population. 2012 , 81, 293-9	34
1926	The resistive index is a marker of renal function, pathology, prognosis, and responsiveness to steroid therapy in chronic kidney disease patients. 2012 , 2012, 139565	52
1925	Evaluation of serum creatinine- and cystatin C-based equations for the estimation of glomerular filtration rate in a Chinese population. 2012 , 46, 223-31	12
1924	A slightly low hemoglobin level is beneficially associated with arterial stiffness in Japanese community-dwelling women. 2012 , 34, 92-8	31
1923	Relationship of ambulatory blood pressure and the heart rate profile with renal function parameters in hypertensive patients with chronic kidney disease. 2012 , 34, 264-9	10
1922	Association of body mass index with glomerular filtration rate in Japanese: a cross-sectional study in work-site population. 2012 , 34, 140-4	1
1921	Positive association between serum level of glyceraldehyde-derived advanced glycation end products and vascular inflammation evaluated by [(18)F]fluorodeoxyglucose positron emission tomography. 2012 , 35, 2618-25	67
1920	Plasma B-type natriuretic peptide level predicts kidney prognosis in patients with predialysis chronic kidney disease. 2012 , 27, 3885-91	32
1919	Short-term administration of diclofenac sodium affects renal function after laparoscopic radical nephrectomy in elderly patients. 2012 , 42, 1073-8	6
1918	Low glomerular density with glomerulomegaly in obesity-related glomerulopathy. 2012 , 7, 735-41	69
1917	Microalbuminuria is independently associated with deep or infratentorial brain microbleeds in hypertensive adults. 2012 , 25, 430-6	36
1916	Aliskiren reduces morning blood pressure in hypertensive patients with diabetic nephropathy. 2012 , 34, 243-8	5
1915	Factors Associated with the Decline of Kidney Function Differ among eGFR Strata in Subjects with Type 2 Diabetes Mellitus. 2012 , 2012, 687867	9

1914	Influence of renin angiotensin system gene polymorphisms on visit-to-visit blood pressure variability in hypertensive patients. 2012 , 25, 1249-55	6
1913	Association of high pulse pressure with proteinuria in subjects with diabetes, prediabetes, or normal glucose tolerance in a large Japanese general population sample. 2012 , 35, 1310-5	19
1912	A genome-wide association study of nephrolithiasis in the Japanese population identifies novel susceptible Loci at 5q35.3, 7p14.3, and 13q14.1. 2012 , 8, e1002541	54
1911	Effectiveness of etanercept vs cyclophosphamide as treatment for patients with amyloid A amyloidosis secondary to rheumatoid arthritis. 2012 , 51, 2064-9	16
1910	The long-term antiproteinuric effect of eplerenone, a selective aldosterone blocker, in patients with non-diabetic chronic kidney disease. 2012 , 13, 113-7	14
1909	Management of ureteral obstruction in advanced testicular tumor with lymph node metastasis. 2012 , 42, 748-52	6
1908	Impact of chronic kidney disease on a re-percutaneous coronary intervention for sirolimus-eluting stent restenosis. 2012 , 23, 528-32	6
1907	Association of sleep onset of acute coronary syndrome with sleep apnea syndrome and abnormal diurnal variation of hemostasis and adipokine levels. 2012 , 23, 590-6	4
1906	Histological and morphometric properties of skeletonized gastroepiploic artery and risk factors for intimal hyperplasia. 2012 , 7, 191-4	5
1905	Microvascular coronary artery spasm presents distinctive clinical features with endothelial dysfunction as nonobstructive coronary artery disease. 2012 , 1, e002485	48
1904	Combined use of vitamin D status and FGF23 for risk stratification of renal outcome. 2012 , 7, 810-9	87
1903	Blood pressure control in a Japanese population with chronic kidney disease: a baseline survey of a nationwide cohort. 2012 , 25, 342-7	21
1902	A candidate gene approach to genetic contributors to the development of IgA nephropathy. 2012 , 27, 1020-30	11
1901	WITHDRAWN: Enhanced cardio-renal protective effects of valsartan in high-risk hypertensive patients with chronic kidney disease: A sub-analysis of KYOTO HEART Study. 2012 ,	1
1900	Composition and plaque patterns of coronary culprit lesions and clinical characteristics of patients with chronic kidney disease. 2012 , 82, 344-51	40
1899	Associations of proteinuria, fluid volume imbalance, and body mass index with circadian ambulatory blood pressure in chronic kidney disease patients. 2012 , 36, 231-41	8
1898	Renal hypothermia with ice slush in laparoscopic partial nephrectomy: the outcome of renal function. 2012 , 26, 1483-8	16
1897	Endothelial impairment and bone marrow-derived CD34(+)/133(+) cells in diabetic patients with erectile dysfunction. 2012 , 3, 526-33	10

1896	Fluctuations in HbA1c are associated with a higher incidence of cardiovascular disease in Japanese patients with type 2 diabetes. 2012 , 3, 148-55	26
1895	Association between urinary angiotensinogen levels and renal and cardiovascular prognoses in patients with type 2 diabetes mellitus. 2012 , 3, 318-324	34
1894	The impact of CKD identification in large countries: the burden of illness. 2012 , 27 Suppl 3, iii32-8	74
1893	Association of coffee consumption with serum adiponectin, leptin, inflammation and metabolic markers in Japanese workers: a cross-sectional study. 2012 , 2, e33	78
1892	Spot urine-guided salt reduction is effective in Japanese cardiology outpatients. 2012 , 35, 1069-71	11
1891	Decreased glomerular filtration rate is a significant and independent risk for in-hospital mortality in Japanese patients with acute myocardial infarction: report from the Hokkaido acute myocardial infarction registry. 2012 , 35, 463-9	4
1890	The impact of visit-to-visit variability in blood pressure on renal function. 2012 , 35, 239-43	73
1889	Ethnic factors of the glomerular filtration rate estimating equation. 2012 , 81, 799; author reply 799-800	3
1888	Comparison of medium-dose losartan/hydrochlorothiazide and maximal-dose angiotensin II receptor blockers in the treatment of Japanese patients with uncontrolled hypertension: the Kobe-CONNECT Study. 2012 , 35, 1080-6	5
1887	Impact of serum uric acid on renal function and cardiovascular events in hypertensive patients treated with losartan. 2012 , 35, 867-73	28
1886	Association of severe hypertension with pneumonia in elderly patients with acute ischemic stroke. 2012 , 35, 648-53	21
1885	Estimating glomerular filtration rates by use of both cystatin C and standardized serum creatinine avoids ethnicity coefficients in Asian patients with chronic kidney disease. 2012 , 58, 450-7	25
1884	Significance of urinary full-length and ectodomain forms of megalin in patients with type 2 diabetes. 2012 , 35, 1112-8	41
1883	The great East Japan earthquake: blood pressure control in patients with chronic kidney disease. 2012 , 25, 951-4	22
1882	Surgical and oncologic outcomes of laparoscopic partial nephrectomy: a Japanese multi-institutional study of 1375 patients. 2012 , 26, 652-9	9
1881	Renal hyperfiltration in prediabetes confirmed by fasting plasma glucose and hemoglobin A1c. 2012 , 34, 1084-90	13
1880	Prevalence of and factors associated with chronic kidney disease (CKD) in Japanese subjects without notable chronic diseases, undergoing an annual health checkup. 2012 , 36, 139-48	12
1879	Nephronectin expression in glomeruli of renal biopsy specimens from various kidney diseases: nephronectin is expressed in the mesangial matrix expansion of diabetic nephropathy. 2012 , 122, 114-21	12

1878	Low urine pH is a predictor of chronic kidney disease. 2012 , 35, 77-81	40
1877	Circulating fibrocytes in ischemia-reperfusion injury and chronic renal allograft fibrosis. 2012 , 121, c16-24	6
1876	Serum antibody to Porphyromonas gingivalis in chronic kidney disease. 2012 , 91, 828-33	15
1875	Serum cystatin C as a marker for early detection of chronic kidney disease and grade 2 nephropathy in Japanese patients with type 2 diabetes. 2012 , 50, 1833-9	10
1874	Low contrast agent and radiation dose protocol for hepatic dynamic CT of thin adults at 256-detector row CT: effect of low tube voltage and hybrid iterative reconstruction algorithm on image quality. 2012 , 264, 445-54	112
1873	The Relationship between Renal Dysfunction and Abnormalities of the Immune System in Patients with Decompensated Cirrhosis. 2012 , 2012, 123826	2
1872	Effects of sex and age on serum 1,5-anhydroglucitol in nondiabetic subjects. 2012 , 120, 288-95	20
1871	Risk factors predisposing to stroke recurrence within one year of non-cardioembolic stroke onset: the Fukuoka Stroke Registry. 2012 , 33, 141-9	47
1870	Glomerular expression of hydrogen peroxide-inducible clone-5 in human and rat progressive mesangial proliferative glomerulonephritis. 2012 , 120, e59-68	5
1869	Hypobilirubinemia might be a possible risk factor of end-stage kidney disease independently of estimated glomerular filtration rate. 2012 , 36, 47-54	13
1868	Superimposed preeclampsia in women with chronic kidney disease. 2012 , 74, 274-81	42
1867	Comparison of renal function after partial nephrectomy and radical nephrectomy for renal cell carcinoma. 2012 , 89, 227-32	16
1866	Effects of carvedilol as third-line add-on therapy on blood pressure and glucose metabolism in type 2 diabetic patients with chronic renal disease stage 3 and above. 2012 , 36, 131-8	2
1865	Aldosterone-to-renin ratio as a predictor of stroke under conditions of high sodium intake: the Ohasama study. 2012 , 25, 777-83	19
1864	Relationship between serum adiponectin levels and disability-free survival among community-dwelling elderly individuals: The Tsurugaya Project. 2012 , 67, 530-6	18
1863	Comparison of the clinical outcome and systemic inflammatory marker levels between retroperitoneal and transperitoneal laparoscopic donor nephrectomy. 2012 , 26, 1038-43	4
1862	Association of home blood pressure variability with progression of chronic kidney disease. 2012 , 17, 1-7	30
1861	KDIGO (Kidney Disease: Improving Global Outcomes) criteria could be a useful outcome predictor of cisplatin-induced acute kidney injury. 2012 , 82, 354-9	20

1860	Pharmacokinetic study of S-1 in patients in whom inulin clearance was measured. 2012 , 83, 38-44	5
1859	Effects of carperitide on contrast-induced acute kidney injury with a minimum volume of contrast in chronic kidney disease patients. 2012 , 2, 303-10	10
1858	Pre-existing arteriosclerotic intimal thickening in living-donor kidneys reflects allograft function. 2012 , 36, 127-35	19
1857	A cross-sectional study of glucose regulation in young adults with very low birth weight: impact of male gender on hyperglycaemia. 2012 , 2, e000327	10
1856	Impact of chronic kidney disease on the incidence of peri-procedural myocardial injury in patients undergoing elective stent implantation. 2012 , 27, 1059-63	16
1855	Gender differences in the association between HDL cholesterol and the progression of diabetic kidney disease in type 2 diabetic patients. 2012 , 27, 1070-5	26
1854	Pre-hypertension as a significant predictor of chronic kidney disease in a general population: the Ohasama Study. 2012 , 27, 3218-23	35
1853	Serum hepcidin-25 levels predict the progression of renal anemia in patients with non-dialysis chronic kidney disease. 2012 , 27, 4378-85; discussion 4384-5	13
1852	Predicting outcomes after myocardial infarction by using the Chronic Kidney Disease Epidemiology Collaboration equation in comparison with the Modification of Diet in Renal Disease study equation: results from the Korea Acute Myocardial Infarction Registry. 2012 , 27, 3868-74	10
1851	Glomerular hyperfiltration in prediabetes and prehypertension. 2012 , 27, 1821-5	98
1850	Changes in urinary albumin excretion, inflammatory and oxidative stress markers in ADPKD patients with hypertension. 2012 , 343, 46-51	29
1849	Glycohemoglobin not as predictive as fasting glucose as a measure of prediabetes in predicting proteinuria. 2012 , 27, 3862-8	11
1848	Comparisons between validated estimated glomerular filtration rate equations and isotopic glomerular filtration rate in HIV patients. 2012 , 26, 1781-8	30
1847	Hyperuricemia at 1 year after renal transplantation, its prevalence, associated factors, and graft survival. 2012 , 94, 145-51	25
1846	DNA hypermethylation and inflammatory markers in incident Japanese dialysis patients. 2012 , 2, 159-68	15
1845	Evaluation of validity of the urine dipstick test for identification of reduced glomerular filtration rate in Japanese male workers aged 40 years and over. 2012 , 54, 176-80	2
1844	CYP2D6 genotype affects age-related decline in flecainide clearance: a population pharmacokinetic analysis. 2012 , 22, 777-83	11
1843	Plasma renin activity and the aldosterone-to-renin ratio are associated with the development of chronic kidney disease: the Ohasama Study. 2012 , 30, 1632-8	24

1842	Evaluation of risk for incident hypertension using glomerular filtration rate in the normotensive general population. 2012 , 30, 505-12	20
1841	Diagnostic performance of cardiac fusion images from myocardial perfusion imaging and multislice computed tomography coronary angiography for assessment of hemodynamically significant coronary artery lesions: an observational study. 2012 , 33, 60-8	9
1840	Impact of chronic kidney disease on outcomes of surgical resection for primary colorectal cancer: a retrospective cohort review. 2012 , 55, 948-56	14
1839	Association between asleep blood pressure and brain natriuretic peptide during antihypertensive treatment: the Japan Morning Surge-Target Organ Protection (J-TOP) study. 2012 , 30, 1015-21	13
1838	[Chronic kidney disease (CKD)--recent progress. Editorial: CKD initiative in these 10 years and perspective]. 2012 , 101, 1233-5	
1837	[Chronic kidney disease (CKD)--recent progress. Topics: I. Concept and importance]. 2012 , 101, 1236-42	
1836	[Chronic kidney disease (CKD)--recent progress. Topics: III. New classification of chronic kidney disease (CKD)]. 2012 , 101, 1253-8	
1835	[Chronic kidney disease (CKD)--recent progress. Topics: IV. Assessment of renal function]. 2012 , 101, 1259-65	1
1834	High serum thioredoxin levels are reduced after tonsillectomy in patients with IgA nephropathy. 2012 , 51, 559-65	3
1833	Deep echo attenuation without calcification increases the risk of periprocedural myonecrosis after elective percutaneous coronary intervention in patients with coronary artery disease. 2012 , 51, 691-8	4
1832	A case of diaphragmatic paralysis complicated by herpes-zoster virus infection. 2012 , 51, 1259-63	15
1831	Non-alcoholic fatty liver disease is an independent predictor for macroangiopathy in Japanese type 2 diabetic patients: a cross-sectional study. 2012 , 51, 1667-75	10
1830	The limited usefulness of the treadmill test or a risk-guided approach in screening for asymptomatic coronary heart disease in Japanese patients with type 2 diabetes. 2012 , 51, 3337-42	2
1829	[Chronic kidney disease (CKD)--recent progress. Topics: II. Epidemiology of CKD]. 2012 , 101, 1243-52	
1828	Low serum free testosterone level is associated with carotid intima-media thickness in middle-aged Japanese men. 2012 , 59, 809-15	14
1827	Job stress strengthens the link between metabolic risk factors and renal dysfunction in adult men. 2012 , 226, 101-8	13
1826	Pre-procedural glucose levels and the risk for contrast-induced acute kidney injury in patients undergoing emergency coronary intervention. 2012 , 76, 1848-55	16
1825	Predictor of early mortality for severe heart failure patients with left ventricular assist device implantation: significance of INTERMACS level and renal function. 2012 , 76, 1631-8	62

1824	Plasma endothelin-1 level is a predictor of 10-year mortality in a general population: the Tanushimaru study. 2012 , 76, 2779-84	19
1823	[Clinical usefulness of the new Japanese glomerular filtration rate equation for initial and individualized dosage adjustment concentrations of vancomycin]. 2012 , 132, 125-33	
1822	[What is chronic kidney disease (CKD)?: early check and early treatment of kidney disease]. 2012 , 132, 441-2	
1821	[Appropriate pharmacotherapy in patients with chronic kidney disease - new approach -]. 2012 , 132, 461-70	1
1820	Usefulness of urinary biomarkers in early detection of acute kidney injury after cardiac surgery in adults. 2012 , 76, 213-20	72
1819	Impact of body mass index on clinical outcome in patients hospitalized with congestive heart failure. 2012 , 76, 145-51	34
1818	Cardiac event risk in Japanese subjects estimated using gated myocardial perfusion imaging, in conjunction with diabetes mellitus and chronic kidney disease. 2012 , 76, 168-75	33
1817	Gender-specific relationship between serum uric acid level and atrial fibrillation prevalence. 2012 , 76, 607-11	54
1816	Impact of low levels of vascular endothelial growth factor after myocardial infarction on 6-month clinical outcome. Results from the Nagoya Acute Myocardial Infarction Study. 2012 , 76, 1509-16	27
1815	Functional limitations predict the risk of rehospitalization among patients with chronic heart failure. 2012 , 76, 1654-61	22
1814	Clinical significance of reactive post-capillary pulmonary hypertension in patients with left heart disease. 2012 , 76, 1235-44	37
1813	Impact of metabolic syndrome on various aspects of microcirculation and major adverse cardiac events in patients with ST-segment elevation myocardial infarction. 2012 , 76, 1972-9	16
1812	Impact of diabetes mellitus on coronary atherosclerosis and plaque composition under statin therapy: Subanalysis of the TRUTH study [2012, 76, 2188-96	17
1811	Radiofrequency catheter ablation of persistent atrial fibrillation decreases a sleep-disordered breathing parameter during a short follow-up period. 2012 , 76, 2096-103	25
1810	Effect of obesity on declining beta cell function after diagnosis of type 2 diabetes: a possible link suggested by cross-sectional analysis. 2012 , 59, 187-95	16
1809	Impact of chronic kidney disease on left main coronary artery disease and prognosis in Japanese patients. 2012 , 76, 2266-72	12
1808	Contribution of extracardiac factors to the inconsistency between plasma B-type natriuretic peptide levels and the severity of pulmonary congestion on chest X-rays in the diagnosis of heart failure. 2012 , 51, 239-48	14
1807	An elderly patient with diabetic nephropathy complicated by ANCA-associated nephritis. 2012 , 51, 1227-32	3

1806	Impact of decreased estimated glomerular filtration rate on Japanese acute stroke and its subtype. 2012 , 51, 1661-6	24
1805	Retrospective comparison of the efficacy of tonsillectomy with and without steroid-pulse therapy in IgA nephropathy patients. 2012 , 51, 1323-8	8
1804	Prevalence of chronic kidney disease and blood pressure control status in elderly hypertensive patients. 2012 , 51, 1473-8	4
1803	Risk assessment for acute kidney injury after allogeneic hematopoietic stem cell transplantation based on Acute Kidney Injury Network criteria. 2012 , 51, 2105-10	38
1802	Matrix metalloproteinase-9 gene polymorphisms and chronic kidney disease. 2012 , 36, 444-50	9
1801	Relationships between levels of urinary podocalyxin, number of urinary podocytes, and histologic injury in adult patients with IgA nephropathy. 2012 , 7, 1385-93	53
1800	Type II diabetes mellitus is a risk factor for heart failure in pre-dialysis patients. 2012 , 16, 541-7	2
1799	Impacts of estimated glomerular filtration rate on coronary atherosclerosis and plaque composition before and during statin therapy in patients with normal to mild renal dysfunction: subanalysis of the TRUTH study. 2012 , 17, 628-35	3
1798	Impact of hyponatremia on survival of patients with metastatic renal cell carcinoma treated with molecular targeted therapy. 2012 , 19, 1050-7	21
1797	Influence of endoscopic submucosal dissection on serum levels of pepsinogens in patients with early gastric cancer. 2012 , 24, 339-42	3
1796	Arterial stiffness and renal impairment in non-proteinuric type 2 diabetic patients. 2012 , 3, 86-91	3
1795	Duration until nighttime blood pressure fall indicates excess sodium retention. 2012 , 29, 1412-7	12
1794	Urinary N-acetyl-β-D-glucosaminidase levels are positively correlated with 2-hr plasma glucose levels during oral glucose tolerance testing in prediabetes. 2012 , 26, 473-80	16
1793	The result of adjuvant chemotherapy for localized pT3 upper urinary tract carcinoma in a multi-institutional study. 2012 , 30, 701-6	24
1792	Blood cadmium and moderate-to-severe glomerular dysfunction in Korean adults: analysis of KNHANES 2005-2008 data. 2012 , 85, 885-93	19
1791	Relationship between kidney damage and stroke types in Japanese patients. 2012 , 16, 564-9	10
1790	Evaluation of estimated creatinine clearance before steady state in acute kidney injury by creatinine kinetics. 2012 , 16, 570-9	19
1789	Renal disease progression in autosomal dominant polycystic kidney disease. 2012 , 16, 622-8	25

1788	Serum levels of soluble secreted β Klotho are decreased in the early stages of chronic kidney disease, making it a probable novel biomarker for early diagnosis. 2012 , 16, 722-9	127
1787	Relationship between renal hemodynamic status and aging in patients without diabetes evaluated by renal Doppler ultrasonography. 2012 , 16, 786-91	10
1786	Differences in the local and national prevalences of chronic kidney disease based on annual health check program data. 2012 , 16, 749-54	1
1785	Long-term improvement of oxidative stress via kidney transplantation ameliorates serum sulfatide levels. 2012 , 16, 959-67	8
1784	Effects of genetic variants in SLC22A2 organic cation transporter 2 and SLC47A1 multidrug and toxin extrusion 1 transporter on cisplatin-induced adverse events. 2012 , 16, 843-51	70
1783	Complete remission within 2 years predicts a good prognosis after methylprednisolone pulse therapy in patients with IgA nephropathy. 2012 , 16, 883-91	23
1782	Plasma accumulation of fondaparinux 2.5mg in patients after total hip arthroplasty. 2012 , 34, 526-32	6
1781	Circulating high-mobility group box 1 and cardiovascular mortality in unstable angina and non-ST-segment elevation myocardial infarction. 2012 , 221, 490-5	41
1780	Non-linear association between ankle-brachial pressure index and prevalence of silent cerebral infarction in Japanese patients with type 2 diabetes. 2012 , 222, 490-4	13
1779	Prevalence and risk factors for cerebral infarction and carotid artery stenosis in peripheral arterial disease. 2012 , 223, 473-7	23
1778	Serum levels of pigment epithelium-derived factor, a novel marker of insulin resistance, are independently associated with fasting apolipoprotein B48 levels in humans. 2012 , 45, 1404-8	6
1777	Differential effects of strict blood pressure lowering by losartan/hydrochlorothiazide combination therapy and high-dose amlodipine monotherapy on microalbuminuria: the ALPHABET study. 2012 , 6, 73-82	10
1776	The effects of blood pressure control levels on the renoprotection of type 2 diabetic patients without overt proteinuria. 2012 , 6, 124-31	9
1775	Chronic kidney disease in patients with ischemic stroke. 2012 , 21, 547-50	25
1774	Lower levels of high-density lipoprotein cholesterol on admission and a recurrence of ischemic stroke: a 12-month follow-up of the Fukuoka Stroke Registry. 2012 , 21, 561-8	8
1773	A decade after the KDOQI CKD guidelines: a perspective from Japan. <i>American Journal of Kidney Diseases</i> , 2012 , 60, 729-30	7-4 5
1772	Ongoing myocardial damage in patients with heart failure and preserved ejection fraction. 2012 , 60, 454-61	24
1771	Prognostic impact of chronic kidney disease on 10-year clinical outcomes among patients with acute coronary syndrome. 2012 , 60, 438-42	18

1770	Risk stratification of cardiovascular events in patients at all stages of chronic kidney disease using myocardial perfusion SPECT. 2012 , 60, 377-82	27
1769	Prognostic impact of systolic blood pressure at admission on in-hospital outcome after primary percutaneous coronary intervention for acute myocardial infarction. 2012 , 60, 139-44	20
1768	Obesity may attenuate the HbA1c-lowering effect of sitagliptin in Japanese type 2 diabetic patients. 2012 , 3, 170-4	27
1767	Asymptomatic diverticulosis identified by computed tomography is not a risk factor for enteric peritonitis. 2012 , 27, 2511-6	19
1766	Efficacy of aliskiren in Japanese chronic kidney disease patients with hypertension. 2012 , 34, 442-7	4
1765	Association of hypoalbuminemia with severe anemia in patients with diabetic nephrosclerosis. 2012 , 34, 189-93	5
1764	Additive renoprotective effects of aliskiren on angiotensin receptor blocker and calcium channel blocker treatments for type 2 diabetic patients with albuminuria. 2012 , 35, 874-81	15
1763	microRNA and the pathogenesis of ovarian cancer--a new horizon for molecular diagnostics and treatment?. 2012 , 50, 601-15	26
1762	Serum hepcidin-25 levels and anemia in non-dialysis chronic kidney disease patients: a cross-sectional study. 2012 , 27, 1076-83	51
1761	Frequency of class III and IV nephritis in systemic lupus erythematosus without clinical renal involvement: an analysis of predictive measures. 2012 , 39, 79-85	62
1760	The usefulness of the revised classification for chronic kidney disease by the KDIGO for determining the frequency of diabetic micro- and macroangiopathies in Japanese patients with type 2 diabetes mellitus. 2012 , 26, 286-90	21
1759	Relationship between chronic kidney disease and white matter hyperintensities on magnetic resonance imaging. 2012 , 21, 18-23	28
1758	Chronic kidney disease and carotid atherosclerosis. 2012 , 21, 47-51	29
1757	Deep white matter hyperintensities, decreased serum low-density lipoprotein, and dilative large arteriopathy. 2012 , 21, 225-30	7
1756	Clinical features and neurologic severity in stroke patients with mild to moderate renal dysfunction. 2012 , 21, 343-9	8
1755	HbA1c and atrial fibrillation: a cross-sectional study in Japan. 2012 , 156, 156-9	30
1754	High-sensitive troponin T is associated with atrial fibrillation in a general population. 2012 , 156, 98-100	23
1753	Intact fibroblast growth factor 23 levels predict incident cardiovascular event before but not after the start of dialysis. 2012 , 50, 1266-74	71

1752	Prognostic value of serum tenascin-C levels on long-term outcome after acute myocardial infarction. 2012 , 18, 480-6	37
1751	Antialbuminuric advantage of cilnidipine compared with L-type calcium channel blockers in type 2 diabetic patients with normoalbuminuria and microalbuminuria. 2012 , 97, 91-8	12
1750	High serum S100A8/A9 levels and high cardiovascular complication rate in type 2 diabetics with ultrasonographic low carotid plaque density. 2012 , 97, 82-90	16
1749	Plasminogen activator inhibitor-1 is associated with renal dysfunction independent of BMI and serum lipid levels in patients with type 2 diabetes. 2012 , 97, e9-12	4
1748	Increased circulating plasma lysophosphatidic acid in patients with acute coronary syndrome. 2012 , 413, 207-12	35
1747	Predicting the outcome of chronic kidney disease by the estimated nephron number: the rationale and design of PRONEP, a prospective, multicenter, observational cohort study. 2012 , 13, 11	9
1746	Significant association between renal function and amyloid-positive area in renal biopsy specimens in AL amyloidosis. 2012 , 13, 118	10
1745	Hypertension resistant to antihypertensive agents commonly occurs with the progression of diabetic nephropathy in Japanese patients with type 2 diabetes mellitus: a prospective observational study. 2012 , 13, 48	10
1744	Latent associations of low serum amylase with decreased plasma insulin levels and insulin resistance in asymptomatic middle-aged adults. 2012 , 11, 80	26
1743	Effect of erythropoietin-stimulating agent on uremic inflammation. 2012 , 9, 17	12
1742	Effects of ezetimibe on visceral fat in the metabolic syndrome: a randomised controlled study. 2012 , 42, 1287-94	34
1741	Simplified dosing regimens of teicoplanin for patient groups stratified by renal function and weight using Monte Carlo simulation. 2012 , 40, 344-8	8
1740	Predictors of worsening renal function after computed tomography coronary angiography: assessed by cystatin C. 2012 , 6, 31-6	6
1739	A HILIC-MS/MS method for the simultaneous determination of seven organic acids in rat urine as biomarkers of exposure to realgar. 2012 , 905, 37-42	37
1738	Renal function and effect of statin therapy on cardiovascular outcomes in patients undergoing coronary revascularization (from the CREDO-Kyoto PCI/CABG Registry Cohort-2). 2012 , 110, 1568-77	28
1737	Dialysis vintage and parathyroid hormone level, not fibroblast growth factor-23, determines chronic-phase phosphate wasting after renal transplantation. 2012 , 51, 729-36	16
1736	The factors associated with a functional outcome after ischemic stroke in diabetic patients: the Fukuoka Stroke Registry. 2012 , 313, 110-4	11
1735	Clinical significance of the measurements of urinary liver-type fatty acid binding protein levels in patients with acute coronary syndrome. 2012 , 60, 168-73	13

1734	Depressive symptoms are common and associated with adverse clinical outcomes in heart failure with reduced and preserved ejection fraction. 2012 , 60, 23-30	42
1733	Three-year experience with alendronate treatment in postmenopausal osteoporotic Japanese women with or without renal dysfunction: a retrospective study. 2012 , 29, 133-42	2
1732	Upright T waves in lead aVR are associated with cardiac death or hospitalization for heart failure in patients with a prior myocardial infarction. 2012 , 27, 548-52	15
1731	Association between deteriorated renal function and long-term clinical outcomes after percutaneous coronary intervention. 2012 , 27, 460-7	13
1730	Population pharmacokinetics of transdermal fentanyl in patients with cancer-related pain. 2012 , 26, 98-104	19
1729	Interactions between serum vitamin D levels and vitamin D receptor gene FokI polymorphisms for renal function in patients with type 2 diabetes. 2012 , 7, e51171	15
1728	Chronic kidney disease, dyslipidemia, and atherosclerosis. 2012 , 19, 299-315	50
1727	Adding thiazide to a rennin-angiotensin blocker regimen to improve left ventricular relaxation in diabetes and nondiabetes patients with hypertension. 2012 , 6, 225-33	2
1726	Use of Antiarrhythmic Drug in Patients with Impaired Renal Function. 2012 , 32, 327-332	
1725	Systematic analysis of risk factors for coronary heart disease in Japanese patients with type 2 diabetes: a matched case-control study. 2012 , 19, 918-23	3
1724	Impact of low-and high-density lipoprotein cholesterol levels on carotid intima-media thickness differs by smoking status in middle-aged men. 2012 , 19, 664-72	2
1723	Age-dependent decline in renal function in patients with lung cancer. 2012 , 4, 38-42	5
1722	Chronic kidney disease in patients with ileal conduit urinary diversion. 2012 , 4, 962-966	9
1721	A reduction of coronary flow reserve is associated with chronic kidney disease and long-term cardio-cerebrovascular events in patients with non-obstructive coronary artery disease and vasospasm. 2012 , 58, 136-43	8
1720	Uric acid increases the incidence of ventricular arrhythmia in patients with left ventricular hypertrophy. 2012 , 58, 101-6	15
1719	Associations between healthy lifestyle behaviors and proteinuria and the estimated glomerular filtration rate (eGFR). 2012 , 19, 932-40	24
1718	Aliskiren suppresses the renin-angiotensin-aldosterone system and reduces blood pressure and albuminuria in elderly chronic kidney disease patients with hypertension. 2012 , 5, 125-33	2
1717	Effectiveness of metformin and lifestyle interventions as an initial treatment in Japanese patients with newly diagnosed type 2 diabetes: a prospective observational study. 2012 , 59, 166-73	9

1716	Plasma B-type natriuretic peptide is useful for cardiovascular risk assessment in community-based diabetes subjects: comparison with albuminuria. 2012 , 53, 176-81	16
1715	Reduction in urinary albumin excretion with a moderate low-carbohydrate diet in patients with type 2 diabetes: a 12-month intervention. 2012 , 5, 283-91	7
1714	The impact of different GFR estimating equations on the prevalence of CKD and risk groups in a Southeast Asian cohort using the new KDIGO guidelines. 2012 , 13, 1	30
1713	Tonsillectomy has beneficial effects on remission and progression of IgA nephropathy independent of steroid therapy. 2012 , 27, 2806-13	54
1712	Risk of overestimation of kidney function using GFR-estimating equations in patients with low inulin clearance. 2012 , 26, 248-53	9
1711	Proteinuria and clinical outcomes after ischemic stroke. 2012 , 78, 1909-15	126
1710	Hypomagnesemia in type 2 diabetic nephropathy: a novel predictor of end-stage renal disease. 2012 , 35, 1591-7	81
1709	Use of a new high-sensitivity assay for cardiac troponin T to stratify the risk of cardiovascular disease in outpatients with type-2 diabetes. 2012 , 3, 29-36	5
1708	Chronic kidney disease categories and renal-cardiovascular outcomes in type 2 diabetes without prevalent cardiovascular disease: a prospective cohort study (JDDM25). 2012 , 55, 1911-8	33
1707	Cigarette smoking, proteinuria, and renal function in middle-aged Japanese men from an occupational population. 2012 , 17, 147-56	7
1706	Association between frequency of drinking alcohol and chronic kidney disease in men. 2012 , 17, 199-204	16
1705	A novel Doppler echocardiographic score reflecting cardiac functional status can predict adverse outcome in acute myocardial infarction. 2012 , 10, 41-7	0
1704	Steady-state B-type natriuretic peptide levels in patients with atrial fibrillation of various clinical backgrounds. 2012 , 27, 280-6	8
1703	Plasma endoglin as a marker to predict cardiovascular events in patients with chronic coronary artery diseases. 2012 , 27, 344-51	39
1702	Sunitinib-induced nephrotic syndrome and irreversible renal dysfunction. 2012 , 16, 310-5	31
1701	Risk factor profiles based on estimated glomerular filtration rate and dipstick proteinuria among participants of the Specific Health Check and Guidance System in Japan 2008. 2012 , 16, 244-9	44
1700	Gene polymorphisms contributing to hypertension in immunoglobulin A nephropathy. 2012 , 16, 250-8	12
1699	Weight gain after 20 years of age is associated with prevalence of chronic kidney disease. 2012 , 16, 259-68	16

1698	How to manage HIV-infected patients with chronic kidney disease in the HAART era. 2012 , 16, 363-72		26
1697	Performance of the Japanese GFR equation in potential kidney donors. 2012 , 16, 415-20		14
1696	Vitamin D deficiency in elderly women in nursing homes: investigation with consideration of decreased activation function from the kidneys. 2012 , 60, 251-5		7
1695	Histological predictors for renal prognosis in diabetic nephropathy in diabetes mellitus type 2 patients with overt proteinuria. 2012 , 17, 68-75		66
1694	Renal impairment after laparoscopic radical nephrectomy affects hypoglycaemic therapy. 2012 , 37, 49-52		2
1693	Ideal timing and predialysis nephrology care duration for dialysis initiation: from analysis of Japanese dialysis initiation survey. 2012 , 16, 54-62		22
1692	Beneficial effect of Astragalus membranaceus on estimated glomerular filtration rate in patients with progressive chronic kidney disease. 2012 , 14, 17-23		9
1691	Relation between estimated glomerular filtration rate and composition of coronary arterial atherosclerotic plaques. 2012 , 109, 1131-6		21
1690	Off-pump bilateral skeletonized internal thoracic artery grafting in elderly patients. 2012 , 93, 531-6		40
1689	Periodontal disease and decreased kidney function in Japanese elderly. <i>American Journal of Kidney Diseases</i> , 2012 , 59, 202-9	7.4	65
1688	A case of kidney failure and glycosuria. <i>American Journal of Kidney Diseases</i> , 2012 , 59, xxxiii-xxxvi	7.4	2
1687	Time course and prognostic implications of QT interval in patients with coronary artery disease undergoing coronary bypass surgery. 2012 , 23, 645-9		6
1686	Effects of aliskiren-based therapy on ambulatory blood pressure profile, central hemodynamics, and arterial stiffness in nondiabetic mild to moderate hypertensive patients. 2012 , 14, 522-9		14
1685	Late dialysis start did not affect long-term outcome in Japanese dialysis patients: long-term prognosis from Japanese Society for [corrected] Dialysis Therapy Registry. 2012 , 16, 111-20		28
1684	High predictive accuracy of Aldosteronoma Resolution Score in Japanese patients with aldosterone-producing adenoma. 2012 , 151, 437-43		36
1683	Conversion of stable kidney transplant recipients from a twice-daily prograf to a once-daily tacrolimus formulation: a short-term study on its effects on glucose metabolism. 2012 , 44, 128-33		16
1682	Pharmacokinetics of mizoribine in adult living donor liver transplantation. 2012 , 44, 1329-35		7
1681	Visualization of coronary plaque in type 2 diabetes mellitus patients using a new 40 MHz intravascular ultrasound imaging system. 2012 , 59, 42-9		17

1680	The number of leads with fragmented QRS is independently associated with cardiac death or hospitalization for heart failure in patients with prior myocardial infarction. 2012 , 59, 36-41	50
1679	Clinical impact of thrombus aspiration during primary percutaneous coronary intervention: results from Korea Acute Myocardial Infarction Registry. 2012 , 59, 249-57	19
1678	Urinary liver-type fatty acid-binding protein level as a predictive biomarker of contrast-induced acute kidney injury. 2012 , 42, 557-63	48
1677	Development of self-care educational material for patients with heart failure in Japan: a pilot study. 2012 , 14, 156-64	7
1676	Impact of renal function on cardiovascular events in patients undergoing radical nephrectomy for renal cancer. 2012 , 19, 722-8	16
1675	Predictors for postoperative renal function after open partial nephrectomy: including postoperative biomarkers. 2012 , 19, 823-8	16
1674	Cytoreductive nephrectomy for pT3b-T4 M1 renal cell carcinoma in 39 patients: single center analysis. 2012 , 19, 875-9	2
1673	Similar functional outcomes after partial nephrectomy for clinical T1b and T1a renal cell carcinoma. 2012 , 19, 980-6	13
1672	Skin autofluorescence is associated with severity of vascular complications in Japanese patients with Type 2 diabetes. 2012 , 29, 492-500	39
1671	Seasonal variations of urinary albumin creatinine ratio in Japanese subjects with Type 2 diabetes and early nephropathy. 2012 , 29, 506-8	11
1670	Pro-/anti-inflammatory cytokine gene polymorphisms and chronic kidney disease: a cross-sectional study. 2012 , 13, 2	27
1669	Distribution of estimated glomerular filtration rate (eGFR) values in patients receiving contrast-enhanced magnetic resonance imaging. 2012 , 30, 116-9	
1668	C-reactive protein levels, airflow obstruction, and chronic kidney disease. 2012 , 17, 18-26	4
1667	The correlation of common carotid arterial diameter with atherosclerosis and diabetic retinopathy in patients with type 2 diabetes mellitus. 2012 , 49, 63-8	25
1666	Visualization of coronary plaque in arterial remodeling using a new 40-MHz intravascular ultrasound imaging system. 2013 , 81, 471-80	14
1665	Low-molecular-weight lipoprotein (a) and low relative lymphocyte concentration are significant and independent risk factors for coronary heart disease in patients with type 2 diabetes mellitus: Lp(a) phenotype, lymphocyte, and coronary heart disease. 2013 , 12, 31	11
1664	Reduced fractional shortening of right ventricular outflow tract is associated with adverse outcomes in patients with left ventricular dysfunction. 2013 , 11, 19	9
1663	Low serum eicosapentaenoic acid / arachidonic acid ratio in male subjects with visceral obesity. 2013 , 10, 25	33

1662	Correlation of circulating C1q and C1q-binding adiponectin concentrations with aging in males: a preliminary report. 2013 , 5, 17	2
1661	T/L-type calcium channel blocker reduces the composite ranking of relative risk according to new KDIGO guidelines in patients with chronic kidney disease. 2013 , 14, 135	6
1660	Determinants of the HbA1c-lowering effect of sitagliptin when added to ongoing insulin therapy in Japanese patients with type 2 diabetes. 2013 , 4, 251-256	2
1659	The area of abdominal subcutaneous adipose tissue is independently correlated with C-peptide increment during glucagon load in Japanese patients with type 2 diabetes. 2013 , 4, 243-250	
1658	Analysis of clinical factors contributing to the postglucagon increment in C-peptide levels in Japanese patients with type 2 diabetes: comparison with basal C-peptide levels and 24-h urinary C-peptide excretion. 2013 , 4, 60-65	3
1657	A replication study for three nephrolithiasis loci at 5q35.3, 7p14.3 and 13q14.1 in the Japanese population. 2013 , 58, 588-93	23
1656	Comparison of the prevalence of convulsions associated with the use of cefepime and meropenem. 2013 , 35, 683-7	17
1655	Subclinical cerebral abnormalities in chronic kidney disease. 2013 , 179, 24-34	15
1654	Stroke feature and management in dialysis patients. 2013 , 179, 100-9	7
1653	Baseline characteristics and prevalence of cardiovascular disease in newly visiting or referred chronic kidney disease patients to nephrology centers in Japan: a prospective cohort study. 2013 , 14, 152	19
1652	Validation of a newly proposed histopathological classification in Japanese patients with anti-neutrophil cytoplasmic antibody-associated glomerulonephritis. 2013 , 14, 125	46
1651	Combined effect of mitochondrial DNA 5178 C/A polymorphism and alcohol consumption on estimated glomerular filtration rate in male Japanese health check-up examinees: a cross-sectional study. 2013 , 14, 35	12
1650	Impact of the clinical syntax score on 5-year clinical outcomes after sirolimus-eluting stents implantation. 2013 , 28, 258-66	7
1649	Role of arterial stiffness and impaired renal function in the progression of new coronary lesions after percutaneous coronary intervention. 2013 , 28, 56-62	16
1648	Influencing factors on cardiac structure and function beyond glycemic control in patients with type 2 diabetes mellitus. 2013 , 12, 38	29
1647	Remifentanyl temporarily improves renal function in adult patients with chronic kidney disease undergoing orthopedic surgery. 2013 , 27, 340-5	9
1646	Impaired renal function is a major determinant of left ventricular diastolic dysfunction: assessment by stress myocardial perfusion imaging. 2013 , 27, 729-36	6
1645	Iodine-123 metaiodobenzylguanidine imaging can predict future cardiac events in Japanese patients with Parkinson's disease. 2013 , 27, 123-31	8

1644	Distribution of and factors contributing to chronic kidney disease in a middle-aged working population. 2013 , 18, 466-76	4
1643	Japan Renal Biopsy Registry and Japan Kidney Disease Registry: Committee Report for 2009 and 2010. 2013 , 17, 155-73	88
1642	Multiple liver cyst infection caused by <i>Salmonella ajiobo</i> in autosomal dominant polycystic kidney disease. 2013 , 19, 530-3	2
1641	Performance of GFR equations in Japanese subjects. 2013 , 17, 352-8	40
1640	Association between AST-120 and abdominal aortic calcification in predialysis patients with chronic kidney disease. 2013 , 17, 365-71	30
1639	Effect of glomerular filtration rate and proteinuria on medical cost among screened subjects. 2013 , 17, 372-8	3
1638	Clinical features and long-term renal outcomes of Japanese patients with obesity-related glomerulopathy. 2013 , 17, 379-85	33
1637	Aliskiren reduces home blood pressure and albuminuria in patients with hypertensive nephrosclerosis. 2013 , 17, 386-95	3
1636	Anti-centromere antibody is an independent risk factor for chronic kidney disease in patients with primary biliary cirrhosis. 2013 , 17, 405-10	6
1635	Elevated levels of systemic pentraxin 3 are associated with thin-cap fibroatheroma in coronary culprit lesions: assessment by optical coherence tomography and intravascular ultrasound. 2013 , 6, 945-54	32
1634	Serum β_2 -microglobulin level is a useful indicator of disease activity and hemophagocytic syndrome complication in systemic lupus erythematosus and adult-onset Still's disease. 2013 , 32, 999-1005	16
1633	Efficacy of alogliptin, a dipeptidyl peptidase-4 inhibitor, on glucose parameters, the activity of the advanced glycation end product (AGE) - receptor for AGE (RAGE) axis and albuminuria in Japanese type 2 diabetes. 2013 , 29, 624-30	49
1632	Assessment of glomerular filtration rate in elderly patients with chronic kidney disease. 2013 , 45, 1475-82	17
1631	MTHFR, MTR and MTRR polymorphisms and risk of chronic kidney disease in Japanese: cross-sectional data from the J-MICC Study. 2013 , 45, 1613-20	8
1630	Acidic urine is associated with poor prognosis in patients with chronic heart failure. 2013 , 28, 735-41	9
1629	Abnormal P-wave terminal force in lead V1 is associated with cardiac death or hospitalization for heart failure in prior myocardial infarction. 2013 , 28, 690-5	32
1628	Prognostic impact of pulse pressure at admission on in-hospital outcome after primary percutaneous coronary intervention for acute myocardial infarction. 2013 , 28, 434-41	14
1627	Efficacy analysis of the renoprotective effects of aliskiren in hypertensive patients with chronic kidney disease. 2013 , 28, 442-52	6

1626	Prognostic value of highly sensitive troponin T on cardiac events in patients with chronic kidney disease not on dialysis. 2013 , 28, 473-9	15
1625	L/N-type calcium channel blocker cilnidipine reduces plasma aldosterone, albuminuria, and urinary liver-type fatty acid binding protein in patients with chronic kidney disease. 2013 , 28, 480-9	17
1624	Characterization of the effect of serum bilirubin concentrations on coronary endothelial function via measurement of high-sensitivity C-reactive protein and high-density lipoprotein cholesterol. 2013 , 28, 157-65	14
1623	Prognostic implications of chronic kidney disease and anemia after percutaneous coronary intervention in acute myocardial infarction patients. 2013 , 28, 19-26	31
1622	Efficacy and effects on lipid metabolism of combination treatment with losartan + hydrochlorothiazide versus losartan + amlodipine: a 48-week prospective, multicenter, randomized, open-label trial. 2013 , 35, 461-73	8
1621	Oral alendronate can suppress bone turnover but not fracture in kidney transplantation recipients with hyperparathyroidism and chronic kidney disease. 2013 , 31, 116-22	11
1620	Estimated glomerular filtration rate and proteinuria are associated with persistent form of atrial fibrillation: analysis in Japanese patients. 2013 , 61, 53-7	13
1619	Serum KL-6 concentrations are associated with molecular sizes and efflux behavior of KL-6/MUC1 in healthy subjects. 2013 , 424, 148-52	5
1618	Assessment of body composition using dry mass index and ratio of total body water to estimated volume based on bioelectrical impedance analysis in chronic kidney disease patients. 2013 , 23, 28-36	11
1617	20-Hour preprocedural hydration is not superior to 5-hour preprocedural hydration in the prevention of contrast-induced increases in serum creatinine and cystatin C. 2013 , 167, 2200-3	10
1616	Intra- and extracranial atherosclerotic disease in acute spontaneous intracerebral hemorrhage. 2013 , 332, 116-20	15
1615	High serum C1q-adiponectin/total adiponectin ratio correlates with coronary artery disease in Japanese type 2 diabetics. 2013 , 62, 578-85	15
1614	Glucose intolerance is associated with increased intimal-medial thickness of the carotid artery and increased pulse-wave velocity in renal transplant recipients. 2013 , 45, 1535-9	3
1613	Utility of cystatin C as a marker for the severity of aortic regurgitation in hypertensive patients. 2013 , 425, 259-64	
1612	Renal function and bone mineral density in community-dwelling elderly Japanese men: the Fujiwara-kyo Osteoporosis Risk in Men (FORMEN) Study. 2013 , 56, 61-6	14
1611	Tolvaptan reduces the risk of worsening renal function in patients with acute decompensated heart failure in high-risk population. 2013 , 61, 169-74	72
1610	Factors determining achievement of early postoperative cardiac rehabilitation goal in patients with or without preoperative kidney dysfunction undergoing isolated cardiac surgery. 2013 , 61, 299-303	15
1609	Prognostic value of B-type natriuretic peptide and its amino-terminal proBNP fragment for cardiovascular events with stratification by renal function. 2013 , 61, 410-6	35

1608	Impact of contrast-induced acute kidney injury on outcomes in patients with ST-segment elevation myocardial infarction undergoing primary percutaneous coronary intervention. 2013 , 14, 253-7	24
1607	Urinary free light chain is a potential biomarker for ISN/RPS class III/IV lupus nephritis. 2013 , 52, 2149-57	16
1606	Usefulness of HPLC assay for early detection of microalbuminuria in chronic kidney disease. 2013 , 27, 333-8	5
1605	Risk factors for increased left ventricular hypertrophy in patients with chronic kidney disease. 2013 , 17, 730-742	19
1604	Acute changes in home blood pressure after the Great East Japan Earthquake among patients with chronic kidney disease in Fukushima City. 2013 , 17, 718-724	13
1603	An inverted J-shaped association of serum uric acid with muscle strength among Japanese adult men: a cross-sectional study. 2013 , 14, 258	26
1602	The predictive value of attenuated proteinuria at 1 year after steroid therapy for renal survival in patients with IgA nephropathy. 2013 , 17, 555-62	21
1601	A retrospective cohort study of panipenem/betamipron for adult pneumococcal bacteremia at three teaching hospitals in Japan. 2013 , 19, 607-14	4
1600	Low contrast- and low radiation dose protocol for cardiac CT of thin adults at 256-row CT: usefulness of low tube voltage scans and the hybrid iterative reconstruction algorithm. 2013 , 29, 913-23	32
1599	Validation of the 2010 histopathological classification of ANCA-associated glomerulonephritis in a Japanese single-center cohort. 2013 , 1	4
1598	Comparison of the clinical utility of tocilizumab and anti-TNF therapy in AA amyloidosis complicating rheumatic diseases. 2013 , 1	1
1597	Impact of dietary fiber intake on glycemic control, cardiovascular risk factors and chronic kidney disease in Japanese patients with type 2 diabetes mellitus: the Fukuoka Diabetes Registry. 2013 , 12, 159	89
1596	Short-term functional and oncological outcomes of partial nephrectomy for renal cell carcinoma in patients with an anatomically or functionally solitary kidney: single-center experience. 2013 , 18, 1049-53	2
1595	Replication study for the association of 3 SNP loci identified in a genome-wide association study for diabetic nephropathy in European type 1 diabetes with diabetic nephropathy in Japanese patients with type 2 diabetes. 2013 , 17, 866-71	10
1594	Japan Diabetic Nephropathy Cohort Study: study design, methods, and implementation. 2013 , 17, 819-26	7
1593	Albuminuria is an independent predictor of all-cause and cardiovascular mortality in the Japanese population: the Takahata study. 2013 , 17, 805-10	30
1592	Serum level of soluble (pro)renin receptor is modulated in chronic kidney disease. 2013 , 17, 848-56	60
1591	Unexpected combined effects of NADH dehydrogenase subunit-2 237 Leu/Met polymorphism and green tea consumption on renal function in male Japanese health check-up examinees: a cross-sectional study. 2013 , 12, 17	3

1590	Gender difference in relationship between body mass index and development of chronic kidney disease. 2013 , 6, 463	16
1589	Long-term results of dose-dense paclitaxel and carboplatin versus conventional paclitaxel and carboplatin for treatment of advanced epithelial ovarian, fallopian tube, or primary peritoneal cancer (JGOG 3016): a randomised, controlled, open-label trial. 2013 , 14, 1020-6	325
1588	Relationship between estimated glomerular filtration rate, albuminuria, and oxidant status in the Japanese population. 2013 , 14, 191	13
1587	Ethnic differences in creatinine kinetics in a New Zealand end-stage kidney disease cohort. 2013 , 18, 222-8	1
1586	Moderate-intensity single exercise session does not induce renal damage. 2013 , 27, 177-80	12
1585	Impact of central venous pressure on cardiorenal interactions in adult patients with congenital heart disease after biventricular repair. 2013 , 8, 103-10	19
1584	Latent IgA deposition from donor kidneys does not affect transplant prognosis, irrespective of mesangial expansion. 2013 , 27 Suppl 26, 14-21	18
1583	Utility of cystatin C for renal function in amyotrophic lateral sclerosis. 2013 , 128, 386-90	15
1582	The association between chronic kidney disease and cardiovascular disease risk factors in atomic bomb survivors. 2013 , 179, 46-52	38
1581	Intracranial deep white matter lesions (DWLs) are associated with chronic kidney disease (CKD) and cognitive impairment: a 5-year follow-up magnetic resonance imaging (MRI) study. 2013 , 56, 55-60	28
1580	Hydroxy-itraconazole pharmacokinetics is similar to that of itraconazole in immunocompromised patients receiving oral solution of itraconazole. 2013 , 415, 128-32	14
1579	Chronic renal outcome after living donor liver transplantation. 2013 , 27, 90-7	10
1578	Morbidity and mortality among newly hospitalized patients with community-acquired pneumococcal bacteremia: a retrospective cohort study in three teaching hospitals in Japan. 2013 , 13, 607-15	7
1577	Renal transporters in drug development. 2013 , 53, 503-29	226
1576	Clinical practice guidelines for nursing- and healthcare-associated pneumonia (NHCAP) [complete translation]. 2013 , 51, 103-26	108
1575	Intravoxel incoherent motion imaging of the kidney: alterations in diffusion and perfusion in patients with renal dysfunction. 2013 , 31, 414-7	76
1574	Frequency and implication of ST-T abnormalities on hospital admission electrocardiograms in patients with type A acute aortic dissection. 2013 , 112, 424-9	24
1573	Carotid intima-media thickness progression predicts cardiovascular events in Japanese patients with type 2 diabetes. 2013 , 101, 286-92	17

1572	Clinical features associated with a rapid decline in renal function among Japanese patients with type 2 diabetes mellitus: microscopic hematuria coexisting with diabetic retinopathy. 2013 , 100, e39-41		3
1571	Influence of daily alcohol consumption on serum adiponectin levels in men. 2013 , 62, 411-6		11
1570	Concordance of CKD stages in estimation by the CKD-EPI equation and estimation by the MDRD equation in the Japanese general population: the Iwate KENCO Study. 2013 , 165, 377-9		4
1569	Statins and the risks of stroke recurrence and death after ischemic stroke: the Fukuoka Stroke Registry. 2013 , 231, 211-5		11
1568	A comparison of the performances of an artificial neural network and a regression model for GFR estimation. <i>American Journal of Kidney Diseases</i> , 2013 , 62, 1109-15	7-4	17
1567	Clinical impact of coronary artery spasm in patients with no significant coronary stenosis who are experiencing acute coronary syndrome. 2013 , 61, 404-9		12
1566	Obesity paradox in Japanese patients after percutaneous coronary intervention: an observation cohort study. 2013 , 62, 18-24		45
1565	Relationship between BNP level and renal function in diabetic nephropathy with microalbuminuria. 2013 , 27, 92-7		12
1564	Bleeding events and activated partial thromboplastin time with dabigatran in clinical practice. 2013 , 62, 121-6		23
1563	Nighttime blood pressure, nighttime glucose values, and target-organ damages in treated type 2 diabetes patients. 2013 , 227, 135-9		6
1562	Endothelial dysfunction measured by peripheral arterial tonometry predicts prognosis in patients with heart failure with preserved ejection fraction. 2013 , 168, 36-40		47
1561	Change in high-sensitive cardiac troponin T on hypertensive treatment. 2013 , 35, 40-4		9
1560	Comparative efficacy and adverse effects of the addition of ezetimibe to statin versus statin titration in chronic kidney disease patients. 2013 , 7, 306-15		16
1559	Combination therapy for hypertension in patients with CKD: a subanalysis of the Combination Therapy of Hypertension to Prevent Cardiovascular Events trial. 2013 , 36, 947-58		22
1558	Diabetic retinopathy and microalbuminuria can predict macroalbuminuria and renal function decline in Japanese type 2 diabetic patients: Japan Diabetes Complications Study. 2013 , 36, 2803-9		35
1557	Relation of estimated glomerular filtration rate and body mass index to mortality in non-dialysis patients with coronary artery disease: a report from the ET-CHD registry, 1997-2003. 2013 , 62, 144-50		3
1556	Clinical characteristics and long-term clinical outcomes of Japanese heart failure patients with preserved versus reduced left ventricular ejection fraction: a prospective cohort of Shinken Database 2004-2011. 2013 , 62, 102-9		30
1555	Indoxyl sulfate, a uremic toxin, and carotid intima-media thickness in patients with coronary artery disease. 2013 , 163, 214-6		19

1554	Relation of plasma indoxyl sulfate levels and estimated glomerular filtration rate to left ventricular diastolic dysfunction. 2013 , 111, 712-6	43
1553	Japanese patients with Fabry disease predominantly showing cardiac and neurological manifestation with novel missense mutation: R220P. 2013 , 62, 63-9	3
1552	Correlation between serum C1q-adiponectin/total adiponectin ratio and polyvascular lesions detected by vascular ultrasonography in Japanese type 2 diabetics. 2013 , 62, 376-85	16
1551	Kidney stone formation is positively associated with conventional risk factors for coronary heart disease in Japanese men. 2013 , 189, 1340-6	30
1550	Age-relevant renal effects of cadmium exposure through consumption of home-harvested rice in female Japanese farmers. 2013 , 56, 1-9	50
1549	Atrial fibrillation-induced endothelial dysfunction improves after restoration of sinus rhythm. 2013 , 168, 1280-5	33
1548	Serum levels of advanced glycation end products (AGEs) are independently correlated with circulating levels of dipeptidyl peptidase-4 (DPP-4) in humans. 2013 , 46, 300-3	31
1547	Prognostic implications of anemia with or without chronic kidney disease in patients undergoing elective percutaneous coronary intervention. 2013 , 168, 5221-8	27
1546	Enhancement of ascites during abdominal oncological endovascular intervention without iatrogenic extravasation in patients with chronic liver disease. 2013 , 68, e460-6	1
1545	Serum uric acid in relation to serum 1,5-anhydroglucitol levels in patients with and without type 2 diabetes mellitus. 2013 , 46, 1436-41	10
1544	The cardio-ankle vascular index predicts chronic kidney disease in Japanese subjects. 2013 , 7, 48	6
1543	A preliminary study of the potential role of FGF-23 in coronary calcification in patients with suspected coronary artery disease. 2013 , 226, 228-33	26
1542	Dipstick proteinuria as a surrogate marker of long-term mortality after acute myocardial infarction. 2013 , 62, 277-82	11
1541	GFR estimation using standardized serum cystatin C in Japan. <i>American Journal of Kidney Diseases</i> , 2013 , 61, 197-203	7.4 162
1540	Low-dose abdominal CT: comparison of low tube voltage with moderate-level iterative reconstruction and standard tube voltage, low tube current with high-level iterative reconstruction. 2013 , 68, 1008-15	13
1539	Recurrent proliferative glomerulonephritis with monoclonal IgG deposits of IgG2 κ subtype in a transplanted kidney: a case report. <i>American Journal of Kidney Diseases</i> , 2013 , 62, 587-90	7.4 17
1538	The effects of continuing and discontinuing smoking on the development of chronic kidney disease (CKD) in the healthy middle-aged working population in Japan. 2013 , 18, 24-32	14
1537	Chapter 1: Definition and classification of CKD. 2013 , 3, 19-62	565

1536	Fibroblast growth factor 23 is associated with carotid artery calcification in chronic kidney disease patients not undergoing dialysis: a cross-sectional study. 2013 , 14, 22	26
1535	Serum galectin-9 levels are elevated in the patients with type 2 diabetes and chronic kidney disease. 2013 , 14, 23	30
1534	Patients with adult minimal change nephrotic syndrome treated with long-term cyclosporine did not experience a reduction in their eGFR. 2013 , 79, 101-6	2
1533	References. 2013 , 3, 136-150	66
1532	Insulin resistance and insulin secretion in renal transplant recipients with hepatitis C. 2013 , 45, 1540-3	3
1531	Systemic inflammatory response syndrome criteria and vancomycin dose requirement in patients with sepsis. 2013 , 39, 1247-52	58
1530	Comparison of effects of angiotensin II receptor blocker on morning home blood pressure and cardiorenal protection between morning administration and evening administration in hypertensive patients: the COMPATIBLE study. 2013 , 36, 202-7	15
1529	Chronic kidney disease: global dimension and perspectives. 2013 , 382, 260-72	2280
1528	An association between uric acid levels and renal arteriolopathy in chronic kidney disease: a biopsy-based study. 2013 , 36, 43-9	93
1527	Association between cardiopulmonary exercise and dobutamine stress testing in ambulatory patients with idiopathic dilated cardiomyopathy: a comparison with peak VO ₂ and VE/VCO ₂ slope. 2013 , 162, 234-9	8
1526	Fluctuation of serum C3 levels reflects disease activity and metabolic background in patients with IgA nephropathy. 2013 , 26, 708-15	16
1525	Association of nocturnal hypoxemia with progression of CKD. 2013 , 8, 1502-7	43
1524	Left ventricular reverse remodeling in long-term (>12 years) survivors with idiopathic dilated cardiomyopathy. 2013 , 111, 106-10	26
1523	Current status of the treatment of microscopic polyangiitis and granulomatosis with polyangiitis in Japan. 2013 , 17, 51-8	29
1522	Effects of atorvastatin on renal function in patients with dyslipidemia and chronic kidney disease: rationale and design of the ASsessment of clinical Usefulness in CKD patients with Atorvastatin (ASUCA) trial. 2013 , 17, 211-7	11
1521	Decreased physical function in pre-dialysis patients with chronic kidney disease. 2013 , 17, 225-31	70
1520	Prognostic significance of left ventricular hypertrophy observed at dialysis initiation depends on the pre-dialysis use of erythropoiesis-stimulating agents. 2013 , 17, 294-303	7
1519	Development of enzyme-linked immunosorbent assays for urinary thiazide-sensitive Na-Cl cotransporter measurement. 2013 , 305, F1374-81	30

1518	Which is a better treatment for hypertensive patients with diabetes: a combination of losartan and hydrochlorothiazide or a maximum dose of losartan?. 2013 , 35, 582-8	2
1517	Chronic kidney disease (CKD) is an independent risk factor for long-term care insurance (LTCI) need certification among older Japanese adults: a two-year prospective cohort study. 2013 , 57, 328-32	21
1516	The impact of cigarette smoking on 24-hour blood pressure, inflammatory and hemostatic activity, and cardiovascular risk in Japanese hypertensive patients. 2013 , 15, 234-40	10
1515	Predictive value of lipoprotein indices for residual risk of acute myocardial infarction and sudden death in men with low-density lipoprotein cholesterol levels . 2013 , 112, 1063-8	9
1514	Prediction of cardiovascular disease mortality by proteinuria and reduced kidney function: pooled analysis of 39,000 individuals from 7 cohort studies in Japan. 2013 , 178, 1-11	45
1513	Relationship between normal serum creatinine concentration and periodontal disease in Japanese middle-aged males. 2013 , 84, 94-9	5
1512	Impact of chronic kidney disease on carotid atherosclerosis according to blood pressure category: the Suita study. 2013 , 44, 3537-9	26
1511	Identification of chromosome 3q28 and ALPK1 as susceptibility loci for chronic kidney disease in Japanese individuals by a genome-wide association study. 2013 , 50, 410-8	41
1510	Thyroid hormone replacement therapy for primary hypothyroidism leads to significant improvement of renal function in chronic kidney disease patients. 2013 , 17, 525-31	27
1509	Effect of allopurinol on cardiovascular incidence among hypertensive nephropathy patients: the Gonryo study. 2013 , 17, 549-53	28
1508	The association between serum uric acid and renal damage in a community-based population: the Takahata study. 2013 , 17, 541-8	16
1507	Comparison of the loss of renal function after cold ischemia open partial nephrectomy, warm ischemia laparoscopic partial nephrectomy and laparoscopic partial nephrectomy using microwave coagulation. 2013 , 6, 118-23	2
1506	Independent association between low serum amylase and non-alcoholic fatty liver disease in asymptomatic adults: a cross-sectional observational study. 2013 , 3,	13
1505	Hyperuricemia is independently associated with endothelial dysfunction in postmenopausal women but not in premenopausal women. 2013 , 3, e003659	29
1504	Kidney-protective effects of azelnidipine versus a diuretic in combination with olmesartan in hypertensive patients with diabetes and albuminuria: a randomized study. 2013 , 28, 1802-10	5
1503	Evaluation of the Chronic Kidney Disease Epidemiology Collaboration equation for estimating glomerular filtration rate in the Chinese population. 2013 , 28, 641-51	119
1502	Predictive role of C reactive protein in stroke recurrence after cardioembolic stroke: the Fukuoka Stroke Registry. 2013 , 3, e003678	11
1501	Polymorphisms in the organic anion transporting polypeptide genes influence liver parenchymal enhancement in gadoteric acid-enhanced MRI. 2013 , 14, 1573-82	7

1500	The significant impact of acute kidney injury on CKD in patients who survived over 10 years after myeloablative allogeneic SCT. 2013 , 48, 80-4	30
1499	New-onset microalbuminuria following allogeneic myeloablative SCT is a sign of near-term decrease in renal function. 2013 , 48, 972-6	5
1498	Polymorphisms in PPAR Genes (PPARD, PPARG, and PPARGC1A) and the Risk of Chronic Kidney Disease in Japanese: Cross-Sectional Data from the J-MICC Study. 2013 , 2013, 980471	6
1497	Increased serum leukocyte cell-derived chemotaxin 2 (LECT2) levels in obesity and fatty liver. 2013 ,	4
1496	Predictive effects of urinary liver-type fatty acid-binding protein for deteriorating renal function and incidence of cardiovascular disease in type 2 diabetic patients without advanced nephropathy. 2013 , 36, 1248-53	71
1495	Oxidative stress/angiotensinogen/renin-angiotensin system axis in patients with diabetic nephropathy. 2013 , 14, 23045-62	51
1494	L/N-type calcium channel blocker cilnidipine added to renin-angiotensin inhibition improves ambulatory blood pressure profile and suppresses cardiac hypertrophy in hypertension with chronic kidney disease. 2013 , 14, 16866-81	11
1493	Prediction of glomerular filtration rate in cancer patients by an equation for Japanese estimated glomerular filtration rate. 2013 , 43, 271-7	14
1492	Association of a polymorphism of with chronic kidney disease in community-dwelling individuals. 2013 , 1, 868-872	7
1491	Addition of aliskiren to Angiotensin receptor blocker improves ambulatory blood pressure profile and cardiorenal function better than addition of benazepril in chronic kidney disease. 2013 , 14, 15361-75	4
1490	Past Obesity as well as Present Body Weight Status Is a Risk Factor for Diabetic Nephropathy. 2013 , 2013, 590569	10
1489	Night-time blood pressure is associated with the development of chronic kidney disease in a general population: the Ohasama Study. 2013 , 31, 2410-7	29
1488	Long-term outcomes of Japanese type 2 diabetic patients with biopsy-proven diabetic nephropathy. 2013 , 36, 3655-62	101
1487	Renal glucose handling: impact of chronic kidney disease and sodium-glucose cotransporter 2 inhibition in patients with type 2 diabetes. 2013 , 36, 1260-5	60
1486	High level of plasma remnant-like particle cholesterol may predispose to development of hypertension in normotensive subjects. 2013 , 26, 793-8	6
1485	Neutrophil extracellular trap components in fibrinoid necrosis of the kidney with myeloperoxidase-ANCA-associated vasculitis. 2013 , 6, 308-12	44
1484	Early evolution and correlates of urine albumin excretion in patients presenting with acutely decompensated heart failure. 2013 , 6, 227-32	23
1483	Clinical correlates of ambulatory BP monitoring among patients with CKD. 2013 , 8, 721-30	46

1482	Preoperative SYNTAX score and graft patency after off-pump coronary bypass surgery. 2013 , 44, e25-31	5
1481	Relationship between coffee consumption, oxidant status, and antioxidant potential in the Japanese general population. 2013 , 51, 1951-9	15
1480	Effects of obstructive sleep apnea and its treatment on signal-averaged P-wave duration in men. 2013 , 6, 287-93	24
1479	Relationship between retinal blood flow and renal function in patients with type 2 diabetes and chronic kidney disease. 2013 , 36, 957-61	32
1478	Elevated Serum Gamma-Glutamyltransferase (GGT) Activity and the Development of Chronic Kidney Disease (CKD) in Cigarette Smokers. 2013 , 5, 967-73	6
1477	Impact of vancomycin or linezolid therapy on development of renal dysfunction and thrombocytopenia in Japanese patients. 2013 , 59, 319-24	15
1476	Skin autofluorescence reflects integration of past long-term glycemic control in patients with type 1 diabetes. 2013 , 36, 2339-45	42
1475	Prognostic factors for metastatic urothelial carcinoma undergoing cisplatin-based salvage chemotherapy. 2013 , 43, 923-8	21
1474	The RENAL nephrometry score and the PADUA classification for the prediction of perioperative outcomes in patients receiving nephron-sparing surgery: feasible tools to predict intraoperative conversion to nephrectomy. 2013 , 91, 261-8	25
1473	Development and validation of a prediction rule using the Oxford classification in IgA nephropathy. 2013 , 8, 2082-90	74
1472	S100A12 gene expression is increased in peripheral leukocytes in chronic kidney disease stage 4-5 patients with cardiovascular disease. 2013 , 123, 202-8	8
1471	The impact of renal tubular damage, as assessed by urinary β_2 -microglobulin-creatinine ratio, on cardiac prognosis in patients with chronic heart failure. 2013 , 6, 662-8	18
1470	Relationship between urinary fractional excretion of sodium and life prognosis in liver cirrhosis patients. 2013 , 43, 1156-62	8
1469	Thromboembolic complications after endovascular repair of abdominal aortic aneurysm with neck thrombus. 2013 , 47, 172-8	3
1468	Population pharmacokinetic analysis of linezolid in low body weight patients with renal dysfunction. 2013 , 53, 967-73	30
1467	Adalimumab treatment in patients with rheumatoid arthritis with renal insufficiency. 2013 , 65, 471-5	18
1466	Risk factors for chronic kidney disease after chemotherapy for testicular cancer. 2013 , 20, 716-22	9
1465	Efficacy of mizoribine followed by low-dose prednisone in patients with idiopathic membranous nephropathy and nephrotic-range proteinuria. 2013 , 35, 936-41	3

1464	Preoperative masked renal damage in Japanese patients with primary aldosteronism: identification of predictors for chronic kidney disease manifested after adrenalectomy. 2013 , 20, 685-91	22
1463	Efficacy of N-terminal pro-brain natriuretic peptide digit number for screening of cardiac disease in new haemodialysis patients. 2013 , 18, 497-504	5
1462	Clinical significance of microscopic haematuria in diabetic nephropathy in type 2 diabetes patients with overt proteinuria. 2013 , 18, 563-8	17
1461	Serum cystatin C as a biomarker of cardiac diastolic dysfunction in patients with cardiac disease and preserved ejection fraction. 2013 , 19, E35-9	10
1460	Distribution of glomerular density in different cortical zones of the human kidney. 2013 , 63, 169-75	22
1459	Creatinine/cystatinC ratio as a surrogate marker of residual muscle mass in amyotrophic lateral sclerosis. 2013 , 1, 32-37	34
1458	B-type natriuretic peptide levels are decreased by reducing dietary salt intake in patients with compensated heart failure with preserved ejection fraction. 2013 , 43, 663-7	9
1457	The impact of predisposing factors on long-term outcome after stroke in diabetic patients: the Fukuoka Stroke Registry. 2013 , 20, 921-7	17
1456	Comparison of spironolactone and trichlormethiazide as add-on therapy to renin-angiotensin blockade for reduction of albuminuria in diabetic patients. 2013 , 4, 316-9	11
1455	Defining criteria for the introduction of liraglutide using the glucagon stimulation test in patients with type2 diabetes. 2013 , 4, 571-5	14
1454	Effects of valsartan on progression of kidney disease in Japanese hypertensive patients with advanced, predialysis, chronic kidney disease: Kanagawa Valsartan Trial (KVT). 2013 , 36, 240-6	10
1453	Independent association of elevated serum hepatocyte growth factor levels with development of insulin resistance in a 10-year prospective study. 2013 , 79, 43-8	15
1452	Translocation of dendrin to the podocyte nucleus in acute glomerular injury in patients with IgA nephropathy. 2013 , 28, 1762-72	15
1451	Assessment and management of renal impairment in chemotherapy for urogenital cancer. 2013 , 43, 1055-63	2
1450	Aging and chronic kidney disease. 2013 , 38, 109-20	69
1449	Effect of tablets with a combination of telmisartan and amlodipine on patients with hypertension: the Cotalo study. 2013 , 36, 620-6	11
1448	Differences between daytime and nighttime blood pressure variability regarding systemic atherosclerotic change and renal function. 2013 , 36, 232-9	34
1447	Carotid plaque score and intima media thickness as predictors of stroke and mortality in hypertensive patients. 2013 , 36, 902-9	25

1446	Fibroblast growth factor 23 and 25-hydroxyvitamin D levels are associated with estimated glomerular filtration rate decline. 2013 , 3, 469-475	18
1445	The angiotensin II type 1 receptor blocker olmesartan preferentially improves nocturnal hypertension and proteinuria in chronic kidney disease. 2013 , 36, 262-9	20
1444	Chronic kidney disease is associated with neovascularization and intraplaque hemorrhage in coronary atherosclerosis in elders: results from the Hisayama Study. 2013 , 84, 373-80	26
1443	A combination of healthy lifestyle factors is associated with a decreased incidence of chronic kidney disease: a population-based cohort study. 2013 , 36, 328-33	45
1442	Impact of visit-to-visit variability of blood pressure on deterioration of renal function in patients with non-diabetic chronic kidney disease. 2013 , 36, 151-7	59
1441	Long-term compliance with salt restriction assessed using the spot urine method in Japanese cardiology outpatients. 2013 , 36, 1096-9	6
1440	Association between dietary behavior and risk of hypertension among Japanese male workers. 2013 , 36, 374-80	11
1439	Effects of dual blockade of the renin-angiotensin system on renal and cardiovascular outcomes in type 2 diabetes with overt nephropathy and hypertension in the ORIENT: a post-hoc analysis (ORIENT-Hypertension). 2013 , 36, 1051-9	22
1438	A community-based cross-sectional and longitudinal study uncovered asymptomatic proteinuria in Japanese adults with low body weight. 2013 , 84, 1254-61	13
1437	High salt intake promotes a decline in renal function in hypertensive patients: a 10-year observational study. 2013 , 36, 172-6	30
1436	Distribution of central blood pressure values estimated by Omron HEM-9000AI in the Japanese general population. 2013 , 36, 50-7	32
1435	Changes in serum uric acid have a reciprocal effect on eGFR change: a 10-year follow-up study of community-based screening in Okinawa, Japan. 2013 , 36, 650-4	34
1434	Association between hypertension status and the screening test for frailty in elderly community-dwelling Japanese. 2013 , 36, 639-44	16
1433	Mycophenolate mofetil versus intravenous cyclophosphamide for induction treatment of proliferative lupus nephritis in a Japanese population: a retrospective study. 2013 , 23, 89-96	8
1432	Presence of diastolic dysfunction in patients with peripheral artery disease. 2013 , 64, 540-3	10
1431	Association between remission of macroalbuminuria and preservation of renal function in patients with type 2 diabetes with overt proteinuria. 2013 , 36, 3227-33	36
1430	An angiotensin II receptor blocker-calcium channel blocker combination prevents cardiovascular events in elderly high-risk hypertensive patients with chronic kidney disease better than high-dose angiotensin II receptor blockade alone. 2013 , 83, 167-76	39
1429	Low salt intake and changes in serum sodium levels in the combination therapy of low-dose hydrochlorothiazide and angiotensin II receptor blocker. 2013 , 77, 2567-72	5

1428	High mean corpuscular volume is a new indicator of prognosis in acute decompensated heart failure. 2013 , 77, 2766-71	32
1427	Comparison of predictability of future cardiovascular events between chronic kidney disease (CKD) stage based on CKD epidemiology collaboration equation and that based on modification of diet in renal disease equation in the Japanese general population--Iwate KENCO Study. 2013 , 77, 1315-25	23
1426	Dipeptidyl peptidase-4 inhibitor, sitagliptin, improves endothelial dysfunction in association with its anti-inflammatory effects in patients with coronary artery disease and uncontrolled diabetes. 2013 , 77, 1337-44	157
1425	Trend of clinical outcome and surrogate markers during titration of β -blocker in heart failure patients with reduced ejection fraction: relevance of achieved heart rate and β -blocker dose. 2013 , 77, 1001-8	18
1424	Angiotensin-like protein 2 is associated with chronic kidney disease in a general Japanese population: the Hisayama Study. 2013 , 77, 2311-7	28
1423	Comparison of febuxostat and allopurinol for hyperuricemia in cardiac surgery patients (NU-FLASH Trial). 2013 , 77, 2043-9	77
1422	Impact of adiponectin and leptin on long-term adverse events in Japanese patients with acute myocardial infarction. Results from the Nagoya Acute Myocardial Infarction Study (NAMIS). 2013 , 77, 2778-85	18
1421	Effects of smoking habit on the prevalence of atrial fibrillation in Japanese patients with special reference to sex differences. 2013 , 77, 2948-53	14
1420	Association of serum YKL-40 levels with urinary albumin excretion rate in young Japanese patients with type 1 diabetes mellitus. 2013 , 60, 73-9	22
1419	Association of myeloperoxidase G-463A gene polymorphism with diabetic nephropathy in Japanese type 2 diabetic subjects. 2013 , 60, 457-471	7
1418	Relationship between metformin use, vitamin B12 deficiency, hyperhomocysteinemia and vascular complications in patients with type 2 diabetes. 2013 , 60, 1275-80	42
1417	Association between beta cell function and future glycemic control in patients with type 2 diabetes. 2013 , 60, 517-523	24
1416	Acute kidney injury in a patient with nephrotic syndrome due to focal segmental glomerular nephritis induced by a single oral administration of high-dose bisphosphonate (minodronate). 2013 , 52, 1383-7	4
1415	Increased uric acid promotes decline of the renal function in hypertensive patients: a 10-year observational study. 2013 , 52, 1467-72	12
1414	Determination of the B-type natriuretic peptide level as a criterion for abnormalities in Japanese individuals in routine clinical practice: the J-ABS Multi-Center Study (Japan Abnormal BNP Standard). 2013 , 52, 171-7	25
1413	The clinical applicability of albuminuria testing in Japanese hypertensive patients: the AVA-E study. 2013 , 52, 425-30	1
1412	Abacavir/lamivudine versus tenofovir/emtricitabine with atazanavir/ritonavir for treatment-naive Japanese patients with HIV-1 infection: a randomized multicenter trial. 2013 , 52, 735-44	12
1411	Mild renal dysfunction on admission is an important prognostic predictor in patients with infective endocarditis: a retrospective single-center study. 2013 , 52, 1013-8	8

1410	Association between indoxyl sulfate and cardiac dysfunction and prognosis in patients with dilated cardiomyopathy. 2013 , 77, 390-6	47
1409	[Prevalence of chronic kidney disease among HIV-infected individuals in Japan--a report from two tertiary hospitals]. 2013 , 87, 14-21	3
1408	Renal interstitial fibrosis in 0-hour biopsy as a predictor of post-transplant anemia. 2013 , 38, 267-74	11
1407	Home-based cardiac rehabilitation decreases red cell distribution width in chronic heart failure. 2013 , 68, 615-9	8
1406	Association of chronic kidney disease and cerebral small vessel disease with cognitive impairment in elderly patients with type 2 diabetes. 2013 , 3, 212-22	23
1405	Relation of estimated glomerular filtration rate to the presence of coronary plaque and obstructive coronary artery disease in a zero or low coronary artery calcium score. 2013 , 126, 153-8	1
1404	Alteration of vascular function is an important factor in the correlation between visit-to-visit blood pressure variability and cardiovascular disease. 2013 , 31, 1387-95; discussion 1395	33
1403	Prevalence and risk factors for loss of bone mineral density in male Japanese patients with HIV. 2013 , 62, e25-7	2
1402	The risk factors of severe acute kidney injury induced by cisplatin. 2013 , 85, 364-9	36
1401	Low-contrast dose protection protocol for diagnostic computed tomography in patients at high-risk for contrast-induced nephropathy. 2013 , 37, 289-96	6
1400	Combination of antihypertensive therapy in the elderly, multicenter investigation (CAMUI) trial: results after 1 year. 2013 , 31, 1245-55	10
1399	Limited renal prophylaxis in regular plasmapheresis for heritable ADAMTS13 deficiency. 2013 , 60, 1557-8	2
1398	Diabetes with preserved renal function is an independent risk factor for renal function deterioration after coronary computed tomography angiography. 2013 , 37, 750-4	9
1397	High-normal diastolic blood pressure is a risk for development of microalbuminuria in the general population: the Watari study. 2013 , 31, 798-804	11
1396	Systemic risk factors associated with polypoidal choroidal vasculopathy and neovascular age-related macular degeneration. 2013 , 33, 841-5	33
1395	[Diabetes mellitus related common medical disorders recent progress in diagnosis and treatment. Topics: I. Pathophysiology, diagnosis and treatment; 7. Chronic kidney disease.]. 2013 , 102, 875-81	
1394	Long-term outcome and efficacy of cyclophosphamide therapy in Japanese patients with ANCA-associated microscopic polyangiitis: a retrospective study. 2013 , 52, 2503-9	7
1393	Atrial fibrillation (AF) and complete right bundle branch block (RBBB) are independently associated with increased hemoglobin levels in apparently healthy subjects. 2013 , 52, 37-43	2

- 1392 An elevated ratio of placental growth factor to soluble fms-like tyrosine kinase-1 predicts adverse outcomes in patients with stable coronary artery disease. **2013**, 52, 1019-27 17
- 1391 [New era of laboratory testing. Topics: II. Particulars; 3. Laboratory examination for the diagnosis of CKD or AKI]. **2013**, 102, 3125-32
- 1390 Association of serum carotenoid levels with N-terminal pro-brain-type natriuretic peptide: a cross-sectional study in Japan. **2013**, 23, 163-8 8
- 1389 Post-stress perfusion abnormalities detected on myocardial perfusion single-photon emission computed tomography predict long-term mortality after elective abdominal aortic aneurysm repair. **2013**, 77, 1229-34 1
- 1388 Comparison of once daily versus twice daily olmesartan in patients with chronic kidney disease. **2013**, 6, 223-7 6
- 1387 Comparison of the antialbuminuric effects of L-/N-type and L-type calcium channel blockers in hypertensive patients with diabetes and microalbuminuria: the study of assessment for kidney function by urinary microalbumin in randomized (SAKURA) trial. **2013**, 10, 1209-16 16
- 1386 Estimating glomerular filtration rates in elderly Chinese patients with chronic kidney disease: performance of six modified formulae developed in Asian populations. **2013**, 8, 899-904 4
- 1385 Renal insufficiency without albuminuria is associated with peripheral artery atherosclerosis and lipid metabolism disorders in patients with type 2 diabetes. **2013**, 20, 790-7 12
- 1384 Polymorphisms in genes encoding antioxidant enzymes (SOD2, CAT, GPx, TXNRD, SEPP1, SEP15 and SELS) and risk of chronic kidney disease in Japanese - cross-sectional data from the J-MICC study. **2013**, 53, 15-20 10
- 1383 Renal dysfunction on admission as a predictor for in-hospital mortality of patients with stanford type B acute aortic dissection. **2013**, 6, 624-30 5
- 1382 Adjusted Anion Gap Is Associated with Glomerular Filtration Rate Decline in Chronic Kidney Disease. **2013**, 3, 113-117 4
- 1381 Association of low serum iron levels with low-grade inflammation and hyperadiponectinemia in community-living elderly women. **2013**, 20, 670-7 7
- 1380 Renal dysfunction and atherosclerosis of the neointima following bare metal stent implantation. **2013**, 38, 58-65 6
- 1379 Carotid artery plaque and LDL-to-HDL cholesterol ratio predict atherosclerotic status in coronary arteries in asymptomatic patients with type 2 diabetes mellitus. **2013**, 20, 452-64 29
- 1378 Glucagon-like peptide-1 secretory function as an independent determinant of blood pressure: analysis in the Tanno-Sobetsu study. **2013**, 8, e67578 11
- 1377 Association between circulating fibroblast growth factor 23, Klotho, and the left ventricular ejection fraction and left ventricular mass in cardiology inpatients. **2013**, 8, e73184 42
- 1376 Serum uric acid is associated with left ventricular hypertrophy independent of serum parathyroid hormone in male cardiac patients. **2013**, 8, e82735 16
- 1375 Skin autofluorescence is associated with the progression of chronic kidney disease: a prospective observational study. **2013**, 8, e83799 29

1374	A decreased level of serum soluble Klotho is an independent biomarker associated with arterial stiffness in patients with chronic kidney disease. 2013 , 8, e56695	134
1373	Association between cystatin C and arteriosclerosis in the absence of chronic kidney disease. 2013 , 20, 548-56	18
1372	Development and psychometric properties of the Japanese heart failure knowledge scale. 2013 , 54, 228-33	14
1371	Prognosis of elderly Japanese patients aged 80 years undergoing hemodialysis. 2013 , 2013, 693514	9
1370	The Prevalence of the Risk Factors for Atherosclerosis among Type 2 Diabetic Patients Is Greater in the Progressive Stages of Chronic Kidney Disease. 2013 , 3, 66-72	14
1369	Recent characteristics and outcomes of Japanese stable angina pectoris after percutaneous coronary intervention. An observational cohort study using the Shinken Database. 2013 , 54, 335-40	10
1368	Differences in brain natriuretic peptide and other factors between Japanese peripheral arterial disease patients with critical limb ischemia and intermittent claudication. 2013 , 20, 798-806	12
1367	Therapeutic Strategies in Amyloid A Amyloidosis Secondary to Rheumatoid Arthritis. 2013 ,	
1366	Anemia and reduced renal function are independent predictors of elevated serum adiponectin in elderly women. 2013 , 20, 568-74	11
1365	Effects of adding ezetimibe to fluvastatin on kidney function in patients with hypercholesterolemia: a randomized control trial. 2013 , 20, 245-56	6
1364	Efficacy and safety of febuxostat in the treatment of hyperuricemia in stable kidney transplant recipients. 2014 , 8, 245-53	18
1363	Comparison of the antialbuminuric effects of benidipine and hydrochlorothiazide in Renin-Angiotensin System (RAS) inhibitor-treated hypertensive patients with albuminuria: the COSMO-CKD (COMbination Strategy on Renal Function of Benidipine or Diuretics TreatMent with RAS inhibitor) in Chronic Kidney Disease (Hypertensive Population) study. 2014 , 11, 887-894	7
1362	Efficacy and safety of febuxostat in elderly female patients. 2014 , 9, 1489-93	5
1361	Predictive significance of kidney myeloid-related protein 8 expression in patients with obesity- or type 2 diabetes-associated kidney diseases. 2014 , 9, e88942	6
1360	The total urine protein-to-creatinine ratio can predict the presence of microalbuminuria. 2014 , 9, e91067	13
1359	Prognosis in IgA nephropathy: 30-year analysis of 1,012 patients at a single center in Japan. 2014 , 9, e91756	117
1358	Predictive properties of plasma amino acid profile for cardiovascular disease in patients with type 2 diabetes. 2014 , 9, e101219	34
1357	Plasma brain-derived neurotrophic factor and reverse dipping pattern of nocturnal blood pressure in patients with cardiovascular risk factors. 2014 , 9, e105977	22

1356	Importance of fatty acid compositions in patients with peripheral arterial disease. 2014 , 9, e107003	13
1355	Association between combined lifestyle factors and non-restorative sleep in Japan: a cross-sectional study based on a Japanese health database. 2014 , 9, e108718	37
1354	Intravenous thrombolysis with recombinant tissue plasminogen activator for ischemic stroke patients over 80 years old: the Fukuoka Stroke Registry. 2014 , 9, e110444	13
1353	Urinary neutrophil gelatinase-associated lipocalin: a useful biomarker for tacrolimus-induced acute kidney injury in liver transplant patients. 2014 , 9, e110527	30
1352	Correlation between renal function and common risk factors for chronic kidney disease in a healthy middle-aged population: a prospective observational 2-year study. 2014 , 9, e113263	9
1351	Significance of urinary full-length megalin in patients with IgA nephropathy. 2014 , 9, e114400	18
1350	Urinary excretion of fatty acid-binding protein 4 is associated with albuminuria and renal dysfunction. 2014 , 9, e115429	23
1349	Chronic kidney disease screening methods and its implication for Malaysia: an in depth review. 2015 , 7, 96-109	8
1348	Waist to height ratio is an independent predictor for the incidence of chronic kidney disease. 2014 , 9, e88873	25
1347	Pulse-Wave Analysis of Optic Nerve Head Circulation Is Significantly Correlated with Kidney Function in Patients with and without Chronic Kidney Disease. 2014 , 2014, 291687	12
1346	Plasma pentraxin 3 is a more potent predictor of endothelial dysfunction than high-sensitive C-reactive protein. 2014 , 55, 160-4	40
1345	Prognostic impact of chronic kidney disease and anemia at admission on in-hospital outcomes after primary percutaneous coronary intervention for acute myocardial infarction. 2014 , 55, 301-6	28
1344	Application of creatinine- and/or cystatin C-based glomerular filtration rate estimation equations in elderly Chinese. 2014 , 9, 1539-49	9
1343	Comparison of Ten Creatinine-based Equations for Estimation of Glomerular Filtration Rate in Chinese Postmenopausal Women with Normal or Mildly Reduced Renal Function. 2014 , 03,	
1342	Evaluation of the Time to Terminate Continuous Renal Replacement Therapy in Patients with Acute Kidney Injury Following Cardiovascular Surgery in Association with or without a Mechanical Ventilation Support. 2014 , 05,	
1341	Entity of Stage 4 CKD over 75 Years Old is Different from that of Less than 75 Years of Age: Characteristics of Elderly Patients with Stage 4 CKD. 2014 , 04,	
1340	Risk Factors for Acute Kidney Injury Requiring Continuous Renal Replacement Therapy after Off-Pump Coronary Surgery. 2014 , 04,	
1339	Effects of the T/L-type calcium channel blocker benidipine on albuminuria and plasma aldosterone concentration. A pilot study involving switching from L-type calcium channel blockers to benidipine. 2014 , 55, 519-25	8

1338	Circulating TNF receptor 2 is closely associated with the kidney function in non-diabetic Japanese subjects. 2014 , 21, 730-8	7
1337	Comparison of the Framingham risk score, UK Prospective Diabetes Study (UKPDS) Risk Engine, Japanese Atherosclerosis Longitudinal Study-Existing Cohorts Combine (JALS-ECC) and maximum carotid intima-media thickness for predicting coronary artery stenosis in patients with asymptomatic type 2 diabetes. 2014 , 21, 799-815	23
1336	The asymmetric dimethylarginine level is associated with the predicted stroke risk in Japanese women. 2014 , 21, 640-7	3
1335	Kidney function, cholesterol absorption and remnant lipoprotein accumulation in patients with diabetes mellitus. 2014 , 21, 346-54	11
1334	Coagulation and Fibrinolysis Markers and Their Use for the Prediction of High Risk Patients with Venous Thromboembolism Following Total Hip Arthroplasty. 2014 ,	
1333	Serum apolipoprotein B-48 concentration is associated with a reduced estimated glomerular filtration rate and increased proteinuria. 2014 , 21, 974-82	7
1332	Impact of blood urea nitrogen for long-term risk stratification in patients with coronary artery disease undergoing percutaneous coronary intervention. 2014 , 4, 116-121	3
1331	Polymorphisms of genes involved in lipid metabolism and risk of chronic kidney disease in Japanese - cross-sectional data from the J-MICC study. 2014 , 13, 162	5
1330	Pharmacokinetic and pharmacodynamic profiles of canagliflozin in Japanese patients with type 2 diabetes mellitus and moderate renal impairment. 2014 , 34, 731-42	24
1329	Association of serum levels of FGF23 and Klotho with glomerular filtration rate and proteinuria among cardiac patients. 2014 , 15, 147	22
1328	Pathological similarities between low birth weight-related nephropathy and nephropathy associated with mitochondrial cytopathy. 2014 , 9, 181	7
1327	Evaluation of urinary tissue inhibitor of metalloproteinase-2 in acute kidney injury: a prospective observational study. 2014 , 18, 716	38
1326	High blood pressure after acute ischemic stroke is associated with poor clinical outcomes: Fukuoka Stroke Registry. 2014 , 63, 54-60	71
1325	Risk factors for acute kidney injury in aortic arch surgery with selective cerebral perfusion and mild hypothermic lower body circulatory arrest. 2014 , 19, 955-61	23
1324	Time Course of Calcium Concentrations and Risk Factors for Hypocalcemia in Patients Receiving Denosumab for the Treatment of Bone Metastases From Cancer. 2014 , 48, 1159-1165	20
1323	Association between blood pressure and disability-free survival among community-dwelling elderly patients receiving antihypertensive treatment. 2014 , 37, 772-8	9
1322	References. 2014 , 37, 362-387	37
1321	Comparison of renal ischemic damage during laparoscopic partial nephrectomy with artery-vein and artery-only clamping. 2014 , 28, 306-11	18

1320	Serum soluble (pro)renin receptor levels in patients with essential hypertension. 2014 , 37, 642-8	47
1319	Interrelationship between brachial artery function and renal small artery sclerosis in chronic kidney disease. 2014 , 37, 863-9	11
1318	Aldehyde dehydrogenase 2 as a potential protective factor for renal insufficiency in Japanese subjects with heart failure: a pilot study. 2014 , 28, 279-81	2
1317	Serum high-sensitivity C-reactive protein levels are associated with high risk of development, not progression, of diabetic nephropathy among Japanese type 2 diabetic patients: a prospective cohort study (Diabetes Distress and Care Registry at Tenri [DDCRT7]). 2014 , 37, 2947-52	21
1316	Ischemic diabetic retinopathy as a possible prognostic factor for chronic kidney disease progression. 2014 , 28, 1119-25	20
1315	Sex differences in the association between serum uric acid levels and cardiac hypertrophy in patients with chronic kidney disease. 2014 , 37, 246-52	31
1314	Postprandial glycemic control conditions in relation to urinary N-acetyl-β-D-glucosaminidase in patients with type 2 diabetes mellitus without low glomerular filtration rate. 2014 , 16, 41-7	9
1313	Hyperuricemia predicts future metabolic syndrome: a 4-year follow-up study of a large screened cohort in Okinawa, Japan. 2014 , 37, 232-8	30
1312	Milder clinical aspects of X-linked Alport syndrome in men positive for the collagen IV α chain. 2014 , 85, 1208-13	53
1311	Comparison of the clinical utility of tocilizumab and anti-TNF therapy in AA amyloidosis complicating rheumatic diseases. 2014 , 24, 137-43	41
1310	Modified glomerular filtration rate-estimating equations developed in asiatic population for chinese patients with type 2 diabetes. 2014 , 2014, 521071	13
1309	X-linked Alport syndrome caused by splicing mutations in COL4A5. 2014 , 9, 1958-64	26
1308	Risk factors for an early dialysis start in patients with diabetic nephropathy end-stage renal disease. 2014 , 10, 73-6	1
1307	Associations of anemia and renal dysfunction with outcomes among patients hospitalized for acute decompensated heart failure with preserved or reduced ejection fraction. 2014 , 9, 1912-21	12
1306	Telaprevir-based triple therapy for hepatitis C null responders among living donor liver transplant recipients. 2014 , 8, 339-45	6
1305	Validation of the 2010 histopathological classification of ANCA-associated glomerulonephritis in a Japanese single-center cohort. 2014 , 24, 300-3	21
1304	Plasma Antimicrobial Peptide LL-37 Level Is Inversely Associated with HDL Cholesterol Level in Patients with Type 2 Diabetes Mellitus. 2014 , 2014, 703696	6
1303	Acoustic radiation force impulse elastography of the kidneys: is shear wave velocity affected by tissue fibrosis or renal blood flow?. 2014 , 33, 793-801	79

1302	Effects of single pill-based combination therapy of amlodipine and atorvastatin on within-visit blood pressure variability and parameters of renal and vascular function in hypertensive patients with chronic kidney disease. 2014 , 2014, 437087	11
1301	Association between the Hemoglobin Level and Cardiothoracic Ratio in Patients on Incident Dialysis. 2014 , 4, 189-200	15
1300	Drinking pattern and risk of chronic kidney disease: the kansai healthcare study. 2014 , 40, 516-22	24
1299	The relationship between serum uric acid levels and β -cell functions in nondiabetic subjects. 2014 , 46, 950-4	8
1298	Selective pharmacological prophylaxis based on individual risk assessment using plasma levels of soluble fibrin and plasminogen-activator inhibitor-1 following total hip arthroplasty. 2014 , 24, 835-9	4
1297	Contrast material and radiation dose reduction strategy for triple-rule-out cardiac CT angiography: feasibility study of non-ECG-gated low kVp scan of the whole chest following coronary CT angiography. 2014 , 55, 1186-96	9
1296	Association of pulse pressure with serum TNF- α and neutrophil count in the elderly. 2014 , 2014, 972431	4
1295	Eldecalcitol improves chair-rising time in postmenopausal osteoporotic women treated with bisphosphonates. 2014 , 10, 51-9	13
1294	Osteoprotegerin is secreted into the coronary circulation: a possible association with the renin-angiotensin system and cardiac hypertrophy. 2014 , 46, 581-6	10
1293	Repulsive guidance cue semaphorin 3A in urine predicts the progression of acute kidney injury in adult patients from a mixed intensive care unit. 2014 , 29, 73-80	16
1292	Renal histological heterogeneity and functional progress in normoalbuminuric and microalbuminuric Japanese patients with type 2 diabetes. 2014 , 2, e000029	21
1291	Association between serum β -glutamyltranspeptidase and atherosclerosis: a population-based cross-sectional study. 2014 , 4, e005413	13
1290	Impact of albuminuria on early neurological deterioration and lesion volume expansion in lenticulostriate small infarcts. 2014 , 45, 587-90	17
1289	The increasing impact of a higher body mass index on the decrease in plasma B-type natriuretic peptide levels. 2014 , 4, 39-46	4
1288	Does increased water intake prevent disease progression in autosomal dominant polycystic kidney disease?. 2014 , 29, 1710-9	45
1287	ABCD3 and ABCD3-I scores are superior to ABCD2 score in the prediction of short- and long-term risks of stroke after transient ischemic attack. 2014 , 45, 418-25	68
1286	Renal prognosis a long time after renal biopsy on patients with diabetic nephropathy. 2014 , 29, 109-18	59
1285	A multicenter cross-sectional study of circulating soluble urokinase receptor in Japanese patients with glomerular disease. 2014 , 85, 641-8	66

1284	A multicenter randomized controlled trial of tonsillectomy combined with steroid pulse therapy in patients with immunoglobulin A nephropathy. 2014 , 29, 1546-53	120
1283	Pharmacokinetic and CYP3A5 pharmacogenetic differences between once- and twice-daily tacrolimus from the first dosing day to 1 year after renal transplantation. 2014 , 15, 1495-506	11
1282	Histomorphology of right versus left internal thoracic artery and risk factors for intimal hyperplasia. 2014 , 45, 726-31	17
1281	Renal function after radical nephrectomy: development and validation of predictive models in Japanese patients. 2014 , 21, 238-42	15
1280	Prediction of partial nephrectomy outcomes by using the diameter-axis-polar nephrometry score. 2014 , 21, 442-6	10
1279	Low serum potassium concentration is a predictor of chronic kidney disease. 2014 , 68, 700-4	8
1278	Chronic kidney disease score for predicting postoperative masked renal insufficiency in patients with primary aldosteronism. 2014 , 81, 665-70	13
1277	Obesity as an effect modifier of the association between leptin and diabetic kidney disease. 2014 , 5, 213-20	7
1276	Decreased mobility after starting dialysis is an independent risk factor for short-term mortality after initiation of dialysis. 2014 , 19, 227-33	17
1275	Effect of renal impairment on cognitive function during a 3-year follow up in elderly patients with type 2 diabetes: Association with microinflammation. 2014 , 5, 597-605	14
1274	Visit-to-visit variability of blood pressure and renal function decline in patients with diabetic chronic kidney disease. 2014 , 16, 362-6	15
1273	Rho-associated kinase activity is a predictor of cardiovascular outcomes. 2014 , 63, 856-64	39
1272	Correlations between serum bilirubin levels and diabetic nephropathy progression among Japanese type 2 diabetic patients: a prospective cohort study (Diabetes Distress and Care Registry at Tenri [DDCRT 5]). 2014 , 37, 252-8	43
1271	Plasma polyunsaturated fatty acid profile and delta-5 desaturase activity are altered in patients with type 2 diabetes. 2014 , 63, 1432-8	23
1270	Editorial comment to prediction of partial nephrectomy outcomes by using the diameter-axis-polar nephrometry score. 2014 , 21, 446-7	
1269	Comparative effects of telmisartan and valsartan as add-on agents for hypertensive patients with morning blood pressure insufficiently controlled by amlodipine monotherapy. 2014 , 37, 225-31	5
1268	Comparison of warm and cold ischemia on renal function after partial nephrectomy. 2014 , 84, 1408-12	38
1267	Effect of renal impairment on the pharmacokinetics, pharmacodynamics, and safety of empagliflozin, a sodium glucose cotransporter 2 inhibitor, in Japanese patients with type 2 diabetes mellitus. 2014 , 36, 1606-15	22

1266	Anti-albuminuric effect of the aldosterone blocker eplerenone in non-diabetic hypertensive patients with albuminuria: a double-blind, randomised, placebo-controlled trial. 2014 , 2, 944-53	72
1265	Better recovery of kidney function in patients with de novo chronic kidney disease after partial nephrectomy compared with those with pre-existing chronic kidney disease. 2014 , 21, 613-6	5
1264	Association between serum immunoglobulin G4 concentration and cardiac function among elderly cardiology inpatients. 2014 , 14, 582-8	1
1263	Chapter 2. Measurement and clinical evaluation of blood pressure. 2014 , 37, 266-278	12
1262	Kidney function crucially affects B-type natriuretic peptide (BNP), N-terminal proBNP and their relationship. 2014 , 44, 303-8	63
1261	Association of high-sensitivity cardiac troponin T and N-terminal pro-brain-type natriuretic peptide with left ventricular structure: J-HOP study. 2014 , 16, 354-61	5
1260	Two-hour post-load plasma glucose levels are associated with carotid intima-media thickness in subjects with normal glucose tolerance. 2014 , 31, 76-83	10
1259	Functional implications of regulatory B cells in human IgA nephropathy. 2014 , 79, 51-60	17
1258	Pathophysiological impact of serum fibroblast growth factor 23 in patients with nonischemic cardiac disease and early chronic kidney disease. 2014 , 307, H1504-11	35
1257	Clinical pharmacokinetics and pharmacodynamics of the novel SGLT2 inhibitor ipragliflozin. 2014 , 53, 975-88	23
1256	Association of serum bilirubin levels with development and progression of albuminuria, and decline in estimated glomerular filtration rate in patients with type 2 diabetes mellitus. 2014 , 5, 228-35	31
1255	WASP neuronet activated by bipolar-sigmoid functions and applied to glomerular-filtration-rate estimation. 2014 ,	2
1254	Practical approach to evaluate asymptomatic coronary artery disease in end-stage renal disease patients at the initiation of dialysis. 2014 , 18, 167-73	
1253	Hyperuricemia is a significant risk factor for the onset of chronic kidney disease. 2014 , 126, 33-8	25
1252	Predictive value of asymmetric dimethylarginine and C-reactive protein for the risk of developing metabolic syndrome in middle-aged men. 2014 , 5, 42-47	2
1251	A slight increase within the normal range of serum uric acid and the decline in renal function: associations in a community-based population. 2014 , 29, 2286-92	37
1250	Clinical value of procalcitonin for patients with suspected bloodstream infection. 2014 , 141, 43-51	36
1249	Dapagliflozin as monotherapy or combination therapy in Japanese patients with type 2 diabetes: an open-label study. 2014 , 5, 415-33	53

1248	Usefulness of presepsin in the diagnosis of sepsis in patients with or without acute kidney injury. 2014 , 14, 88	55
1247	Visit-to-visit blood pressure variability is a marker of cardiac diastolic function and carotid atherosclerosis. 2014 , 14, 188	18
1246	Chronic kidney disease is associated with dementia independent of cerebral small-vessel disease. 2014 , 82, 1051-7	71
1245	Clinical features and prognosis of patients with coronary spasm-induced non-ST-segment elevation acute coronary syndrome. 2014 , 3, e000795	36
1244	Efficacy of low-dose rosuvastatin in patients with type 2 diabetes and hypo high-density lipoprotein cholesterolaemia. 2014 , 42, 457-67	4
1243	Seasonal variations of severe hypoglycemia in patients with type 1 diabetes mellitus, type 2 diabetes mellitus, and non-diabetes mellitus: clinical analysis of 578 hypoglycemia cases. 2014 , 93, e148	20
1242	Worsening of renal function during 1 year after hospital discharge is a strong and independent predictor of all-cause mortality in acute decompensated heart failure. 2014 , 3, e001174	16
1241	A high normal ankle-brachial index is associated with proteinuria in a screened cohort of Japanese: the Okinawa Peripheral Arterial Disease Study. 2014 , 32, 1435-43; discussion 1443	10
1240	Age-adjusted level of circulating elastin as a cardiovascular risk factor in medical check-up individuals. 2014 , 15, 364-70	7
1239	Long-term exposure to tenofovir continuously decrease renal function in HIV-1-infected patients with low body weight: results from 10 years of follow-up. 2014 , 28, 1903-10	54
1238	Subacute kidney injury in hospitalized patients. 2014 , 9, 457-61	19
1237	Extensive late gadolinium enhancement on cardiovascular magnetic resonance predicts adverse outcomes and lack of improvement in LV function after steroid therapy in cardiac sarcoidosis. 2014 , 100, 1165-72	89
1236	Serum cystatin C, creatinine-based estimated glomerular filtration rate, and the risk of incident hypertension in middle-aged men. 2014 , 27, 596-602	6
1235	Impact of serum bilirubin levels on carotid atherosclerosis in patients with coronary artery disease. 2014 , 5, 24-27	6
1234	Retrospective study of salazosulfapyridine in eight patients with rheumatoid arthritis on hemodialysis. 2014 , 24, 285-90	5
1233	Morningness-eveningness questionnaire score and metabolic parameters in patients with type 2 diabetes mellitus. 2014 , 31, 1017-23	38
1232	A new modified CKD-EPI equation for Chinese patients with type 2 diabetes. 2014 , 9, e109743	17
1231	Age- and sex-related effects on ankle-brachial index in a screened cohort of Japanese: the Okinawa Peripheral Arterial Disease Study (OPADS). 2014 , 21, 712-8	20

1230	Sex differences in the prevalence, progression, and improvement of chronic kidney disease. 2014 , 39, 279-88	14
1229	Trend of office and home blood pressure control in treated hypertensive patients: changes in antihypertensive medication and salt intake. 2014 , 36, 103-7	3
1228	Incremental prognostic significance of the elevated levels of pentraxin 3 in patients with heart failure with normal left ventricular ejection fraction. 2014 , 3,	18
1227	Short sleep duration increases the risk of chronic kidney disease in shift workers. 2014 , 56, 1243-8	18
1226	Reduced-dose telaprevir-based triple antiviral therapy for recurrent hepatitis C after living donor liver transplantation. 2014 , 98, 994-9	7
1225	Impact of albuminuria on the incidence of periprocedural myocardial injury in patients undergoing elective coronary stent implantation. 2014 , 114, 42-6	7
1224	Mineral and bone disorder is temporary in patients treated with early rapid corticosteroid reduction after kidney transplantation: a single-center experience. 2014 , 46, 514-20	7
1223	Impact of atrial fibrillation on long-term clinical outcomes in outpatients with heart failure. 2014 , 30, 186-191	
1222	Poor glycemic control and decreased renal function are associated with increased intrarenal RAS activity in Type 2 diabetes mellitus. 2014 , 105, 40-6	13
1221	Characterization of in-stent neointimal tissue components following drug-eluting stent implantation according to the phase of restenosis using a 40-MHz intravascular ultrasound imaging system. 2014 , 64, 423-9	8
1220	Performance of the Japanese glomerular filtration rate equation based on standardized serum cystatin C in potential kidney donors. 2014 , 46, 314-7	6
1219	Presentations and outcomes of patients with acute decompensated heart failure admitted in the winter season. 2014 , 64, 470-5	8
1218	Efficacy and safety of febuxostat, a novel nonpurine selective inhibitor of xanthine oxidase for the treatment of hyperuricemia in kidney transplant recipients. 2014 , 46, 511-3	16
1217	Impact of continuous deterioration of kidney function 6 to 8 months after percutaneous coronary intervention for acute coronary syndrome. 2014 , 113, 1647-51	32
1216	Comparison of characteristics and healing course of diabetic foot ulcers by etiological classification: neuropathic, ischemic, and neuro-ischemic type. 2014 , 28, 528-35	36
1215	The utility of urinary myo-inositol as a marker of glucose intolerance. 2014 , 103, 88-96	13
1214	Impact of statin therapy on patients with coronary heart disease and aortic aneurysm or dissection. 2014 , 60, 604-12.e2	10
1213	Association of the triglycerides to high-density lipoprotein cholesterol ratio with the risk of chronic kidney disease: analysis in a large Japanese population. 2014 , 233, 260-7	50

1212	Association of serum IgG4 and soluble interleukin-2 receptor levels with epicardial adipose tissue and coronary artery calcification. 2014 , 428, 63-69	8
1211	Pregnancy in 4 women with childhood-onset steroid-sensitive nephrotic syndrome. 2014 , 3, 63-67	3
1210	Renal impairment and coronary collaterals in patients with acute coronary syndrome. 2014 , 39, 379-83	2
1209	Hypertriglyceridemia accompanied by increased serum complement component 3 and proteinuria in non-nephrotic chronic kidney disease. 2014 , 18, 453-60	3
1208	U-shaped association between body mass index and proteinuria in a large Japanese general population sample. 2014 , 18, 75-86	14
1207	Prevalence and risk factor analysis of nephrosclerosis and ischemic nephropathy in the Japanese general population. 2014 , 18, 461-8	7
1206	Kidney volume and function in autosomal dominant polycystic kidney disease. 2014 , 18, 157-65	29
1205	Lower serum albumin level is associated with higher fractional excretion of creatinine. 2014 , 18, 469-74	9
1204	Vital signs, QT prolongation, and newly diagnosed cardiovascular disease during severe hypoglycemia in type 1 and type 2 diabetic patients. 2014 , 37, 217-25	74
1203	Long-term incidence and prognostic factors of the progression of new coronary lesions in Japanese coronary artery disease patients after percutaneous coronary intervention. 2014 , 29, 437-42	19
1202	Relation of omega-3 fatty acid and C-reactive protein to peripheral artery disease in patients with coronary artery disease. 2014 , 29, 449-55	16
1201	Impact of aging on the clinical outcomes of Japanese patients with coronary artery disease after percutaneous coronary intervention. 2014 , 29, 156-64	31
1200	Serum tenascin-C level is associated with coronary plaque rupture in patients with acute coronary syndrome. 2014 , 29, 165-70	16
1199	Low high-density lipoprotein cholesterol is a residual risk factor associated with long-term clinical outcomes in diabetic patients with stable coronary artery disease who achieve optimal control of low-density lipoprotein cholesterol. 2014 , 29, 35-41	20
1198	Accelerated decline in renal function after acute myocardial infarction in patients with high low-density lipoprotein-cholesterol to high-density lipoprotein-cholesterol ratio. 2014 , 29, 7-14	4
1197	Long-term changes in renal function outcomes following radical cystectomy and urinary diversion. 2014 , 19, 1105-11	26
1196	GCK, GCKR polymorphisms and risk of chronic kidney disease in Japanese individuals: data from the J-MICC Study. 2014 , 27, 143-9	8
1195	Electrocardiographic left ventricular hypertrophy Cornell product is a feasible predictor of cardiac prognosis in patients with chronic heart failure. 2014 , 103, 275-84	8

1194	Health-related quality of life and prognosis in patients with chronic kidney disease: a 3-year follow-up study. 2014 , 18, 697-703	14
1193	Effects of atorvastatin and ezetimibe on endothelial function in dyslipidemic patients with chronic kidney disease. 2014 , 18, 704-10	8
1192	Intravenous salt supplementation with low-dose furosemide for treatment of acute decompensated heart failure. 2014 , 20, 295-301	19
1191	Telaprevir impairs renal function and increases blood ribavirin concentration during telaprevir/pegylated interferon/ribavirin therapy for chronic hepatitis C. 2014 , 21, 341-7	36
1190	Thresholds of various glycemic measures for diagnosing diabetes based on prevalence of retinopathy in community-dwelling Japanese subjects: the Hisayama Study. 2014 , 13, 45	27
1189	Predictive clinical parameters for the hemoglobin A1c-lowering effect of vildagliptin in Japanese patients with type 2 diabetes. 2014 , 5, 229-233	
1188	Dietary phosphorus restriction by a standard low-protein diet decreased serum fibroblast growth factor 23 levels in patients with early and advanced stage chronic kidney disease. 2014 , 18, 925-31	25
1187	Plasma homocysteine and folate levels and dietary folate intake in adolescents and young adults who underwent kidney transplantation during childhood. 2014 , 18, 151-6	4
1186	The role of a low glomerular density and being overweight in the etiology of proteinuria in CKD patients without known glomerular diseases. 2014 , 18, 911-7	1
1185	Clinical effectiveness of tolvaptan in patients with acute decompensated heart failure and renal failure: design and rationale of the AQUAMARINE study. 2014 , 28, 73-7	15
1184	An association between serum γ -glutamyltransferase and proteinuria in drinkers and non-drinkers: a Japanese nationwide cross-sectional survey. 2014 , 18, 899-910	4
1183	Preoperative estimated glomerular filtration rate as a significant predictor of long-term outcomes after coronary artery bypass grafting in Japanese patients. 2014 , 62, 95-102	11
1182	The eGFR-C study: accuracy of glomerular filtration rate (GFR) estimation using creatinine and cystatin C and albuminuria for monitoring disease progression in patients with stage 3 chronic kidney disease--prospective longitudinal study in a multiethnic population. 2014 , 15, 13	28
1181	Estimated glomerular filtration rate and daily amount of urinary protein predict the clinical remission rate of tonsillectomy plus steroid pulse therapy for IgA nephropathy. 2014 , 18, 606-12	6
1180	Predictors of septic shock in obstructive acute pyelonephritis. 2014 , 32, 803-11	40
1179	Long-term prognosis of 4 children with steroid-sensitive nephrotic syndrome and relapse after 30 years of age. 2014 , 3, 106-109	
1178	Safety and efficacy of skin patches containing loxoprofen sodium in diabetic patients with overt nephropathy. 2014 , 18, 487-91	5
1177	Overestimation of the risk of progression to end-stage renal disease in the poor prognosis' group according to the 2002 Japanese histological classification for immunoglobulin A nephropathy. 2014 , 18, 475-80	1

1176	Diuretics enhance effects of increased dose of candesartan on ambulatory blood pressure reduction in Japanese patients with uncontrolled hypertension treated with medium-dose angiotensin II receptor blockers. 2014 , 36, 40-5		2
1175	Low serum bilirubin concentration is a predictor of chronic kidney disease. 2014 , 234, 421-5		31
1174	Predictors of decreasing glomerular filtration rate and prevalence of chronic kidney disease after treatment of primary aldosteronism: renal outcome of 213 cases. 2014 , 99, 1593-8		65
1173	Impact of admission glycemia and glycosylated hemoglobin A1c on long-term clinical outcomes of non-diabetic patients with acute coronary syndrome. 2014 , 63, 106-11		17
1172	How to estimate kidney function in kidney transplant recipients with mild to moderate kidney impairment: comparison of estimated glomerular filtration (eGFR) values between creatinine-based GFR equations and cystatin C-based GFR equations for Japanese population. 2014 , 18, 130-4		9
1171	Estimation of GFR in South Asians: a study from the general population in Pakistan. <i>American Journal of Kidney Diseases</i> , 2014 , 63, 49-58	7.4	49
1170	A microarray analysis of urinary microRNAs in renal diseases. 2014 , 18, 711-7		14
1169	A higher frequency of CD4+CXCR5+ T follicular helper cells in patients with newly diagnosed IgA nephropathy. 2014 , 158, 101-8		23
1168	Lower serum endogenous secretory receptor for advanced glycation end product level as a risk factor of metabolic syndrome among Japanese adult men: a 2-year longitudinal study. 2014 , 99, 587-93		15
1167	Relationship between serum uric acid and serum oxidative stress markers in the Japanese general population. 2014 , 128, 49-56		22
1166	Urinary liver-type fatty acid-binding protein linked with increased risk of acute kidney injury after allogeneic stem cell transplantation. 2014 , 20, 2010-4		14
1165	Living donor liver transplantation for non-alcoholic steatohepatitis: A single center experience. 2014 , 44, E3-E10		15
1164	Urinary type IV collagen is related to left ventricular diastolic function and brain natriuretic peptide in hypertensive patients with prediabetes. 2014 , 28, 824-30		4
1163	Positive correlation between malondialdehyde-modified low-density lipoprotein cholesterol and vascular inflammation evaluated by 18F-FDG PET/CT. 2014 , 237, 404-9		8
1162	Assessing kidney function in transplant recipients: time to work together and address the most relevant questions. <i>American Journal of Kidney Diseases</i> , 2014 , 64, 818	7.4	1
1161	Creatinine-based GFR estimating equations in kidney transplant recipients. <i>American Journal of Kidney Diseases</i> , 2014 , 64, 818	7.4	2
1160	Mortality prediction by acute kidney injury biomarkers in comparison with serum creatinine. 2014 , 19, 646-51		5
1159	Safety, efficacy and renal effect of febuxostat in patients with moderate-to-severe kidney dysfunction. 2014 , 37, 919-25		51

1158	Aldosterone-to-renin ratio and nocturnal blood pressure decline assessed by self-measurement of blood pressure at home: the Ohasama Study. 2014 , 36, 108-14	14
1157	High plasma fetuin-A levels are associated with metabolic syndrome among males but not females in a Japanese general population. 2014 , 106, 128-35	17
1156	β-cell telomere attrition in diabetes: inverse correlation between HbA1c and telomere length. 2014 , 99, 2771-7	29
1155	Uric acid levels predict future blood pressure and new onset hypertension in the general Japanese population. 2014 , 28, 529-34	14
1154	Empagliflozin monotherapy in Japanese patients with type 2 diabetes mellitus: a randomized, 12-week, double-blind, placebo-controlled, phase II trial. 2014 , 31, 621-38	59
1153	V-shaped relationship between HbA1c and all-cause mortality in the elderly with type 2 diabetes. 2014 , 5, 319-322	
1152	Heart-type fatty acid-binding protein is a novel prognostic marker in patients with essential hypertension. 2014 , 176, 1323-5	8
1151	Clinical characteristics of resistant hypertension evaluated by ambulatory blood pressure monitoring. 2014 , 36, 454-8	4
1150	The right internal thoracic artery: how much do we know?. 2014 , 45, 731-2	3
1149	Ferric citrate hydrate for the treatment of hyperphosphatemia in nondialysis-dependent CKD. 2014 , 9, 543-52	86
1148	Serum under-O-glycosylated IgA1 level is not correlated with glomerular IgA deposition based upon heterogeneity in the composition of immune complexes in IgA nephropathy. 2014 , 15, 89	9
1147	DPP-4 inhibitor and alpha-glucosidase inhibitor equally improve endothelial function in patients with type 2 diabetes: EDGE study. 2014 , 13, 110	92
1146	Evaluation of split renal function before and after renal arterial embolization for angiomyolipoma using absolute ethanol. 2014 , 37, 1220-5	4
1145	Myocardial perfusion imaging for predicting cardiac events in Japanese patients with advanced chronic kidney disease: 1-year interim report of the J-ACCESS 3 investigation. 2014 , 41, 1701-9	23
1144	Short-term and long-term outcomes after gastrectomy for gastric cancer in patients with chronic kidney disease. 2014 , 38, 1453-60	23
1143	Development of a novel nomogram to predict hypertension cure after laparoscopic adrenalectomy in patients with primary aldosteronism. 2014 , 38, 2640-4	32
1142	Decreased glomerular filtration as the primary factor of elevated circulating suPAR levels in focal segmental glomerulosclerosis. 2014 , 29, 1553-60	28
1141	Knee extensor muscle strength and index of renal function associated with an exercise capacity of 5 metabolic equivalents in male chronic heart failure patients with chronic kidney disease. 2014 , 18, 313-9	1

1140	Classification of human immunodeficiency virus-infected patients with chronic kidney disease using a combination of proteinuria and estimated glomerular filtration rate. 2014 , 18, 600-5	10
1139	Clinical impact of albuminuria and glomerular filtration rate on renal and cardiovascular events, and all-cause mortality in Japanese patients with type 2 diabetes. 2014 , 18, 613-20	105
1138	Relation of epicardial fat to central aortic pressure and left ventricular diastolic function in patients with known or suspected coronary artery disease. 2014 , 30, 1393-8	10
1137	Higher serum soluble receptor for advanced glycation end product levels and lower prevalence of metabolic syndrome among Japanese adult men: a cross-sectional study. 2014 , 6, 33	13
1136	Effects of three kinds of erythropoiesis-stimulating agents on renal anemia in Japanese non-dialysis chronic kidney disease patients. 2014 , 18, 755-62	11
1135	Validation of the Kidney Disease Improving Global Outcomes criteria for AKI and comparison of three criteria in hospitalized patients. 2014 , 9, 848-54	95
1134	Impact of combined losartan/hydrochlorothiazide on proteinuria in patients with chronic kidney disease and hypertension. 2014 , 37, 993-8	10
1133	Novel urinary biomarkers in early diabetic kidney disease. 2014 , 14, 513	19
1132	Safety margin of minimized contrast volume during percutaneous coronary intervention in patients with chronic kidney disease. 2014 , 29, 209-15	19
1131	Follow-up study of unilateral renal function after nephrectomy assessed by glomerular filtration rate per functional renal volume. 2014 , 12, 59	15
1130	The impact of elevation of serum uric acid level on the natural history of glomerular filtration rate (GFR) and its sex difference. 2014 , 29, 1932-9	22
1129	Poor glycemic control is a major factor in the overestimation of glomerular filtration rate in diabetic patients. 2014 , 37, 596-603	25
1128	The revised Lund-Malm GFR estimating equation outperforms MDRD and CKD-EPI across GFR, age and BMI intervals in a large Swedish population. 2014 , 52, 815-24	94
1127	Ankle-brachial blood pressure index predicts cardiovascular events and mortality in Japanese patients with chronic kidney disease not on dialysis. 2014 , 37, 1050-5	7
1126	Generation of a new cystatin C-based estimating equation for glomerular filtration rate by use of 7 assays standardized to the international calibrator. 2014 , 60, 974-86	182
1125	Statins combined with exercise are associated with the increased renal function mediated by high-molecular-weight adiponectin in coronary artery disease patients. 2014 , 64, 91-7	4
1124	Outcome of elective total aortic arch replacement in patients with non-dialysis-dependent renal insufficiency stratified by estimated glomerular filtration rate. 2014 , 147, 966-972.e2	15
1123	Associations of proteinuria and the estimated glomerular filtration rate with incident hypertension in young to middle-aged Japanese males. 2014 , 60, 48-54	10

1122	Correlation between serum lipids and 1-hour postload plasma glucose levels in normoglycemic individuals. 2014 , 8, 217-22	6
1121	Significance of imbalance in the ratio of serum n-3 to n-6 polyunsaturated fatty acids in patients with acute coronary syndrome. 2014 , 113, 441-5	48
1120	Association of plasma thioredoxin-1 with renal tubular damage and cardiac prognosis in patients with chronic heart failure. 2014 , 64, 353-9	6
1119	Association between HbA1c variability and mortality in patients with type 2 diabetes. 2014 , 28, 494-9	44
1118	Visit-to-visit variability in systolic blood pressure predicts development and progression of diabetic nephropathy, but not retinopathy, in patients with type 2 diabetes. 2014 , 28, 185-90	20
1117	Improvement of HAS-BLED bleeding score predictive capability by changing the definition of renal dysfunction in Japanese atrial fibrillation patients on anticoagulation therapy. 2014 , 64, 482-7	12
1116	Serum vaspin levels are positively associated with carotid atherosclerosis in a general population. 2014 , 233, 248-52	23
1115	Endothelial function and cardiovascular events in chronic kidney disease. 2014 , 173, 481-6	38
1114	Conversion to mycophenolate mofetil from azathioprine shows significant positive effect on graft function in long-term post-kidney transplantation stable-state patients. 2014 , 46, 411-4	3
1113	Impact of diabetic retinopathy on late cardiac events after percutaneous coronary intervention. 2014 , 64, 175-8	3
1112	Effects of conversion from a twice-daily tacrolimus to a once-daily tacrolimus on glucose metabolism in stable kidney transplant recipients. 2014 , 46, 532-6	10
1111	Investigation of urinary angiotensinogen in renal transplant recipients. 2014 , 46, 489-91	5
1110	Reduction of dental metallic artefacts in CT: value of a newly developed algorithm for metal artefact reduction (O-MAR). 2014 , 69, e11-6	76
1109	Stroke and renal dysfunction. 2014 , 25, 18-24	18
1108	Asymmetric dimethylarginine is related to the predicted stroke risk in middle-aged Japanese men. 2014 , 338, 87-91	9
1107	Association of renal tubular damage with cardio-renal anemia syndrome in patients with heart failure. 2014 , 173, 222-8	10
1106	Performance of creatinine-based GFR estimating equations in solid-organ transplant recipients. <i>American Journal of Kidney Diseases</i> , 2014 , 63, 1007-18	7-4 85
1105	Novel preprocedural and acute-phase postprocedural predictive factors for contrast-induced kidney injury in CKD patients. 2014 , 172, e293-6	9

1104	Change in urinary N-acetyl-β-d-glucosaminidase levels relevant to postprandial glycemic control conditions in subjects without diabetes mellitus. 2014 , 433, 88-92	5
1103	Noncardiovascular death, especially infection, is a significant cause of death in elderly patients with acutely decompensated heart failure. 2014 , 20, 174-80	12
1102	Intravascular embolization therapy in patients with enlarged polycystic liver. <i>American Journal of Kidney Diseases</i> , 2014 , 63, 937-44	7.4 27
1101	Association of contrast-induced acute kidney injury with long-term cardiovascular events in acute coronary syndrome patients with chronic kidney disease undergoing emergent percutaneous coronary intervention. 2014 , 174, 57-63	65
1100	Impact of pre-discharge nocturnal pulse oximetry (sleep-disordered breathing) on postdischarge clinical outcomes in hospitalized patients with left ventricular systolic dysfunction after acute decompensated heart failure. 2014 , 113, 697-700	22
1099	Impact of recipient aging on kidney allograft in living donor transplantation. 2014 , 46, 454-6	3
1098	Angiotensin receptor blocker (ARB)-diuretic versus ARB-calcium channel blocker combination therapy for hypertension uncontrolled by ARB monotherapy. 2014 , 36, 244-50	10
1097	Long-term use of protease inhibitors is associated with bone mineral density loss. 2014 , 30, 553-9	21
1096	Conformational change in transfer RNA is an early indicator of acute cellular damage. 2014 , 25, 2316-26	70
1095	Call for the use of a common equation for glomerular filtration rate estimation in East and South-East Asia. 2014 , 47, 1214-9	
1094	Mannitol has no impact on renal function after open partial nephrectomy in solitary kidneys. 2014 , 21, 200-3	27
1093	Muscle mass, visceral fat, and plasma levels of B-type natriuretic peptide in healthy individuals (from the J-SHIP Study). 2014 , 114, 635-40	12
1092	Time-to-effect relationships between systolic blood pressure and the risks of nephropathy and retinopathy in patients with type 2 diabetes. 2014 , 28, 674-8	5
1091	Relation of contrast-induced nephropathy to long-term mortality after percutaneous coronary intervention. 2014 , 114, 362-8	69
1090	Assessment of postoperative changes in antihypertensive drug consumption in patients with primary aldosteronism using the defined daily dose. 2014 , 37, 190-4	7
1089	The number of metabolic syndrome components is a good risk indicator for both early- and late-stage kidney damage. 2014 , 24, 277-85	23
1088	Effects of statin treatment in patients with coronary artery disease and chronic kidney disease. 2014 , 29, 21-8	27
1087	Elevation of urinary liver-type fatty acid-binding protein as predicting factor for occurrence of contrast-induced acute kidney injury and its reduction by hemodiafiltration with blood suction from right atrium. 2014 , 29, 191-7	12

1086	[111th Scientific Meeting of the Japanese Society of Internal Medicine: Invited Lecture: 1. Diabetes and its complication, focusing on diabetic nephropathy]. 2014 , 103, 2040-50	
1085	[111th Scientific Meeting of the Japanese Society of Internal Medicine: Educational Lecture: 3. Clinical epidemiology of CKD]. 2014 , 103, 2242-6	
1084	Renal and vascular protective effects of ezetimibe in chronic kidney disease. 2014 , 53, 307-14	10
1083	Response to clopidogrel and its association with chronic kidney disease in noncardiogenic ischemic stroke patients. 2014 , 53, 215-9	9
1082	The dose-response relationship between body mass index and the risk of incident stage B chronic kidney disease in a general Japanese population: the Ibaraki prefectural health study (IPHS). 2014 , 24, 444-51	32
1081	Relative risk values of age, acrolein, IL-6 and CRP as markers of periventricular hyperintensities: a cross-sectional study. 2014 , 4, e005598	3
1080	Relationship between the accuracy of glycemic markers and the chronic kidney disease stage in patients with type 2 diabetes mellitus. 2014 , 82, 107-14	12
1079	Alogliptin improves steroid-induced hyperglycemia in treatment-naïve Japanese patients with chronic kidney disease by decrease of plasma glucagon levels. 2014 , 20, 587-93	13
1078	Impact of chronic kidney disease and anemia on physical function in patients with chronic heart failure. 2014 , 4, 73-81	3
1077	A cross-sectional study to find out the relationship of methylenetetrahydrofolate reductase (MTHFR) C677T genotype with plasma levels of folate and total homocysteine by daily folate intake in Japanese. 2014 , 60, 231-8	6
1076	Long-term outcome of the carpentier-edwards pericardial valve in the aortic position in Japanese patients. 2014 , 78, 882-9	34
1075	Serum 1,25-dihydroxyvitamin d and the development of kidney dysfunction in a Japanese community. 2014 , 78, 732-7	7
1074	Clinical predictors of contrast-induced acute kidney injury in patients undergoing emergency versus elective percutaneous coronary intervention. 2014 , 78, 85-91	40
1073	Impact of cardiac rehabilitation on renal function in patients with and without chronic kidney disease after acute myocardial infarction. 2014 , 78, 377-84	34
1072	Prognostic value of pulmonary hypertension in ambulatory patients with non-ischemic dilated cardiomyopathy. 2014 , 78, 1245-53	15
1071	Preoperative chronic kidney disease as a strong predictor of postoperative infection and mortality after coronary artery bypass grafting. 2014 , 78, 2225-31	26
1070	Plasma indoxyl sulfate and estimated glomerular filtration rate. 2014 , 78, 2477-82	15
1069	Use of proton pump inhibitors is associated with anemia in cardiovascular outpatients. 2015 , 79, 193-200	10

1068	Regression of Electrocardiographic Signs of Left Ventricular Hypertrophy by Combined Treatment With Thiazide Diuretic and Angiotensin-II Receptor Blocker. 2014 , 78, 2719-26	3
1067	Diagnostic accuracy of Global Registry of Acute Coronary Events (GRACE) risk score in ST-elevation myocardial infarction for in-hospital and 360-day mortality in Japanese patients. 2014 , 78, 2950-4	29
1066	Lower beta cell function relates to sustained higher glycated albumin to glycated hemoglobin ratio in Japanese patients with type 2 diabetes. 2014 , 61, 149-57	17
1065	Effects of alogliptin in chronic kidney disease patients with type 2 diabetes. 2014 , 53, 195-203	14
1064	Type 2 diabetes mellitus complicated by hypertension in Japanese patients: switching treatment from high-dose angiotensin II receptor blockers to losartan plus hydrochlorothiazide. 2014 , 53, 1283-9	2
1063	Aortic arch atherosclerosis in ischaemic stroke of unknown origin affects prognosis. 2014 , 4, 92-101	4
1062	An integrative study of the genetic, social and environmental determinants of chronic kidney disease characterized by tubulointerstitial damages in the North Central Region of Sri Lanka. 2014 , 56, 28-38	73
1061	Determinants of plasma renin activity: role of a human renin gene variant as a genetic factor. 2014 , 93, e354	6
1060	Hypertension increases urinary excretion of immunoglobulin G, ceruloplasmin and transferrin in normoalbuminuric patients with type 2 diabetes mellitus. 2014 , 32, 432-8	15
1059	Short-term prognosis of living-donor kidney transplantation from hypertensive donors with high-normal albuminuria. 2014 , 97, 104-10	15
1058	Association of cognitive dysfunction with cardiovascular disease events in elderly hypertensive patients. 2014 , 32, 423-31	13
1057	Hemoglobin a1c levels and the risk of cardiovascular disease in people without known diabetes: a population-based cohort study in Japan. 2015 , 94, e785	20
1056	Comparison of Renal Scintigraphy and Computed Tomographic Renal Volumetry for Determining Split Renal Function and Estimating Post-Transplant Renal Function. 2015 , 47, 2700-2	13
1055	Estimated glomerular filtration rate by serum creatinine or standardized cystatin C in Japanese patients with Graves' disease. 2015 , 5, 1092-5	
1054	Prediction of renal function after nephroureterectomy in patients with upper tract urothelial carcinoma. 2015 , 45, 1064-8	8
1053	Choice of Antihypertensive Combination Therapy Based on Daily Salt Intake. 2015 , 350, 160-6	3
1052	The relationship of fibroblast growth factors 21 and 23 and Klotho with platelet activity measured by platelet volume indices. 2015 , 53, 1569-74	2
1051	Plasma levels of natriuretic peptides and development of chronic kidney disease. 2015 , 16, 171	11

1050	Effects of a high-pitch protocol and a hybrid iterative reconstruction algorithm on image quality of cerebral subtracted 3D CT angiography. 2015 , 33, 687-93	2
1049	Orally active prostacyclin analogue beraprost sodium in patients with chronic kidney disease: a randomized, double-blind, placebo-controlled, phase II dose finding trial. 2015 , 16, 165	16
1048	Impact of preoperative asymptomatic renal dysfunction on clinical course after pancreatoduodenectomy. 2015 , 22, 810-8	10
1047	FABP4 is secreted from adipocytes by adenylyl cyclase-PKA- and guanylyl cyclase-PKG-dependent lipolytic mechanisms. 2015 , 23, 359-67	59
1046	Differences in risk factors for the onset of albuminuria and decrease in glomerular filtration rate in people with Type 2 diabetes mellitus: implications for the pathogenesis of diabetic kidney disease. 2015 , 32, 1354-60	38
1045	Feasibility and accuracy of computational robot-assisted partial nephrectomy planning by virtual partial nephrectomy analysis. 2015 , 22, 439-46	24
1044	Japanese society for dialysis therapy clinical guideline for "hemodialysis initiation for maintenance hemodialysis". 2015 , 19 Suppl 1, 93-107	38
1043	Early unclamping might reduce the risk of renal artery pseudoaneurysm after robot-assisted laparoscopic partial nephrectomy. 2015 , 22, 1096-102	37
1042	Efficacy and safety of canagliflozin alone or as add-on to other oral antihyperglycemic drugs in Japanese patients with type 2 diabetes: A 52-week open-label study. 2015 , 6, 210-8	55
1041	Circulating Fibroblast Growth Factor 23 Has a U-Shaped Association With Atrial Fibrillation Prevalence. 2015 , 79, 1742-8	11
1040	Clinical outcome of renal artery stenting for hypertension and chronic kidney disease up to 12 months in the J-RAS Study [prospective, single-arm, multicenter clinical study]. 2015 , 79, 351-9	20
1039	Proteinuria and Reduced Estimated Glomerular Filtration Rate Are Independent Risk Factors for Contrast-Induced Nephropathy After Cardiac Catheterization. 2015 , 79, 1624-30	17
1038	SYMPPLICITY HTN-Japan - First Randomized Controlled Trial of Catheter-Based Renal Denervation in Asian Patients -. 2015 , 79, 1222-9	76
1037	Association of Estimated Glomerular Filtration Rate and Proteinuria With Lipid-Rich Plaque in Coronary Artery Disease. 2015 , 79, 2263-70	6
1036	Risk Factors for Progression of Degenerative Aortic Valve Disease in the Japanese- The Japanese Aortic Stenosis Study (JASS) Prospective Analysis. 2015 , 79, 2050-7	12
1035	Twenty-year outcome of aortic valve replacement with St. Jude Medical mechanical valves in Japanese patients. 2015 , 79, 2380-8	10
1034	Ameliorated pancreatic β cell dysfunction in type 2 diabetic patients treated with a sodium-glucose cotransporter 2 inhibitor ipragliflozin. 2015 , 62, 77-86	38
1033	Circulating betatrophin is elevated in patients with type 1 and type 2 diabetes. 2015 , 62, 417-21	71

1032	Early liraglutide treatment improves β -cell function in patients with type 2 diabetes: a retrospective cohort study. 2015 , 62, 971-80	11
1031	HOMA-IR Values are Associated With Glycemic Control in Japanese Subjects Without Diabetes or Obesity: The KOBE Study. 2015 , 25, 407-14	14
1030	Prognostic Significance of Neuroadrenergic Dysfunction for Cardiovascular Events in Patients With Acute Myocardial Infarction. 2015 , 79, 2238-45	9
1029	Nocturia in Patients With Sleep-Disordered Breathing and Cardiovascular Disease. 2015 , 79, 2632-40	19
1028	Microvascular resistance in response to iodinated contrast media in normal and functionally impaired kidneys. 2015 , 42, 1245-50	6
1027	Association of genetic variants with dyslipidemia and chronic kidney disease in a longitudinal population-based genetic epidemiological study. 2015 , 35, 1290-300	12
1026	Age-related changes in clinical parameters and their associations with common complex diseases. 2015 , 3, 767-777	11
1025	Increased Total Homocysteine Levels Predict the Risk of Incident Dementia Independent of Cerebral Small-Vessel Diseases and Vascular Risk Factors. 2016 , 49, 503-13	34
1024	Development and Assessment of a Nomogram to Propose the Initial Dosage Regimen of a Meropenem Infusion Based on Serum Creatinine and Age Using a Monte Carlo Simulation. 2015 , 63, 986-91	4
1023	Cystatin C-based eGFR is a superior prognostic parameter to creatinine-based eGFR in post-endovascular therapy peripheral artery disease patients. 2015 , 79, 2480-6	11
1022	High salt intake is associated with renal involvement in Japanese patients with type 2 diabetes mellitus. 2015 , 54, 311-7	10
1021	A Reduction of HbA1c after 3 Months Predicts 2-year Responsiveness to Sitagliptin Treatment. 2015 , 54, 2981-9	4
1020	Saturated Metabolism of Voriconazole N-Oxidation Resulting in Nonlinearity of Pharmacokinetics of Voriconazole at Clinical Doses. 2015 , 38, 1496-503	16
1019	Risk of chronic kidney disease in non-obese individuals with clustering of metabolic factors: a longitudinal study. 2015 , 54, 375-82	20
1018	Serum Cholinesterase Is Inversely Associated with Body Weight Change in Men Undergoing Routine Health Screening. 2015 , 54, 2427-32	5
1017	Associations between the intake of miso soup and Japanese pickles and the estimated 24-hour urinary sodium excretion: a population-based cross-sectional study. 2015 , 54, 903-10	18
1016	The Optimal Cut-off Value of Plasma BNP to Differentiate Heart Failure in the Emergency Department in Japanese Patients with Dyspnea. 2015 , 54, 2975-80	8
1015	Efficacy and safety of telaprevir, pegylated interferon β and ribavirin triple therapy in Japanese patients infected with hepatitis C virus genotype 1b. 2015 , 54, 2551-60	7

1014	[Kidney diseases and metabolic disorders--Basics and applications required for general physicians. Topics: IX. Impaired drug metabolism]. 2015 , 104, 967-74	
1013	Elevated plasma B-type natriuretic peptide concentration and resistive index, but not decreased aortic distensibility, associate with impaired blood flow at popliteal artery in type 2 diabetic patients. 2015 , 62, 503-11	2
1012	Within-visit blood pressure variability is associated with prediabetes and diabetes. 2015 , 5, 7964	17
1011	The combined effect of visit-to-visit variability in HbA1c and systolic blood pressure on the incidence of cardiovascular events in patients with type 2 diabetes. 2015 , 3, e000129	27
1010	Role of Pulse Wave Velocity in Patients with Chronic Kidney Disease Stages 3-5 on Long-Term Follow-Up. 2014 , 2, 1-10	4
1009	Decline of Renal Function and Progression of Left Ventricular Hypertrophy Are Independently Determined in Chronic Kidney Disease Stages 3-5. 2014 , 2, 29-37	
1008	Approximation of Corrected Calcium Concentrations in Advanced Chronic Kidney Disease Patients with or without Dialysis Therapy. 2015 , 5, 39-49	9
1007	Comparison between Cystatin C- and Creatinine-Estimated Glomerular Filtration Rate in Cardiology Patients. 2015 , 5, 289-96	3
1006	The prevalence, characteristics, and clinical significance of abnormal albuminuria in patients with hypertension. 2014 , 4, 3884	11
1005	Quantification of excretory renal function and urinary protein excretion by determination of body cell mass using bioimpedance analysis. 2015 , 16, 174	7
1004	Echocardiographic characteristics of patients with acute heart failure requiring tolvaptan: a retrospective study. 2015 , 13, 27	5
1003	Development and validation of new glomerular filtration rate predicting models for Chinese patients with type 2 diabetes. 2015 , 13, 317	7
1002	Association of hemoglobin A1c and glycated albumin with carotid atherosclerosis in community-dwelling Japanese subjects: the Hisayama Study. 2015 , 14, 84	21
1001	The Associations of Malnutrition and Aging with Fluid Volume Imbalance between Intra- and Extracellular Water in Patients with Chronic Kidney Disease. 2015 , 19, 986-93	17
1000	Membranoproliferative glomerulonephritis and C3 glomerulonephritis: frequency, clinical features, and outcome in children. 2015 , 20, 286-92	16
999	Coronary microvascular function is independently associated with left ventricular filling pressure in patients with type 2 diabetes mellitus. 2015 , 14, 98	38
998	Plasma leptin level is associated with cardiac autonomic dysfunction in patients with type 2 diabetes: HSCAA study. 2015 , 14, 117	27
997	Effects of pitavastatin add-on therapy on chronic kidney disease with albuminuria and dyslipidemia. 2015 , 14, 161	5

996	Serum uric acid is associated with cardiac diastolic dysfunction among women with preserved ejection fraction. 2015 , 309, H986-94	18
995	Comparing the efficacy of continuous erythropoietin receptor activator and darbepoetin Alfa treatments in Japanese patients with chronic kidney disease during the predialysis period: A propensity-matched analysis. 2015 , 20 Suppl 4, 22-8	7
994	Reduced Glutathione for Prevention of Renal Outcomes in Patients Undergoing Selective Coronary Angiography or Intervention. 2015 , 28, 249-56	5
993	Anaemia management and mortality risk in newly visiting patients with chronic kidney disease in Japan: The CKD-ROUTE study. 2015 , 20, 601-8	32
992	The Pivotal Role of a Novel Biomarker of Reactive Oxygen Species in Chronic Kidney Disease. 2015 , 94, e1040	14
991	Relationship between vascular function indexes, renal arteriosclerosis, and renal clinical outcomes in chronic kidney disease. 2015 , 20, 585-90	11
990	Change of carotid intima-media thickness is associated with age in elderly Japanese patients without a history of cardiovascular disease. 2015 , 15, 1023-30	5
989	Relationship between inter-arm blood pressure differences and future cardiovascular events in coronary artery disease. 2015 , 33, 1780-9; discussion 1790	14
988	Effects of single-session dietary counseling by dieticians on salt reduction in cardiology outpatients who consumed large amounts of salt. 2015 , 10, 113-116	1
987	Incidence and predictors of rehospitalization of acute heart failure patients. 2015 , 56, 219-25	25
986	Relationship Between Retinal Blood Flow and Serum Adiponectin Concentrations in Patients With Type 2 Diabetes Mellitus. 2015 , 56, 4143-9	17
985	High-sensitivity C-reactive Protein, Lipoprotein(a) and Homocysteine are Risk Factors for Coronary Artery Disease in Japanese Patients with Peripheral Arterial Disease. 2015 , 22, 344-54	23
984	Efficacy and safety of sitagliptin in Japanese patients with type 2 diabetes. 2015 , 7, 211-9	14
983	Elevation of urinary liver-type fatty acid binding protein after cardiac catheterization related to cardiovascular events. 2015 , 8, 91-9	3
982	Efficacy And Safety of Coadministration of Tolvaptan And Carperitide for Acute Decompensated Heart Failure Patients. 2015 , 06,	1
981	A Higher Cardiothoracic Ratio Is Associated with 2-Year Mortality after Hemodialysis Initiation. 2015 , 5, 100-10	6
980	A Trial of Pitavastatin Versus Rosuvastatin for Dyslipidemia in Chronic Kidney Disease. 2015 , 22, 1235-47	3
979	Imaging of the Corneal Subbasal Whorl-like Nerve Plexus: More Accurate Depiction of the Extent of Corneal Nerve Damage in Patients With Diabetes. 2015 , 56, 5417-23	31

978	Efficacy and safety of sitagliptin added to insulin in Japanese patients with type 2 diabetes: the EDIT randomized trial. 2015 , 10, e0121988	23
977	Novel Autoantigens Associated with Lupus Nephritis. 2015 , 10, e0126564	9
976	Distinct roles of urinary liver-type fatty acid-binding protein in non-diabetic patients with anemia. 2015 , 10, e0126990	12
975	Individualizing pharmacotherapy in patients with renal impairment: the validity of the Modification of Diet in Renal Disease formula in specific patient populations with a glomerular filtration rate below 60 ml/min. A systematic review. 2015 , 10, e0116403	9
974	Impact of Admission Anemia on Coronary Microcirculation and Clinical Outcomes in Patients With ST-Segment Elevation Myocardial Infarction Undergoing Primary Percutaneous Coronary Intervention. 2015 , 56, 381-8	13
973	Association between Hemoglobin Concentration and the Progression or Development of Albuminuria in Diabetic Kidney Disease. 2015 , 10, e0129192	3
972	Influence of the PNPLA3 rs738409 Polymorphism on Non-Alcoholic Fatty Liver Disease and Renal Function among Normal Weight Subjects. 2015 , 10, e0132640	40
971	Alcohol and Exercise Affect Declining Kidney Function in Healthy Males Regardless of Obesity: A Prospective Cohort Study. 2015 , 10, e0134937	9
970	Relationship between Serum Uric Acid Levels and Chronic Kidney Disease in a Japanese Cohort with Normal or Mildly Reduced Kidney Function. 2015 , 10, e0137449	17
969	Pre- and/or Intra-Operative Prescription of Diuretics, but Not Renin-Angiotensin-System Inhibitors, Is Significantly Associated with Acute Kidney Injury after Non-Cardiac Surgery: A Retrospective Cohort Study. 2015 , 10, e0132507	24
968	Comparative Effects of Statins on the Kidney Function in Patients with Type 2 Diabetes. 2015 , 22, 618-27	3
967	Prevalence of chronic kidney diseases in patients with chronic obstructive pulmonary disease: assessment based on glomerular filtration rate estimated from creatinine and cystatin C levels. 2015 , 10, 1283-9	32
966	The relation between CA 19-9 level and early-phase insulin secretion in normoglycemic and prediabetic subjects. 2015 , 30, e169-73	3
965	Gender Specific Association between Serum Fibroblast Growth Factor 23/ β Klotho and Coronary Artery and Aortic Valve Calcification. 2015 , 22, 1338-46	8
964	Urinary Angiotensinogen Could Be a Prognostic Marker of Renoprotective Effects of Alogliptin in Patients with Type 2 Diabetes. 2015 , 2015, 517472	5
963	Improvement of Renal Function by Long-Term Sustained Eculizumab Treatment in a Patient with Paroxysmal Nocturnal Hemoglobinuria. 2015 , 2015, 673195	1
962	Association of Fish Consumption-Derived Ratio of Serum n-3 to n-6 Polyunsaturated Fatty Acids and Cardiovascular Risk With the Prevalence of Coronary Artery Disease. 2015 , 56, 260-8	20
961	Flow Mediated Dilatation Is Reduced with the Progressive Stages of Glomerular Filtration Rate and Albuminuria in Type 2 Diabetic Patients without Coronary Heart Disease. 2015 , 2015, 728127	16

960	Optimal Hemoglobin A1c Levels for Screening of Diabetes and Prediabetes in the Japanese Population. 2015 , 2015, 932057	17
959	Hyperkalemia develops in some thyroidectomized patients undergoing thyroid hormone withdrawal in preparation for radioactive iodine ablation for thyroid carcinoma. 2015 , 21, 488-94	2
958	Echolucent Carotid Plaque is Associated with Future Renal Dysfunction in Patients with Stable Coronary Artery Disease. 2015 , 22, 685-96	3
957	Urinary albumin as a marker of future blood pressure and hypertension in the general population. 2015 , 94, e511	23
956	Adjusting the starting dose of telaprevir according to renal function decreases adverse effects and affects the sustained virological response rate. 2015 , 27, 55-64	2
955	Sexual networks, HIV, race and bacterial vaginosis. 2015 , 29, 641-2	8
954	High-normal blood pressure is associated with new-onset electrocardiographic left ventricular hypertrophy. 2015 , 29, 9-13	6
953	Glomerular Density in Biopsy-Proven Hypertensive Nephrosclerosis. 2015 , 28, 1164-71	15
952	Gender- and disease-specific urinary thioredoxin in chronic kidney disease patients with or without type 2 diabetic nephropathy. 2015 , 20, 368-74	4
951	Effects of Celiprolol and Bisoprolol on Blood Pressure, Vascular Stiffness, and Baroreflex Sensitivity. 2015 , 28, 858-67	20
950	Uric acid level has a U-shaped association with loss of kidney function in healthy people: a prospective cohort study. 2015 , 10, e0118031	52
949	Serum uric acid and the incidence of CKD and hypertension. 2015 , 19, 1127-34	30
948	Impact of renal function of patients with advanced urothelial cancer on eligibility for first-line chemotherapy and treatment outcomes. 2015 , 45, 867-73	10
947	. 2015 , 185-231	
946	Factors affecting the progression of renal dysfunction and the importance of salt restriction in patients with type 2 diabetic kidney disease. 2015 , 19, 1120-6	9
945	Comparison of predictive value for first cardiovascular event between Japanese GFR equation and coefficient-modified CKD-EPI equation. 2015 , 19, 387-94	7
944	Is the eGFR formula adequate for evaluating renal function before chemotherapy in patients with urogenital cancer? A suggestion for clinical application of eGFR formula. 2015 , 19, 738-45	1
943	Nephrotic range proteinuria as a strong risk factor for rapid renal function decline during pre-dialysis phase in type 2 diabetic patients with severely impaired renal function. 2015 , 19, 1037-43	8

942	High risk of abnormal QT prolongation in the early morning in diabetic and non-diabetic patients with severe hypoglycemia. 2015 , 47, 238-44	12
941	Pretreatment neutrophil-to-lymphocyte ratio as an independent predictor of survival in patients with metastatic urothelial carcinoma: A multi-institutional study. 2015 , 22, 638-43	34
940	Patients with biopsy-proven nephrosclerosis and moderately impaired renal function have a higher risk for cardiovascular disease: 15 years' experience in a single, kidney disease center. 2015 , 9, 77-86	4
939	Effectiveness of Prior Use of Beta-Blockers for Preventing Adverse Influences of Severe Hypoglycemia in Patients With Diabetes: An Observational Study. 2015 , 94, e1629	10
938	Association between pulse wave velocity and a marker of renal tubular damage (N-acetyl-β-D-glucosaminidase) in patients without diabetes. 2015 , 17, 290-7	9
937	Arteriosclerosis in zero-time biopsy is a risk factor for tacrolimus-induced chronic nephrotoxicity. 2015 , 20 Suppl 2, 51-7	9
936	Urinary phosphorus excretion per creatinine clearance as a prognostic marker for progression of chronic kidney disease: a retrospective cohort study. 2015 , 16, 116	11
935	Gender-related alterations in plasma adrenomedullin level and its correlation with body weight gain. 2015 , 4, 43-9	15
934	Preoperative renal function and surgical outcomes in patients with acute type A aortic dissection□	7
933	Ambulatory blood pressure and tubulointerstitial injury in patients with IgA nephropathy. 2015 , 8, 716-21	2
932	Impact of increased visceral adiposity with normal weight on the progression of arterial stiffness in Japanese patients with type 2 diabetes. 2015 , 3, e000081	25
931	ACE insertion/deletion polymorphism (rs1799752) modifies the renoprotective effect of renin-angiotensin system blockade in patients with IgA nephropathy. 2015 , 16, 633-41	10
930	Dietary Sodium Consumption Predicts Future Blood Pressure and Incident Hypertension in the Japanese Normotensive General Population. 2015 , 4, e001959	36
929	Impact of Renal Transplantation and Nephrectomy on Urinary Soluble Klotho Protein. 2015 , 47, 1697-9	9
928	The prognostic impact of worsening renal function in Japanese patients undergoing percutaneous coronary intervention with acute coronary syndrome. 2015 , 66, 326-32	12
927	Fibroblast Growth Factor-23 and Vitamin D Metabolism in Subjects with eGFR ≤ 30 ml/min/1.73 m ² . 2015 , 130, 119-26	6
926	Acute Kidney Injury Facilitates Hypocalcemia by Exacerbating the Hyperphosphatemic Effect of Muscle Damage in Rhabdomyolysis. 2015 , 131, 11-6	7
925	Acute ischemic stroke associated with nephrotic syndrome: Incidence and significance - Retrospective cohort study. 2015 , 1, 47-50	5

924	Do metastatic upper tract urothelial carcinoma and bladder carcinoma have similar clinical responses to systemic chemotherapy? A Japanese multi-institutional experience. 2016 , 46, 163-9	11
923	Bucillamine-induced membranous nephropathy with crescent formation in a patient with rheumatoid arthritis: case report and literature review. 2015 , 5, 30-8	7
922	Cross-sectional and longitudinal associations between serum bilirubin and dyslipidemia in a health screening population. 2015 , 239, 31-7	7
921	Reduced estimated glomerular filtration rate affects outcomes 3 months after intracerebral hemorrhage: the stroke acute management with urgent risk-factor assessment and improvement-intracerebral hemorrhage study. 2015 , 24, 176-82	14
920	Relation between paradoxical decrease in high-density lipoprotein cholesterol levels after statin therapy and adverse cardiovascular events in patients with acute myocardial infarction. 2015 , 115, 411-6	9
919	Impact of low-grade albuminuria on left ventricular diastolic dysfunction. 2015 , 6, 13-16	2
918	Comparison of cystatin C and creatinine to determine the incidence of composite adverse outcomes in HIV-infected individuals. 2015 , 21, 84-9	6
917	Serum total bilirubin concentration is negatively associated with increasing severity of retinopathy in patients with type 2 diabetes mellitus. 2015 , 29, 218-21	23
916	Renoprotective effects of febuxostat in hyperuricemic patients with chronic kidney disease: a parallel-group, randomized, controlled trial. 2015 , 19, 1044-53	55
915	Renal histopathological findings in relation to ambulatory blood pressure in chronic kidney disease patients. 2015 , 38, 116-22	15
914	Current dietary salt intake of Japanese individuals assessed during health check-up. 2015 , 38, 163-8	12
913	Comparing GFR Estimating Equations Using Cystatin C and Creatinine in Elderly Individuals. 2015 , 26, 1982-9	101
912	Endothelial dysfunction is associated with the severity of cerebral small vessel disease. 2015 , 38, 291-7	36
911	Potential use of procalcitonin as biomarker for bacterial sepsis in patients with or without acute kidney injury. 2015 , 21, 257-63	33
910	Serum brain-derived neurotrophic factor level predicts adverse clinical outcomes in patients with heart failure. 2015 , 21, 300-6	21
909	Association between serum IgG4 concentrations and the morphology of the aorta in patients who undergo cardiac computed tomography. 2015 , 65, 150-6	3
908	HDL cholesterol as a predictor for the incidence of lower extremity amputation and wound-related death in patients with diabetic foot ulcers. 2015 , 239, 465-9	20
907	Effect of dipeptidyl peptidase-4 inhibitor, vildagliptin on plasminogen activator inhibitor-1 in patients with diabetes mellitus. 2015 , 115, 454-60	14

906	The relationships between visit-to-visit blood pressure variability and renal and endothelial function in chronic kidney disease. 2015 , 38, 193-8	15
905	Angiotensin II receptor blockers decrease serum concentration of fatty acid-binding protein 4 in patients with hypertension. 2015 , 38, 252-9	35
904	Efficacy and safety of ipragliflozin as an add-on to a sulfonylurea in Japanese patients with inadequately controlled type 2 diabetes: results of the randomized, placebo-controlled, double-blind, phase III EMIT study. 2015 , 6, 125-138	43
903	Ectopic expression of fatty acid-binding protein 4 in the glomerulus is associated with proteinuria and renal dysfunction. 2014 , 128, 345-51	32
902	Hyperhomocysteinemia is one of the risk factors associated with cerebrovascular stiffness in hypertensive patients, especially elderly males. 2014 , 4, 5663	42
901	Effectiveness and safety of tocilizumab therapy for patients with rheumatoid arthritis and renal insufficiency: a real-life registry study in Japan (the ACTRA-RI study). 2015 , 74, 627-30	26
900	CD34+/CD144+ circulating endothelial cells as an indicator of carotid atherosclerosis. 2015 , 24, 583-90	6
899	Different chronotherapeutic effects of valsartan and olmesartan in non-dipper hypertensive patients during valsartan treatment at morning. 2015 , 127, 62-8	8
898	Low-density lipoprotein apheresis for proteinuria in lupus nephritis with intraglomerular foam cells containing cholesterol crystals. <i>American Journal of Kidney Diseases</i> , 2015 , 65, 490-3	7.4 3
897	Use of simeprevir following pre-emptive pegylated interferon/ribavirin treatment for recurrent hepatitis C in living donor liver transplant recipients: a 12-week pilot study. 2015 , 22, 144-50	10
896	Association of a polymorphism of the interleukin 6 receptor gene with chronic kidney disease in Japanese individuals. 2015 , 20, 273-8	10
895	Characteristics of IgA nephropathy in advanced-age patients. 2015 , 47, 137-45	13
894	Association between smoking habits and the first-time appearance of atrial fibrillation in Japanese patients: Evidence from the Shinken Database. 2015 , 66, 73-9	21
893	Novel prediction model of renal function after nephrectomy from automated renal volumetry with preoperative multidetector computed tomography (MDCT). 2015 , 19, 974-81	10
892	Development and assessment of a simple scoring system for the risk of developing diabetic foot. 2015 , 6, 212-218	3
891	Association of arginine vasopressin surrogate marker urinary copeptin with severity of autosomal dominant polycystic kidney disease (ADPKD). 2015 , 19, 1199-205	18
890	Significance of estimated glomerular filtration rate in predicting brain or heart attacks in obese and non-obese populations. 2015 , 19, 790-6	4
889	Prognostic impact of serum bilirubin level on long-term renal survival in IgA nephropathy. 2015 , 19, 1062-70	9

888	Efficacy and safety of ipragliflozin as an add-on to pioglitazone in Japanese patients with inadequately controlled type 2 diabetes: a randomized, double-blind, placebo-controlled study (the SPOTLIGHT study). 2015 , 6, 104-116	40
887	Serum C-peptide to plasma glucose ratio may be associated with efficacy of vildagliptin in Japanese patients with type 2 diabetes mellitus. 2015 , 6, 197-205	2
886	Awareness of salt restriction and actual salt intake in hypertensive patients at a hypertension clinic and general clinic. 2015 , 37, 172-5	14
885	Ethnic disparity in prevalence of diabetic kidney disease in an Asian primary healthcare cluster. 2015 , 20, 216-23	14
884	Safety study of somatostatin analogue octreotide for autosomal dominant polycystic kidney disease in Japan. 2015 , 19, 746-52	4
883	Systemic arteriosclerosis and eating behavior in Japanese type 2 diabetic patients with visceral fat accumulation. 2015 , 14, 8	14
882	Effects of serum albumin and glycated albumin levels on performance of the Japanese GFR equation based on serum cystatin C. 2015 , 19, 626-30	
881	Relationship between low blood pressure and renal/cardiovascular outcomes in Japanese patients with chronic kidney disease under nephrologist care: the Gonryo study. 2015 , 19, 878-86	9
880	Validation of the Japanese histologic classification 2013 of immunoglobulin A nephropathy for prediction of long-term prognosis in a Japanese single-center cohort. 2015 , 19, 411-8	6
879	Impact of kidney function and urinary protein excretion on intima-media thickness in Japanese patients with type 2 diabetes. 2015 , 19, 909-17	8
878	High uric acid level is a risk factor for progression of IgA nephropathy with chronic kidney disease stage G3a. 2015 , 28, 451-6	20
877	Different association of cigarette smoking with GFR estimated from serum creatinine and that from serum cystatin C in the general population. 2015 , 19, 669-77	6
876	Urinary angiotensin-converting enzyme 2 increases in diabetic nephropathy by angiotensin II type 1 receptor blocker olmesartan. 2015 , 16, 159-64	20
875	Safety and efficacy of carbon dioxide and intravascular ultrasound-guided stenting for renal artery stenosis in patients with chronic renal insufficiency. 2015 , 66, 231-6	13
874	The urinary levels of prostanoid metabolites predict acute kidney injury in heterogeneous adult Japanese ICU patients: a prospective observational study. 2015 , 19, 1024-36	4
873	Evaluation of renal function in the elderly, not as easy as it seems: a review. 2015 , 27, 397-401	7
872	Underestimating chronic kidney disease by urine dipstick without serum creatinine as a screening tool in the general Japanese population. 2015 , 19, 474-80	5
871	Effect of empagliflozin monotherapy on postprandial glucose and 24-hour glucose variability in Japanese patients with type 2 diabetes mellitus: a randomized, double-blind, placebo-controlled, 4-week study. 2015 , 14, 11	79

870	Is pentraxin 3 a biomarker, a player, or both in the context of coronary atherosclerosis and metabolic factors?. 2015 , 30, 752-61	15
869	Risk factors associated with abnormal cognition in Japanese outpatients with diabetes, hypertension or dyslipidemia. 2015 , 6, 268-274	15
868	Association of aldosterone-to-renin ratio with hypertension differs by sodium intake: the Ohasama study. 2015 , 28, 208-15	8
867	Urinary ACE2 is associated with urinary L-FABP and albuminuria in patients with chronic kidney disease. 2015 , 75, 421-7	9
866	Off-pump bilateral skeletonized internal thoracic artery grafting in patients with chronic kidney disease. 2015 , 150, 315-21.e3	21
865	The PR interval and QRS duration could be predictors of renal function decline. 2015 , 240, 105-9	4
864	Changes over the last decade in carotid atherosclerosis in patients with end-stage kidney disease. 2015 , 240, 535-43	5
863	Correlation between UGT1A1 polymorphisms and raltegravir plasma trough concentrations in Japanese HIV-1-infected patients. 2015 , 21, 713-7	7
862	Estimated Glomerular Filtration Rate Variability Independently Predicts Renal Prognosis in Advanced Chronic Kidney Disease Patients. 2015 , 130, 256-62	9
861	Extension of the Calvert formula to patients with severe renal insufficiency. 2015 , 76, 53-9	8
860	Effects of a selective aldosterone blocker and thiazide-type diuretic on blood pressure and organ damage in hypertensive patients. 2015 , 37, 569-73	6
859	Effect of N- and T-type calcium channel blocker on proteinuria, blood pressure and kidney function in hypertensive patients: a meta-analysis. 2015 , 38, 847-55	16
858	Relationship between the brain natriuretic peptide (BNP) level and remission of diabetic nephropathy with microalbuminuria: a 3-year follow-up study. 2015 , 47, 138-44	6
857	Impact of the Triglycerides to High-Density Lipoprotein Cholesterol Ratio on the Incidence and Progression of CKD: A Longitudinal Study in a Large Japanese Population. <i>American Journal of Kidney Diseases</i> , 2015 , 66, 972-83	7.4 47
856	Pioglitazone treatment and cardiovascular event and death in subjects with type 2 diabetes without established cardiovascular disease (JDDM 36). 2015 , 109, 485-92	12
855	Associations between blood pressure responses to acute stress and impaired renal function and serum uric acid level. 2015 , 37, 656-60	2
854	Population Pharmacokinetics and Therapeutic Efficacy of Febuxostat in Patients with Severe Renal Impairment. 2015 , 96, 90-8	26
853	Hematocrit and the risk of cardiovascular disease in a Japanese community: The Hisayama Study. 2015 , 242, 199-204	29

852	Perioperative risk factors for acute kidney injury after liver resection surgery: an historical cohort study. 2015 , 62, 753-61	41
851	High-sensitivity C-reactive protein, but not white blood cell count, independently predicted incident diabetes in a Japanese health screening population. 2015 , 52, 983-90	8
850	Longitudinal Study of the Decline in Renal Function in Healthy Subjects. 2015 , 10, e0129036	53
849	MicroRNA-26a inhibits TGF- β -induced extracellular matrix protein expression in podocytes by targeting CTGF and is downregulated in diabetic nephropathy. 2015 , 58, 2169-80	73
848	Association of aging with glomerular filtration changes in cardiac rehabilitation participants with chronic kidney disease. 2015 , 187, 283-5	7
847	Impact of preprocedural high-sensitive C-reactive protein levels on long-term clinical outcomes of patients with stable coronary artery disease and chronic kidney disease who were treated with drug-eluting stents. 2015 , 66, 15-21	8
846	Magnesium modifies the association between serum phosphate and the risk of progression to end-stage kidney disease in patients with non-diabetic chronic kidney disease. 2015 , 88, 833-42	39
845	The relationship between serum calcium level and early-phase insulin secretion in normoglycemic and pre-diabetic individuals. 2015 , 123, 165-9	8
844	Telomere G-tail Length is a Promising Biomarker Related to White Matter Lesions and Endothelial Dysfunction in Patients With Cardiovascular Risk: A Cross-sectional Study. 2015 , 2, 960-7	9
843	Effect of estimated plasma volume reduction on renal function for acute heart failure differs between patients with preserved and reduced ejection fraction. 2015 , 8, 527-32	30
842	Day-by-Day Blood Pressure Variability and Functional Outcome After Acute Ischemic Stroke: Fukuoka Stroke Registry. 2015 , 46, 1832-9	54
841	Effect of Brazilian green propolis in patients with type 2 diabetes: A double-blind randomized placebo-controlled study. 2015 , 3, 355-360	45
840	Blood Pressure, Proteinuria, and Renal Function Decline: Associations in a Large Community-Based Population. 2015 , 28, 1150-6	31
839	Association of serum total bilirubin with renal outcome in Japanese patients with stages 3-5 chronic kidney disease. 2015 , 64, 1096-102	23
838	Metabolically healthy obesity and risk of incident CKD. 2015 , 10, 578-83	94
837	Association of and polymorphisms with ischemic stroke. 2015 , 3, 491-498	3
836	Association between hypouricemia and reduced kidney function: a cross-sectional population-based study in Japan. 2015 , 41, 138-46	37
835	C-peptide immunoreactivity index is associated with improvement of HbA1c: 2-Year follow-up of sitagliptin use in patients with type 2 diabetes. 2015 , 108, 441-7	7

834	Association between post-transplantation immunoglobulin A deposition and reduced allograft function. 2015 , 47, 332-6	7
833	The association between kidney function and cognitive decline in community-dwelling, elderly Japanese people. 2015 , 16, 349.e1-5	19
832	Sleep-Disordered Breathing in Patients with Polycystic Liver and Kidney Disease Referred for Transcatheter Arterial Embolization. 2015 , 10, 949-56	2
831	Prevalence of positive diffusion-weighted imaging findings and ischemic stroke recurrence in transient ischemic attack. 2015 , 24, 1000-7	11
830	Glycated albumin as a diagnostic tool for diabetes in a general Japanese population. 2015 , 64, 698-705	38
829	Serum cystatin C level is associated with left atrial enlargement, left ventricular hypertrophy and impaired left ventricular relaxation in patients with stage 2 or 3 chronic kidney disease. 2015 , 190, 287-92	5
828	Hybrid procedure in living donor liver transplantation. 2015 , 47, 679-82	18
827	Estimated functional renal parenchymal volume predicts the split renal function following renal surgery. 2015 , 33, 1571-7	15
826	The FRAX \square as a predictor of mortality in Japanese incident hemodialysis patients: an observational, follow-up study. 2015 , 33, 674-83	8
825	Efficacy of add-on therapy of aliskiren to an angiotensin II receptor blocker on renal outcomes in advanced-stage chronic kidney disease: a prospective, randomized, open-label study. 2015 , 19, 631-8	3
824	Impact of the new risk stratification in the 2011 Japanese Society of Nephrology clinical guidelines for IgA nephropathy on incidence of early clinical remission with tonsillectomy plus steroid pulse therapy. 2015 , 19, 646-52	9
823	Clinical significance of serum and urinary soluble urokinase receptor (suPAR) in primary nephrotic syndrome and MPO-ANCA-associated glomerulonephritis in Japanese. 2015 , 19, 804-14	10
822	The effect of tolvaptan on autosomal dominant polycystic kidney disease patients: a subgroup analysis of the Japanese patient subset from TEMPO 3:4 trial. 2015 , 19, 867-77	40
821	Renal biopsy findings and clinical indicators of patients with hematuria without overt proteinuria. 2015 , 19, 918-24	11
820	Benefits of a 12-week lifestyle modification program including diet and combined aerobic and resistance exercise on albuminuria in diabetic and non-diabetic Japanese populations. 2015 , 19, 1079-89	7
819	Hemoglobin concentration and incident metabolic syndrome: a population-based large-scale cohort study. 2015 , 50, 390-6	26
818	Efficacy and safety of empagliflozin monotherapy for 52 weeks in Japanese patients with type 2 diabetes: a randomized, double-blind, parallel-group study. 2015 , 32, 306-18	35
817	Clinical impact of thrombus aspiration during primary percutaneous coronary intervention in acute myocardial infarction with occluded culprit. 2015 , 30, 22-8	4

816	Low serum total bilirubin concentration in patients with type 1 diabetes mellitus complicated by retinopathy and nephropathy. 2015 , 6, 300-305	5
815	TWEAK/Fn14 system and crescent formation in IgA nephropathy. 2015 , 16, 27	12
814	Renal resistive index as an indicator of the presence and severity of anemia and its future development in patients with hypertension. 2015 , 16, 45	3
813	Associations between serum cholinesterase and incident hyper-LDL cholesterolemia, hypertriglyceridemia and hypo-HDL cholesterolemia as well as changes in lipid levels in a health screening population. 2015 , 241, 1-5	10
812	Clinical outcomes following percutaneous coronary intervention before and after introduction of drug-eluting stent. 2015 , 30, 338-46	
811	The therapeutic advantage of combination antihypertensive drug therapy using amlodipine and irbesartan in hypertensive patients: Analysis of the post-marketing survey data from PARTNER (Practical combination therapy of Amlodin and angiotensin II Receptor blocker; safety and efficacy in patients with hypertension) study. 2015 , 27, 542-50	4
810	Advanced stage of chronic kidney disease is risk of poor treatment outcome for smear-positive pulmonary tuberculosis. 2015 , 21, 559-63	7
809	Renal function evaluation in patients with cancer who were scheduled to receive carboplatin or S-1. 2015 , 19, 1107-13	6
808	Anti-albuminuric effects of spironolactone in patients with type 2 diabetic nephropathy: a multicenter, randomized clinical trial. 2015 , 19, 1098-106	36
807	Comparison of febuxostat and allopurinol for hyperuricemia in cardiac surgery patients with chronic kidney disease (NU-FLASH trial for CKD). 2015 , 66, 298-303	55
806	Negative associations between arterial stiffness parameter evaluated by cardio-ankle vascular index and serum low-density lipoprotein cholesterol concentration in early-stage atherosclerosis. 2015 , 66, 143-9	8
805	Comparison of the antidiabetic effects of linagliptin among groups with a normal renal function and a mild or severe renal impairment - retrospective observation study of Japanese patients with type 2 diabetes mellitus. 2015 , 16, 289-96	7
804	Plasma Renin Activity and Resting Heart Rate in a Population of Community-Dwelling Japanese: The Tanushimaru Study. 2015 , 28, 894-9	10
803	Ratio of urine and blood urea nitrogen concentration predicts the response of tolvaptan in congestive heart failure. 2015 , 20, 405-12	11
802	Absence of mesangial C1q deposition is associated with resolution of proteinuria and hematuria after tonsillectomy plus steroid pulse therapy for immunoglobulin a nephropathy. 2015 , 130, 1-7	4
801	Biomarkers in chronic kidney disease, from kidney function to kidney damage. 2015 , 4, 57-73	161
800	How does weight influence tenofovir disoproxil-fumarate induced renal function decline?. 2015 , 29, 643-5	2
799	Reply to 'how does weight influence tenofovir disoproxil-fumarate induced renal function decline?'. 2015 , 29, 645-7	

798	Reply to 'sexual networks, HIV, race and bacterial vaginosis'. 2015 , 29, 642-3	
797	Safety and efficacy of LCZ696, a first-in-class angiotensin receptor neprilysin inhibitor, in Japanese patients with hypertension and renal dysfunction. 2015 , 38, 269-75	56
796	Visit-to-visit variability and seasonal variation in blood pressure: Combination of Antihypertensive Therapy in the Elderly, Multicenter Investigation (CAMUI) Trial subanalysis. 2015 , 37, 411-9	14
795	Albuminuria and reduced glomerular filtration rate for predicting the renal outcomes in type 2 diabetic patients. 2015 , 20, 531-8	9
794	Relationships between the risk of cardiovascular disease in type 2 diabetes patients and both visit-to-visit variability and time-to-effect differences in blood pressure. 2015 , 29, 699-706	7
793	Association between estimated glomerular filtration rate and peripheral arterial disease. 2015 , 66, 430-4	17
792	Impact of Combined C-Reactive Protein and High-Density Lipoprotein Cholesterol Levels on Long-Term Outcomes in Patients With Coronary Artery Disease After a First Percutaneous Coronary Intervention. 2015 , 116, 999-1002	12
791	Association between Geriatric Nutrition Risk Index and The Presence of Sarcopenia in People with Type 2 Diabetes Mellitus: A Cross-Sectional Study. 2021 , 13,	1
790	Nutritional status during hospitalization is associated with the long-term prognosis of patients with heart failure. 2021 , 8, 5372	4
789	Bladder filling volume variation between the first and second day of planning computed tomography for prostate cancer radiation therapy and correlation with renal function. 2021 ,	0
788	Changes in serum uric acid levels as a predictor of future decline in renal function in older adults with type 2 diabetes. 2021 , 100, e27420	0
787	Concurrent chemoradiotherapy with cisplatin and docetaxel for head and neck squamous cell carcinoma. 2021 ,	
786	Impact of Area Under the Concentration-Time Curve on the Prevalence of Vancomycin-Induced Nephrotoxicity in Combination With Tazobactam/Piperacillin or Cefepime: A Single-Institution Retrospective Study. 2021 ,	1
785	Impact of switching to tenofovir alafenamide fumarate in patients with entecavir-treated chronic hepatitis B.. 2021 , 33, e898-e904	0
784	Effect of body mass index on the association between alcohol consumption and the development of chronic kidney disease. 2021 , 11, 20440	1
783	Effect of Cardiac Rehabilitation on Glomerular Filtration Rate Using Serum Cystatin C Concentration in Patients With Cardiovascular Disease and Renal Dysfunction. 2021 , 42,	
782	A weight regain of 1.5kg or more and lack of exercise are associated with nonalcoholic fatty liver disease recurrence in men. 2021 , 11, 19992	1
781	Estimated glomerular filtration rate may be an independent predictor for clinical outcomes regardless of acute kidney injury complication in the emergency department. 2021 , 16, e0258665	2

- 780 FAN score comprising fibrosis-4 index, albumin-bilirubin score and neutrophil-lymphocyte ratio is a prognostic marker of urothelial carcinoma patients treated with pembrolizumab. **2021**, 11, 21199 1
- 779 Lifestyle behaviors associated with the initiation of renal replacement therapy in Japanese patients with chronic kidney disease: a retrospective cohort study using a claims database linked with specific health checkup results. **2021**, 26, 102
- 778 Effect of Conventional Medical Therapy or Laparoscopic Sleeve Gastrectomy on Urinary Albumin in Japanese Subjects with Severe Obesity: An Observational Study. **2021**, 14, 613-621
- 777 Accuracy of the methods used to estimate glomerular filtration rate compared to 24-hour urinary creatinine clearance in patients with chronic spinal cord injury. **2021**, 1-6
- 776 Relationship between skin autofluorescence levels and clinical events in patients with heart failure undergoing cardiac rehabilitation. **2021**, 20, 208 1
- 775 A Novel Scoring System Based on Oxford Classification Indicating Steroid Therapy Use for IgA Nephropathy.. **2022**, 7, 99-107 0
- 774 Outcomes of open partial nephrectomy for renal cell carcinoma in the minimally invasive approach era.. **2021**, 15, 198-203
- 773 Safety and Efficacy of Sodium-glucose Cotransporter 2 Inhibitors in Kidney Transplant Recipients With Pretransplant Type 2 Diabetes Mellitus: A Retrospective, Single-center, Inverse Probability of Treatment Weighting Analysis of 85 Transplant Patients. **2021**, 7, e772 0
- 772 Creatinine reduction ratio on postoperative day 2 predicts long-term outcomes after living donor kidney transplantation. **2021**,
- 771 A Novel Prediction Model of Acute Kidney Injury Based on Combined Blood Variables in STEMI. **2021**, 1, 372-372 0
- 770 Association of a polymorphism of ROR2 and ischemic stroke in Japanese individuals with chronic kidney disease. **2010**, 1, 377-384 0
- 769 Evaluation of Japanese GFR Estimation Equation for Setting Initial Dose of Vancomycin-Comparison with Cockcroft & Gault equation-. **2010**, 36, 587-591 1
- 768 Clinical characteristics of 10 HIV-infected patients who developed end-stage renal disease. **2010**, 43, 581-586
- 767 Stroke Acute Management with Urgent Risk-factor Assessment and Improvement (SAMURAI) rt-PA Registry: General results and subanalyses. **2010**, 32, 756-761 6
- 766 What antihypertensive drugs are prescribed for patients with stroke in Japan? Ekuoka Stroke Registry Study□**2010**, 32, 275-281 1
- 765 Diabetic Nephropathy. 119-128
- 764 ??????20mL/min/1.73m2????????????Cr,Ccr,??GFR??GFR???. **2011**, 44, 55-58
- 763 Measurement of glomerular filtration rate by rapid intravenous injection of a newly developed inulin fraction. **2011**, 6, 9-15

- 762 Comparison of muscle strength between subjects with and without proteinuria. **2011**, 03, 698-702
- 761 Comparison of ventilatory threshold between subjects with and without proteinuria in Japanese. **2011**, 03, 394-399 1
- 760 4. ?????????? ?????????? **2011**, 44, 130-132
- 759 2. ?????????? ~?????????????????????. **2011**, 44, 126-127
- 758 3. ??????????. **2011**, 44, 128-129
- 757 Decreasing serum uric acid levels might be associated with improving estimated glomerular filtration rate (eGFR) in Japanese men. **2011**, 03, 498-503 1
- 756 Influence of Different Kidney Function Assessment Methods Used in Carboplatin Dose Setting on Thrombocytopenia. **2011**, 37, 649-652
- 755 Administration of Once-Daily Tacrolimus Extended-Release Formulation in Allogeneic Stem Cell Transplantation. **2012**, 1, 59-65
- 754 Reduced endogenous insulin secretion in diabetic patients with low-titer positive antibodies against GAD. **2012**, 02, 96-100 1
- 753 Evaluation of Predictability for Determining the Initial Dosing of Teicoplanin. **2012**, 38, 461-470 0
- 752 Analysis of the Clinical Covariates Associated with the Risks of Additional Percutaneous Coronary Intervention due to In-stent Restenosis or de Novo Lesions in Patients with Implanted Drug-eluting Stents. **2012**, 38, 795-803 1
- 751 Brachial-ankle pulse wave velocity (baPWV) in young adult recipients who underwent kidney transplantation during childhood. **2012**, 25, 114-119
- 750 Albuminuria-reducing effect of angiotensin II receptor blocker plus hydrochlorothiazide combination therapy in renal transplant recipients. **2012**, 4, 105-108
- 749 Histological and Morphometric Properties of Skeletonized Gastroepiploic Artery and Risk Factors for Intimal Hyperplasia. **2012**, 7, 191-194
- 748 Analysis of Risk Factors Influencing Development of Renal Failure with Cyclosporin A in Hematopoietic Stem Cell Transplantation Recipients. **2013**, 39, 45-51
- 747 ?? **2013**, 46, 1109-1118
- 746 1. ??????????????????. **2013**, 46, 1121-1124
- 745 Decreasing serum uric acid levels are associated with improving estimated glomerular filtration rate (eGFR) in Japanese women. **2013**, 03, 40-43

- 744 The fixed-dose combination of losartan/hydrochlorothiazide elicits potent blood pressure lowering during nighttime in obese hypertensive patients. **2014**, 6, 8-16 3
- 743 The number of antihypertensive agents simply reflects the grade and the risk of atherosclerosis in patients with type 2 diabetes mellitus. **2013**, 03, 161-167
- 742 Usefulness of change in estimated glomerular filtration rate as a predicting factor of progression of chronic kidney disease. **2013**, 2013, 351364
- 741 Declining Prevalence of Coronary Artery Disease in Incident Dialysis Patients Over the Past Two Decades. **2014**, 21, 3
- 740 Survey on the Usage of the Calvert Formula to Calculate AUC-based Carboplatin Dosing in Japan. **2015**, 41, 759-767 1
- 739 Prevalence and Characteristics of Treatment-Resistant Hypertension among Hypertensive Japanese Outpatients. **2015**, 06, 623-634 1
- 738 Clinical Significance of Renal Interstitial Fibrosis in Patients with Lupus Nephritis. **2015**, 61, 418-425
- 737 Clinical and Histological Characteristics in Patients with Non-IgA Mesangioproliferative Glomerulonephritis. **2015**, 61, 41-48
- 736 Pharmacoepidemiologic Investigation of the Risk Factors Associated with Hypocalcemia Induced by Denosumab. **2015**, 41, 578-584 1
- 735 The Prevention of Renal Dysfunction in High-risk CKD Patients Undergoing Carotid Artery Stenting. **2016**, 10, 183-189
- 734 Identification of the Risk Factors for Severe Neutropenia Induced by Pemetrexed Plus Carboplatin Chemotherapy. **2016**, 42, 278-285
- 733 Follow-up rates of living kidney donor in Japan: A single center study. **2016**, 26, 423-426 1
- 732 4. Chronic Kidney Disease Initiatives in Japan and Future Perspective. **2016**, 105, 1600-1610 1
- 731 Acute kidney injury following cardiac surgery under cardiopulmonary bypass. **2016**, 23, 141-147 1
- 730 4. Chronic Kidney Disease Initiatives in Japan and Future Perspective. **2016**, 105, 89a-94a
- 729 A Greater Association of Hyperuricemia than of Metabolic Syndrome with the New Incidence of Chronic Kidney Disease. **2016**, 06, 17-27
- 728 Prevention Strategies of Contrast Medium Induced Nephropathy (CIN): A Review of the Current Literature. **2016**, 06, 98-110 1
- 727 The Significance of Angiotensin Converting Enzyme Inhibitor or Angiotensin II Receptor Blocker Use in Sudden Cardiac Death. **2017**, 08, 496-503

- 726 Outcomes of the Re-classification of WHO Class IV Lupus Nephritis Using the ISN/RPS Classification: a Single Center Retrospective Observational Study from the JUDE Study. **2017**, 63, 362-369
- 725 Mean Corpuscular Volume is a Prognostic Factor for Patients after Curative Resection for Stage II Colorectal Cancer. **2017**, 78, 905-915
- 724 The Cluster of Abnormalities Related to Metabolic Syndrome Is Associated With Reduced Glomerular Filtration Rate and Raised Albuminuria in Patients With Type 2 Diabetes Mellitus. **2017**, 9, 759-764
- 723 Clinical Features of Appendiceal Diverticulitis Comparing with Acute Appendicitis. **2017**, 78, 2391-2397 0
- 722 A study of the effects of hypothermic circulatory arrest has on postoperative renal function in the total arch replacement. **2017**, 44, 14-18
- 721 Autologous Blood Donation Before Hepatectomy in Patients With Chronic Kidney Disease. **2021**, 105, 228-233
- 720 Effects of Once-Weekly Dulaglutide on Glycemic Control and Renal Function in Patients with Advanced-Stage Diabetic Nephropathy. **2017**, 9, 2
- 719 ??????????????????????????????????????. **2018**, 27, 43-48
- 718 Temporal Trends in Patients Characteristics and Clinical Outcomes in the Elderly Following Percutaneous Coronary Intervention. **2018**, 64, 379-385
- 717 Effects of Repaglinide Versus Glimpiride on 1,5-Anhydroglucitol and Glycated Hemoglobin Levels in Japanese Patients With Type 2 Diabetes. **2018**, 10, 838-842
- 716 Is High Central BP but Normal Office Brachial BP a risk?-The ABC-J II Study-. **2018**, 2, 66-73
- 715 [Renal senescence and Geriatric renal function]. **2018**, 55, 338-344
- 714 A Pilot Study of Prediction of Creatinine Clearance by Ellipsoid Volumetry of Kidney Using Noncontrast Computed Tomography. **2019**, 2, 60-66 2
- 713 Essential Points of Treatment and Examination to Prevent Diabetic Kidney Disease Aggravation. **2019**, 65, 529-536
- 712 Brain Natriuretic Peptide (BNP) and N-Terminal-proBNP in Cardio-Renal Anemia Syndrome □ Difference in Prognostic Ability □ **2019**, 1, 71-77 0
- 711 Changes in Serum and Urinary Potassium Handling Associated with Renin-Angiotensin-Aldosterone System Inhibitors in Advanced Chronic Kidney Disease Patients. **2019**, 11, e5561 1
- 710 Significant polar vasculosis in a patient with a 30-year history of type 2 diabetes. **2019**, 2019,
- 709 Clinicopathological Implications of Proteinuria after Long-Term Isolated Hematuria due to Thin Basement Membrane Nephropathy and Focal Segmental Glomerulosclerosis. **2019**, 2019, 1627392

708	Differential Effects of DPP-4 Inhibitors, Anagliptin and Sitagliptin, on PCSK9 Levels in Patients with Type 2 Diabetes Mellitus who are Receiving Statin Therapy. 2020 , 29,	0
707	Clinicopathological Analysis of Medullary Ray Injury in 1-Year Protocol Paediatric Renal Allograft Biopsies. 2020 , 144 Suppl 1, 79-85	1
706	Tongue thickness measured by ultrasonography is associated with tongue pressure in the Japanese elderly.	1
705	Refractory Proteinuria in a Patient with Lupus Nephritis.. 2020 , 1, 580-581	
704	Association between the Door-to-balloon Time and Mid-term Clinical Outcomes in Patients with ST-Segment Elevation Myocardial Infarction. 2020 , 59, 1597-1603	1
703	Lipoprotein (a) and the Risk of Chronic Kidney Disease in Hospitalized Japanese Patients. 2020 , 59, 1705-1710	0
702	Synthesis of Fructose Biosensors and Progressing Their Efficiency Using Californium Colloidal Nanoparticles for Detecting Fructose and Triglycerides. 2020 , 12, 1002-1017	
701	Sex Differences in the Effects of Weight Reduction on Future Blood Pressure Elevation in a Mildly Obese Middle-Aged Population. 2020 , 2, 385-392	1
700	The influence of gut microbiome on progression of overactive bladder symptoms: a community-based 3-year longitudinal study in Aomori, Japan. 2021 , 1	3
699	Changes in Prevalence and Health Checkup Coverage Rate of Chronic Kidney Disease (CKD) after Introduction of Prefecture-Wide CKD Initiative: Results of the Kagawa Association of CKD Initiatives. 2021 , 11,	0
698	Effectiveness of CKD Exacerbation Countermeasures in Izumo City. 2021 , 11,	1
697	Unique Habitual Food Intakes in the Gut Microbiota Cluster Associated with Type 2 Diabetes Mellitus. 2021 , 13,	1
696	Vegetable Preference and Prediction of Proteinuria: A Retrospective Cohort Study. 2021 , 1-7	1
695	Factors associated with difficulty in crossing the culprit lesion of acute myocardial infarction. 2021 , 11, 21403	1
694	Long-Term renal function after implantation of continuous-flow left ventricular assist devices: A single center study. 2021 , 37, 100907	
693	Can distinction between the renal cortex and outer medulla on ultrasonography predict estimated glomerular filtration rate in canine chronic kidney diseases?. 2020 , 21, e58	
692	Peripheral artery disease at the time of dialysis initiation and mortality: a prospective observational multicenter study. 2020 , 10, e042315	3
691	Association between sinusitis and relapse and changes in the myeloperoxidase-antineutrophil cytoplasmic antibody in microscopic polyangiitis. 2020 , 15, e0243572	0

690	Effect of Endovascular Treatment on Systemic Vascular Resistance in Patients with Lower-Limb Peripheral Artery Disease. 2020 , 13, 377-383	
689	Association of impaired oral health status with chronic kidney disease in post-acute rehabilitation. 2021 , 38, 300-307	0
688	Utilization of a three-dimensional printed kidney model for favorable TRIFECTA achievement in early experience of robot-assisted partial nephrectomy. 2020 , 9, 2697-2704	2
687	A fact-finding survey on self-efficacy of foot care behaviour in patients with diabetes: Analysis using the Diabetes Study from the Center of Tokyo Women's Medical University 2017 (DIACET 2017). 2021 , 4, e00219	1
686	Association between platelet distribution width and prognosis in patients with heart failure. 2020 , 15, e0244608	3
685	Prediction of long-term biochemical cure in patients with unilateral primary hyperaldosteronism treated surgically based on the early post-operative plasma aldosterone value. 2021 ,	
684	Effects of daily alcohol intake on glomerular filtration rate over three years. 2021 , 67, 1-7	1
683	Cardio-Ankle Vascular Index Predicts Post-Discharge Stroke in Patients with Heart Failure. 2021 , 28, 766-775	1
682	Reactive Hyperemia Index Associated With Atherosclerotic Cardiovascular Disease Under Treatment for Lifestyle Diseases. 2020 , 12, 293-299	4
681	Predictors of Stroke Outcome Extracted from Multivariate Linear Discriminant Analysis or Neural Network Analysis. 2020 , 29,	2
680	Renal outcome after hemopoietic stem cell transplantation. 2020 , 33, 115-122	
679	Renal function outcomes in the early and intermediate phases after radical cystectomy by ileal conduit. 2020 , 15, 178-182	0
678	Sedentary behavior, physical activity and renal function in middle-aged and older adults: isotemporal substitution modelling.	
677	Examination of Large Artery Atherosclerosis could Reveal Small Artery Retinopathy in Untreated Middle-Aged Individuals. 2020 ,	1
676	Perioperative Acute Kidney Injury Leading to Chronic Kidney Disease Following Major Abdominal Surgery: A Propensity Score-Matched Analysis. 2020 , 11, 9-19	
675	Diagnostic Accuracy of Single Spot Urine for Detecting Renal Uric Acid Underexcretion in Men. 2020 , 12, 443-447	
674	Effect of Long-term Peritoneal Dialysis on Change in Visceral Fat Area: A Single-Center Experience. 2020 , 30, 398-402	
673	Serum Digoxin Concentrations and Outcomes in Patients with Heart Failure and Atrial Fibrillation: A Single-Center Observational Study. 2020 , 51, 57-64	1

- 672 A Clinical Association between an Increasing Renal Resistive Index and the Atherosclerotic Burden in Patients with a Preserved Renal Function. **2020**, 59, 909-916 1
- 671 External Validation of a Prediction Model for Acute Kidney Injury Following Noncardiac Surgery. **2021**, 4, e2127362 0
- 670 Coexisting of aortic arch atheroma and atrial fibrillation for short-term recurrence and poor functional outcome in acute stroke. **2021**, 1 1
- 669 Health inequities and the inappropriate use of race in nephrology. **2021**, 7
- 668 Association between MR-proADM concentration and treatment intensity of antihypertensive agents in chronic kidney disease patients with insufficient blood pressure control. **2021**, 11, 21931
- 667 Kidney outcomes and all-cause mortality in people with type 2 diabetes exhibiting non-albuminuric kidney insufficiency. **2022**, 65, 234-245 1
- 666 Usefulness of subclassification of adult diabetes mellitus among inpatients in Japan. **2021**, 0
- 665 Excretion Patterns of Urinary Sediment and Supernatant Podocyte Biomarkers in Patients with CKD.. **2022**, 3, 63-73 0
- 664 Safety and efficacy of esaxerenone in Japanese hypertensive patients with heart failure with reduced ejection fraction: A retrospective study. **2021**, 16, e0259485 0
- 663 Association between remnant cholesterol and arterial stiffness: A secondary analysis based on a cross-sectional study. **2021**, 2
- 662 Mortality of inactive hepatitis B virus carriers in Japan is similar to that of the general population. **2021**, 52, 81 3
- 661 Heterozygous Urinary Abnormality-Causing Variants of and Affect Severity of Autosomal Recessive Alport Syndrome.. **2020**, 1, 936-942 0
- 660 Machine Learning Approach to Decision Making for Insulin Initiation in Japanese Patients With Type 2 Diabetes (JDDM 58): Model Development and Validation Study (Preprint).
- 659 Association of Fish Intake with Low-density Lipoprotein Cholesterol/high-density Lipoprotein Cholesterol Ratio in Apparently Healthy Males in Japanese. **2020**, 47, 507-515
- 658 Home device-monitored sleep blood pressure reflects large artery stiffness: the Nagahama study. **2020**, 38, 2459-2464 0
- 657 Brain Natriuretic Peptide (BNP) and N-Terminal-proBNP in Cardio-Renal Anemia Syndrome - Difference in Prognostic Ability. **2019**, 1, 71-77
- 656 Seasonal Variations in the Incidence of Ischemic Stroke, Extracranial and Intracranial Hemorrhage in Atrial Fibrillation Patients. **2020**, 84, 1701-1708 1
- 655 Serum IgG titers against periodontal pathogens are associated with cerebral hemorrhage growth and 3-month outcome. **2020**, 15, e0241205 1

654	Factors associated with poor clinical outcomes of ST-elevation myocardial infarction in patients with door-to-balloon time . 2020 , 15, e0241251	3
653	Impact of substratification on predicting oncological outcomes in patients with primary high-risk non-muscle-invasive bladder cancer who underwent transurethral resection of bladder tumor. 2020 , 38, 795.e9-795.e17	3
652	Clinical Decision Support System with Renal Dose Adjustment Did Not Improve Subsequent Renal and Hepatic Function among Inpatients: The Japan Adverse Drug Event Study. 2020 , 11, 846-856	1
651	Clinical impacts of cerebral microbleeds in patients with established coronary artery disease. 2021 , 39, 259-265	1
650	Geriatric Patients Are at a High Risk of Hypokalemia Associated with Yokukansan Preparation: A Retrospective Cohort Study. 2020 , 43, 1742-1748	1
649	Renal function and outcomes in atrial fibrillation patients after catheter ablation. 2020 , 15, e0241449	1
648	Time Course Changes in Urinary Angiotensinogen and Circulating N-Terminal Pro-B-Type Natriuretic Peptide in Patients Hospitalized with Acute Heart Failure. 2020 , 59, 2839-2847	
647	The influence of renal insufficiency on sleep-disordered breathing in patients with symptomatic chronic heart failure. 2010 , 15, 33-6	1
646	Comparison between Three Different Equations for the Estimation of Glomerular Filtration Rate in Omani Patients with Type 2 Diabetes Mellitus. 2014 , 14, e197-203	4
645	Frequency and predictors of bleeding complications associated with anti-coagulant therapy using dabigatran in Japanese patients with atrial fibrillation. 2014 , 4, 70-8	6
644	Carboplatin dosing for adult Japanese patients. 2014 , 76, 1-9	7
643	Long-term effects of tolvaptan in patients requiring recurrent hospitalization for heart failure. 2015 , 77, 355-62	1
642	Randomized control trial for the assessment of the anti-albuminuric effects of topiroxostat in hyperuricemic patients with diabetic nephropathy (the ETUDE study). 2016 , 78, 135-42	8
641	Extremity Radioactive Iodine Uptake on Post-therapeutic Whole Body Scan in Patients with Differentiated Thyroid Cancer. 2015 , 3, 26-34	3
640	Impaired Erythrocyte Deformability in Patients with Coronary Risk Factors: Significance of Nonvalvular Atrial Fibrillation. 2013 , 6, 939	
639	Comprehensive Analysis of Factors Affecting Post-partial Nephrectomy Renal Global Function. 2017 , 60, 94-100	
638	Ultrasound Assessment of Kidney Volume in Patients with Acute Decompensated Heart Failure: A Predictor of Diuretic Resistance. 2017 , 60, 135-144	1
637	Cystatin C is Indispensable for Evaluation of Kidney Disease. 2017 , 28, 268-276	15

- 636 Laboratory prognostic score for predicting 30-day mortality in terminally ill cancer patients. **2018**, 80, 571-582 4
- 635 Impact of the interferon- γ release assay and glomerular filtration rate on the estimation of active tuberculosis risk before bronchoscopic examinations: a retrospective pilot study. **2020**, 12, 5842-5849 1
- 634 Sex differences in clinical outcomes after rotational atherectomy of calcified coronary stenoses: from multicenter registry. **2021**, 11, 12-20
- 633 Effect of smoking on clinical outcomes in patients receiving rotational atherectomy in calcified coronary lesions: from the ROCK Registry, South Korea. **2021**, 41, 191-197
- 632 Initiation and long-term use of tolvaptan for patients with worsening heart failure through hospital and clinic cooperation. **2021**, 83, 431-441
- 631 The impact of physical activity on cardiovascular mortality in the general population. **2021**, 20, 1294-1304 0
- 630 Prevalence and Risk Factors of Silent Cerebral Microbleeds in Patients with Coronary Artery Disease. **2021**, 31, 106211 0
- 629 Effect of smoking on clinical outcomes in patients receiving rotational atherectomy in calcified coronary lesions: from the ROCK Registry, South Korea. **2021**, 41, 191-197
- 628 Sarcopenia Is Associated With a Risk of Mortality in People With Type 2 Diabetes Mellitus. **2021**, 12, 783363 0
- 627 Initiation of dialysis for kidney graft failure: A retrospective single-center cohort study. **2021**,
- 626 Comparison of multiparametric magnetic resonance imaging sequences with laboratory parameters for prognosticating renal function in chronic kidney disease. **2021**, 11, 22129 3
- 625 Physical Exercise Improves Cognitive Function in Older Adults with Stage 3-4 Chronic Kidney Disease: A Randomized Controlled Trial. **2021**, 1-11 1
- 624 Finerenone Dose-Exposure-Response for the Primary Kidney Outcome in FIDELIO-DKD Phase III: Population Pharmacokinetic and Time-to-Event Analysis. **2021**, 1 3
- 623 Time to rethink the current paradigm for assessing kidney function in drug development and beyond. **2021**, 0
- 622 Change in tartrate-resistant acid phosphatase isoform 5b levels, a marker of bone metabolism, in patients with chronic hepatitis B treated with tenofovir alafenamide.. **2022**, 16, 6
- 621 Clinicopathological discordance in biopsy-proven nephrosclerosis: a nationwide cross-sectional study of the Japan Renal Biopsy Registry (J-RBR). **2021**, 1 0
- 620 Echocardiographic Findings and Genotypes in Autosomal Dominant Polycystic Kidney Disease. 1-7 0
- 619 Association of Polypharmacy with Kidney Disease Progression in Adults with CKD. **2021**, 2

618	Validation of predictive model for new baseline renal function after robot-assisted partial nephrectomy or radical nephrectomy in Japanese patients. 2021 ,	
617	Impact of Concomitant Chronic Kidney Disease on Hospitalized Infections and Remission in Patients with Rheumatoid Arthritis: Results from the IORRA Cohort. 2021 ,	
616	Use of biologic agents and methotrexate improves renal manifestation and outcome in patients with rheumatoid arthritis: a retrospective analysis. 2021 , 1	1
615	Is Chronic Kidney Disease Affecting the Postoperative Complications of Vitrectomy for Proliferative Diabetic Retinopathy?. 2021 , 10,	0
614	Impact of Medication Adherence and Glycemic Control on the Risk of Micro/Macrovascular Diseases in Patients with Diabetes. 2021 ,	0
613	Enhanced prediction of renal function decline by replacing waist circumference with "A Body Shape Index (ABSI)" in diagnosing metabolic syndrome: a retrospective cohort study in Japan. 2021 ,	0
612	Association between intraoperative hyperglycemia and postoperative end-organ dysfunctions after cardiac surgery: a retrospective observational study. 2021 , 1	1
611	Chronic kidney disease and subclinical abnormalities of left heart mechanics in the community.	
610	Coronary angiography in patients with acute heart failure: from the KCHF registry. 2021 ,	1
609	Association Between Insulin Resistance, Oxidative Stress, Sympathetic Activity and Coronary Microvascular Function in Patients With Early Stage Impaired Glucose Metabolism. 2021 ,	0
608	Sodium-glucose cotransporter 2 inhibitors and anemia among diabetes patients in real clinical practice. 2021 ,	1
607	Serum uric acid level is associated with an increase in systolic blood pressure over time in female subjects: Linear mixed-effects model analyses. 2021 ,	0
606	Simple risk-score model for in-hospital major bleeding based on multiple blood variables in patients with acute myocardial infarction. 2022 , 346, 1-7	2
605	Tumor volume and tumor crossing of the axial renal midline predict renal function after robotic partial nephrectomy. 2021 , 11, 22526	0
604	Effect of chronic kidney disease on the association between hyperuricemia and new-onset hypertension in the general Japanese population: ISSA-CKD study. 2021 , 23, 2071	3
603	Association of Plasma Branched-Chain and Aromatic Amino Acids with Reduction in Kidney Function Evaluated in Apparently Healthy Adults. 2021 , 10,	0
602	A multiple 24-hour urine collection study indicates that kidney function decline is related to urinary sodium and potassium excretion in patients with chronic kidney disease. 2021 ,	1
601	Gender Differences in the Impact of Plasma Xanthine Oxidoreductase Activity on Coronary Artery Spasm. 2021 , 10,	0

- 600 Long-term impact of β -blocker in elderly patients without myocardial infarction after percutaneous coronary intervention. **2021**, 1
- 599 Prognostic effect of sleep-disordered breathing on hospitalized patients following acute heart failure. **2021**, 1
- 598 Transplant Prognosis in Kidney Transplant Recipients with Diabetes under Mycophenolic Acid-Focused Therapeutic Drug Monitoring. **2021**, 11,
- 597 Clinical Significance of Intermediate-Density Lipoprotein Cholesterol Determination as a Predictor for Coronary Heart Disease Risk in Middle-Aged Men. **2021**, 8, 756057
- 596 Assessment of Future Hypertension Risk by Sex Using Combined Body Mass Index and Waist-to-Height Ratio.. **2022**, 4, 9-16
- 595 Population Pharmacokinetic Analysis and Dosing Optimization of Prophylactic Fluconazole in Japanese Patients with Hematological Malignancy. **2021**, 7,
- 594 Combined impact of residual inflammatory risk and chronic kidney disease on long-term clinical outcomes in patients undergoing percutaneous coronary intervention. **2021**, 0
- 593 Association between enterocyte injury and fluid balance in patients with septic shock: a post hoc exploratory analysis of a prospective observational study. **2021**, 21, 293 0
- 592 Association between pulse pressure and progression of chronic kidney disease. **2021**, 11, 23275
- 591 Safety of Remdesivir for Patients 80 Years of Age or Older with Coronavirus Disease 2019 (COVID-19). **2021**, 38, 1067-1074 2
- 590 Drug-Drug Interaction between Azathioprine and Allopurinol in Patients with Chronic Kidney Disease: A Case Series Study. **2021**, 4, 170-174
- 589 Serum Potassium Levels of 4.5 to Less Than 5.0 mmol/L Are Associated with Better Vascular Function.. **2021**, 0
- 588 Visit-to-Visit Variability in Glomerular Filtration Before the Development of Atrial Fibrillation is Associated with Subsequent Progression of Chronic Kidney Disease: A Longitudinal Study Before and after the Development of Atrial Fibrillation.
- 587 Low plasma renin activity is independently associated with kidney disease progression in patients with type 2 diabetes and overt nephropathy, including those with impaired kidney function: a 2-year prospective study.. **2021**, 0
- 586 Measurement of glomerular filtration rate using endogenous d-serine clearance in living kidney transplant donors and recipients.. **2022**, 43, 101223 0
- 585 Different Kinetics of Activated Clotting Time among Uninterrupted Oral Anticoagulants during Catheter Ablation Procedure.. **2021**,
- 584 The Combination of High Levels of Adiponectin and Insulin Resistance Are Affected by Aging in Non-Obese Old Peoples.. **2021**, 12, 805244 0
- 583 Impact of nutritional status on prognosis in acute myocardial infarction patients undergoing percutaneous coronary intervention.. **2022**, 22, 3 0

582	Comparison of door-to-balloon time and in-hospital outcomes in patients with ST-elevation myocardial infarction between before versus after COVID-19 pandemic.. 2022 , 1	3
581	Impact of the interferon- γ release assay and glomerular filtration rate on the estimation of active tuberculosis risk before bronchoscopic examinations: a retrospective pilot study. 2020 , 12, 5842-5849	2
580	I. Clinical Evaluation of Kidney Function. 2020 , 109, 2466-2470	
579	Living and working environments are important determinants of glycemic control in patients with diabetes during the COVID-19 pandemic: a retrospective observational study.. 2022 ,	1
578	Successful introduction of off-pump coronary artery bypass grafting in Southeastern Asian countries: A single center's experience in Thailand.. 2022 , 8, 27-32	
577	Circulating serum amyloid A levels but not SAA1 variants predict long-term outcomes of angiographically confirmed coronary artery disease. 2022 ,	0
576	Efficacy and feasibility of robot-assisted partial nephrectomy for octogenarians: comparison with younger counterparts.. 2022 , 1	0
575	Association of perceived stress and coping strategies with the renal function in middle-aged and older Japanese men and women.. 2022 , 12, 291	
574	Long-Term Prognosis and Predictive Risk Factors for Polyvascular Disease in Patients with Peripheral Arterial Disease. 2022 , 12, 50-64	
573	Chronic Kidney Disease Patients Visiting Various Hospital Departments: An Analysis in a Hospital in Central Tokyo, Japan.. 2022 , 12,	1
572	HFA-PEFF scores: prognostic value in heart failure with preserved left ventricular ejection fraction.. 2021 ,	1
571	The Association of Postprandial Triglyceride Variability with Renal Dysfunction and Microalbuminuria in Patients with Type 2 Diabetic Mellitus: A Retrospective and Observational Study.. 2022 , 2022, 3157841	0
570	Inferior Vena Cava Size Predicts the Diuretic Response in Acute Decompensated Heart Failure Patients with Severe Aortic Stenosis: Sub-Analysis of the LOHAS Registry.	
569	Polypharmacy Influences the Renal Composite Outcome in Patients Treated with Sodium-Glucose Cotransporter 2 Inhibitors.. 2022 ,	1
568	Association of metabolic syndrome traits with urinary biomarkers in Japanese adults.. 2022 , 14, 9	0
567	Thigh Intramuscular Fat on Prognosis of Patients With Nonischemic Cardiomyopathy.. 2022 ,	1
566	Combined serum free light chain predicts prognosis in acute kidney injury following cardiovascular surgery.. 2022 , 44, 1-10	
565	JCS 2018 Guideline on Revascularization of Stable Coronary Artery Disease.. 2022 , 86,	3

- 564 Circulating level of FABP4 is an independent predictor of metabolic dysfunction-associated fatty liver disease in middle-aged and elderly individuals. **2021**, 0
- 563 External Evaluation of a Bayesian Warfarin Dose Optimization Based on a Kinetic-Pharmacodynamic Model.. **2022**, 45, 136-142 1
- 562 Urine albumin-to-creatinine ratio on admission predicts early rehospitalization in patients with acute decompensated heart failure.. **2022**, 1 0
- 561 Long-term social outcome after pediatric kidney transplantation: a single-center experience.. **2022**, 0
- 560 Issue of Waist Circumference for the Diagnosis of Metabolic Syndrome Regarding Arterial Stiffness: Possible Utility of a Body Shape Index in Middle-Aged Nonobese Japanese Urban Residents Receiving Health Screening.. **2022**, 1-10 0
- 559 Mineralocorticoid Receptor Antagonist Use and Hard Renal Outcomes in Real-World Patients With Chronic Kidney Disease.. **2022**, HYPERTENSIONAHA12118360 1
- 558 Serum Sulfatide Levels as a Biomarker of Active Glomerular Lesion in Patients with Anti-Neutrophil Cytoplasmic Antibody-Associated Vasculitis: A Single Center Pilot Study.. **2022**, 11, 0
- 557 Association between expanded criteria for living kidney donors and renal biopsy findings.. **2022**, 1 0
- 556 Evaluation of the Effect of Topiroxostat on Renal Function in Patients with Hyperuricemia: STOP-C Study, a Retrospective Observational Cohort Study.. **2022**, 1 1
- 555 Predictive Ability of Visit-to-Visit Variability of HbA1c Measurements for the Development of Diabetic Kidney Disease: A Retrospective Longitudinal Observational Study.. **2022**, 2022, 6934188 1
- 554 Sequential HBV treatment with tenofovir alafenamide for patients with chronic hepatitis B: week 96 results from a real-world, multicenter cohort study.. **2022**, 16, 282 0
- 553 Comparison of surgical outcomes after robot-assisted laparoscopic partial nephrectomy between patients continuing and discontinuing aspirin therapy: a Japanese single-centre study.. **2022**, 0
- 552 Clinical Outcomes of Everolimus With Reduced-Dose Tacrolimus vs Mycophenolate Mofetil With Standard-Dose Tacrolimus in De Novo ABO-Incompatible Kidney Transplant Recipients: 1-Year Follow-up.. **2022**, 0
- 551 Usefulness of the 2-year iodine-123 metaiodobenzylguanidine-based risk model for post-discharge risk stratification of patients with acute decompensated heart failure.. **2022**, 1 1
- 550 Daily behavioral and sleep patterns are associated with aging-induced male-specific disorders in individuals with reduced renal function.. **2022**, 161, 111717 0
- 549 Clinicopathologic Features of Mitochondrial Nephropathy.. **2022**, 7, 580-590 0
- 548 Association of urinary angiotensinogen with renal arteriolar remodeling in chronic kidney disease.. **2022**, 40, 0
- 547 Factors associated with atrial fibrillation in Japanese patients with type 2 diabetes mellitus: a cross-sectional study. 1

546	Imeglimin population pharmacokinetics and dose adjustment predictions for renal impairment in Japanese and Western patients with type 2 diabetes.. 2021 ,	3
545	Hyponatremia reflects mortality risks other than myocardial damage in acute coronary syndrome.. 2022 , 1	
544	Effects of luseogliflozin on the secretion of islet hormones and incretins in patients with type 2 diabetes.. 2022 ,	
543	Myeloperoxidase Antineutrophil Cytoplasmic Antibody-Associated Renal-Limited Vasculitis in a Young Adult Woman.. 2022 , 14, e21654	
542	Risk for Proteinuria in Newly Defined Hypertensive People Based on the 2017 American College of Cardiology/American Heart Association Blood Pressure Guideline.. 2022 ,	0
541	Impact of Lipoprotein(a) as a Residual Risk Factor in Long-Term Cardiovascular Outcomes in Patients With Acute Coronary Syndrome Treated With Statins.. 2022 ,	1
540	Impact of Female Sex on the Susceptibility to Hypernatremia Among Older Community-Dwelling Individuals in Japan.. 2022 , 15, 777-785	0
539	Clinical Characteristics and Outcomes of IgA Vasculitis in the Elderly.. 2022 ,	
538	Relationship between glomerular number in fresh kidney biopsy samples and light microscopy samples.. 2022 , 1	0
537	Long-term outcomes of the modest stent expansion strategy for the culprit lesion of acute myocardial infarction.. 2022 , 1	1
536	Cardio-Ankle Vascular Index is Associated with Prevalence and New-Appearance of Atrial Fibrillation in Japanese Urban Residents: A Retrospective Cross-Sectional and Cohort Study.. 2022 , 18, 5-15	1
535	Analysis of the variable factors affecting changes in the blood concentration of cyclosporine before and after transfusion of red blood cell concentrate.. 2022 , 8, 4	
534	Impact of Chronic Kidney Disease on the Associations of Cardiovascular Biomarkers With Adverse Outcomes in Patients With Suspected or Known Coronary Artery Disease: The EXCEED-J Study.. 2022 , e023464	0
533	Long-Term Outcomes of Longitudinal Efficacy Study With Tolvaptan in ADPKD.. 2022 , 7, 270-281	0
532	Cystatin C-based equations for estimating glomerular filtration rate do not require race or sex coefficients.. 2022 , 1-5	1
531	Association between kidney function and intracerebral hematoma volume.. 2022 , 96, 101-106	3
530	Relationship between eosinophils counts and muscle mass decline in older people with type 2 diabetes: A prospective study of the KAMOGAWA-DM cohort.. 2022 , 159, 111671	
529	Population pharmacokinetic analysis and dosage recommendations for digoxin in Japanese patients with atrial fibrillation and heart failure using real-world data.. 2022 , 23, 14	2

- 528 Temporal study of renal volume losses in patients with robotic partial nephrectomies.. **2022**,
- 527 Efficacy of combination therapy for childhood complicated focal IgA nephropathy.. **2022**, 1 0
- 526 Relationship of coffee consumption with a decline in kidney function among patients with type 2 diabetes: the Fukuoka Diabetes Registry.. **2022**,
- 525 Associations of Plasma Concentration Profiles of Dapagliflozin, a Selective Inhibitor of Sodium-Glucose Co-Transporter Type 2, with Its Effects in Japanese Patients with Type 2 Diabetes Mellitus.. **2022**, 15,
- 524 Development of chronic kidney disease influenced by serum urate and body mass index based on young-to-middle-aged Japanese men: a propensity score-matched cohort study.. **2022**, 12, e049540 1
- 523 Association between protein intake and changes in renal function among Japanese community-dwelling older people: The SONIC study.. **2022**,
- 522 Autopsy study examining non-chronic kidney disease versus chronic kidney disease caused by hypertensive-nephrosclerosis in elderly subjects.. **2022**, 1
- 521 Early Use of Everolimus as a Third Immunosuppressive Agent for Intestinal Transplantation: A Report of 2 Cases.. **2022**, 54, 472-472 0
- 520 Comparison of renal outcomes between sodium glucose co-transporter 2 inhibitors and glucagon-like peptide 1 receptor agonists.. **2022**, 185, 109231 0
- 519 Influence of stress induced by the first announced state of emergency due to coronavirus disease 2019 on outpatient blood pressure management in Japan.. **2021**,
- 518 Comparison between febuxostat and allopurinol uric acid-lowering therapy in patients with chronic heart failure and hyperuricemia: a multicenter randomized controlled trial.. **2021**, 49, 3000605211062770 2
- 517 Prediction Accuracy of Area under the Concentration-Time Curve of Vancomycin by Bayesian Approach Using Creatinine-based Equations of Estimated Kidney Function in Bedridden Elderly Japanese Patients.. **2022**,
- 516 Time-Series Change of Serum Soluble T-Cadherin Concentrations and Its Association with Creatine Kinase-MB Levels in ST-Segment Elevation Myocardial Infarction.. **2022**,
- 515 Clinical characteristics associated with renal impairment in children with spina bifida treated with clean intermittent catheterization. **2022**, 35,
- 514 Effects of Suprarenal Aortic Cross-Clamping and Adjunctive Renal Reconstruction on Outcomes of Open Abdominal Aortic Aneurysm Repair.. **2022**,
- 513 Association of Kidney Function with Serum Levels of Cholesterol Absorption and Synthesis Markers: The CACHE Study CKD Analysis.. **2022**,
- 512 In-hospital blood collection increases the rate of indeterminate results in interferon-gamma release assays.. **2022**, 16, 17534666221077817 1
- 511 Association between before-bedtime passive body heating and nocturia during the cold season among older adults.. **2022**,

510	Association of wound, ischemia, and foot infection clinical stage with frailty and malnutrition in chronic limb-threatening ischemia patients undergoing endovascular intervention.. 2022 , 17085381221076943	
509	Long-term Effectiveness of a Primary Care Practice Facilitation Program for Chronic Kidney Disease Management: An Extended Follow-up of a Cluster-Randomized FROM-J Study.. 2022 ,	1
508	Premature Atrial Contractions and Their Determinants in a General Population of Japanese Men.. 2022 ,	0
507	Potential effect of tolvaptan on polycystic liver disease for patients with ADPKD meeting the Japanese criteria of tolvaptan use.. 2022 , 17, e0264065	0
506	Comparison of the association between six different frailty scales and clinical events in patients on hemodialysis.. 2022 ,	0
505	Human Leukocyte Antigens and Biomarkers in Type 1 Diabetes Mellitus Induced by Immune-Checkpoint Inhibitors.. 2022 , 37, 84-95	3
504	Superiority of tenofovir alafenamide fumarate over entecavir for serum HBsAg level reduction in patients with chronic HBV infection: A 144-week outcome study after switching of the nucleos(t)ide analog.. 2022 , 17, e0262764	0
503	Eating Speed Is Associated with the Presence of Sarcopenia in Older Patients with Type 2 Diabetes: A Cross-Sectional Study of the KAMOGAWA-DM Cohort.. 2022 , 14,	1
502	Soluble Interleukin-2 Receptor Predicts Treatment Outcome in Patients With Autoimmune Tubulointerstitial Nephritis. A Preliminary Study.. 2022 , 9, 827388	
501	Dose-response relationships of sarcopenia parameters with incident disability and mortality in older Japanese adults.. 2022 ,	1
500	Self-assessment sheet submission rate predicts technique survival in patients initiating peritoneal dialysis.. 2022 ,	
499	Seasonal Changes in the Prevalence of Hyperkalemia in the Emergency Department: A Single Center Study.. 2022 , 58,	
498	Associations among Ratio of Free Triiodothyronine to Free Thyroxine, Chronic Kidney Disease, and Subclinical Hypothyroidism.. 2022 , 11,	0
497	Association between fluid infusions and the recovery from acute kidney injury in patients administered liposomal amphotericin B: a nationwide observational study.. 2022 , 44, 282-292	0
496	Framework for estimating renal function using magnetic resonance imaging.. 2022 , 9, 024501	
495	Cardiac computed tomography-derived myocardial tissue characterization after anthracycline treatment.. 2022 ,	0
494	Impact of inpatient educational programs on mortality after the start of dialysis therapy.. 2022 , 1	
493	Biopsy-proven CKD etiology and outcomes: Chronic Kidney Disease Japan Cohort (CKD-JAC) study.. 2022 ,	

- 492 Risk Factors of Proteinuria in Patients with Hepatocellular Carcinoma Receiving Lenvatinib.. **2022**, 45, 333-338
- 491 Increased intrarenal post-glomerular blood flow is a key condition for the development of calcineurin inhibitor-induced renal tubular acidosis in kidney transplant recipients.. **2022**, e14648 ○
- 490 Vesicoureteral reflux treatment following kidney transplantation potentially prevents graft function deterioration and allows long-term graft survival.. **2022**, ○
- 489 Association of circulating calciprotein particle levels with skeletal muscle mass and strength in middle-aged and older adults.. **2022**, ○
- 488 Effect of Teriparatide on Bone Mineral Density and Trabecular Bone Score in Type 2 Diabetic Patients with Osteoporosis: A Retrospective Cohort Study.. **2022**, 58, ○
- 487 Association of the serum albumin level with prognosis in chronic kidney disease patients.. **2022**, 1 ○
- 486 Predictors of hematoma as a complication in pediatric kidney biopsies.. **2022**, 64, e15189
- 485 Impact of Diabetes and KL-6 on COVID-19 Severity; A Single Center Study from Japan.. **2022**,
- 484 Urinary L-type fatty acid-binding protein is a predictor of cisplatin-induced acute kidney injury.. **2022**, 23, 125
- 483 Plasma and Urinary Levels of Nerve Growth Factor Are Elevated in Primary Hypertension.. **2022**, 2022, 3003269
- 482 Albuminuria predicts short-term worsening renal function after transcatheter aortic valve replacement.. **2022**, ○
- 481 The Impact of Postoperative Renal Function Recovery after Laparoscopic and Robot-Assisted Partial Nephrectomy in Patients with Renal Cell Carcinoma.. **2022**, 58, ○
- 480 Two-year outcomes of low-exposure extended-release tacrolimus and mycophenolate mofetil regimen in de novo kidney transplantation: A multi-center randomized controlled trial.. **2022**, e14655
- 479 Plasma Tsukushi Concentration Is Associated with High Levels of Insulin and FGF21 and Low Level of Total Cholesterol in a General Population without Medication.. **2022**, 12, ○
- 478 Association between lung function and the risk of atrial fibrillation in a nationwide population cohort study.. **2022**, 12, 4007 ○
- 477 Safety and effectiveness of empagliflozin in Japanese patients with type 2 diabetes: final results of a 3-year post-marketing surveillance study.. **2022**, ○
- 476 Lateral spine dual-energy X-ray absorptiometry and the risk of fragility fractures in long-term kidney graft recipients.. **2022**, 1 ○
- 475 Association of collateral flow with clinical outcomes in patients with acute myocardial infarction.. **2022**, 1 2

- 474 Effects of renin-angiotensin system inhibitors on the incidence of unplanned dialysis.. **2022,** ○
- 473 Predictive factors of postoperative fentanyl consumption in patients with inflammatory bowel disease: a retrospective cohort study.. **2022,** 22, 70 ○
- 472 Etiology and Outcome of Ischemic Stroke in Patients With Renal Impairment Including Chronic Kidney Disease: Japan Stroke Data Bank.. **2022,** ○
- 471 Increased Cerebral Small Vessel Disease Burden With Renal Dysfunction and Albuminuria in Patients Taking Antithrombotic Agents: The Bleeding With Antithrombotic Therapy 2.. **2022,** e024749 ○
- 470 Cystatin C-based estimated GFR performs best in identifying individuals with poorer survival in an unselected Chinese population: results from the China Health and Retirement Longitudinal Study (CHARLS).
- 469 Effect of Anagliptin versus Sitagliptin on Renal Function: Subanalyses from the REASON Trial.. **2022** , 15, 685-694
- 468 Overestimated Renal Function in Patients with Liver Cirrhosis Predicts Poor Prognosis.. **2022,** ○
- 467 Association between the Suita Score and Stroke Recurrence in Patients with First-ever Ischemic Stroke: A Prospective Cohort Study.. **2022,** 61, 773-780
- 466 Epicardial adipose tissue volume is an independent predictor of left ventricular reverse remodeling in patients with non-ischemic cardiomyopathy.. **2022,** ○
- 465 Association Between Urinary Protein-to-Creatinine Ratio and Chronic Kidney Disease Progression: A Secondary Analysis of a Prospective Cohort Study.. **2022,** 9, 854300 ○
- 464 Risk factors of proteinuria and potentially protective effect of renin-angiotensin system inhibitors in patients with renal cell carcinoma receiving axitinib.. **2022,** 1
- 463 Aldosterone-to-Renin Ratio Is Associated with Diabetic Nephropathy in Type 2 Diabetic Patients: A Single-Center Retrospective Study.. **2022,** 28, e935615
- 462 Association between urinary N-acetyl-β-D-glucosaminidase activity-urinary creatinine concentration ratio and risk of disability and all-cause mortality.. **2022,** 17, e0265637
- 461 Dynapenic abdominal obesity is associated with mild cognitive impairment in patients with cardiometabolic disease: a cross-sectional study.. **2022,** 22, 255
- 460 Impact of malnutrition on prognosis and coronary artery calcification in patients with stable coronary artery disease.. **2022,** 101185 ○
- 459 Prospective randomized comparison of effect on coronary endothelial and renal function between febuxostat and benzbromarone in hyperuricemic patients with coronary artery disease: EFEF study.. **2022,** 5, e563
- 458 Time to remission of proteinuria and incidence of relapse in patients with steroid-sensitive minimal change disease and focal segmental glomerulosclerosis: the Japan Nephrotic Syndrome Cohort Study.. **2022,** 1 ○
- 457 High Aspartate Aminotransferase/Alanine Aminotransferase Ratio May Be Associated with All-Cause Mortality in the Elderly: A Retrospective Cohort Study Using Artificial Intelligence and Conventional Analysis.. **2022,** 10, ○

- 456 Assessment of sarcopenia and malnutrition using estimated GFR ratio (eGFR_{cys}/eGFR) in hospitalised adult patients.. **2022**, 48, 456-463 ○
- 455 Usefulness of urinary tubule injury markers for predicting progression of renal dysfunction in patients with type 2 diabetes and albuminuria: The Fukuoka Diabetes Registry.. **2022**, 186, 109840 ○
- 454 Primary aldosteronism with mild autonomous cortisol secretion increases renal complication risk.. **2022**, ○ 1
- 453 Association between proteinuria and incident colorectal cancer: analysis of a nationwide population-based database.. **2022**, 12, e056250 ○
- 452 Protective Effect of Catheter Ablation of Atrial Fibrillation on the Renal Function in Patients With Hypertrophic Cardiomyopathy.. **2022**, ○
- 451 Evaluation of a newly proposed renal risk score for Japanese patients with ANCA-associated glomerulonephritis.. **2022**, 1 ○
- 450 Associations between continuous glucose monitoring-derived metrics and HbA1c in patients with type 2 diabetes mellitus.. **2022**, 109836 ○
- 449 Effect of the response to preoperative treatment for hepatorenal syndrome on the outcome of recipients of living-donor liver transplantation.. **2022**, ○ 1
- 448 Risk Assessment of Neutropenia during Low-Dose Valganciclovir Prophylaxis for Heart Transplant Recipients.. **2022**, 45, 452-459 ○
- 447 Effects of Nitric Oxide Synthase 3 Gene Polymorphisms on Cardiovascular Events in a General Japanese Population [The Yamagata (Takahata) Study] **2022**, ○
- 446 Safety and effectiveness of empagliflozin according to body mass index in Japanese patients with type 2 diabetes: a subgroup analysis of a 3-year post-marketing surveillance study.. **2022**, ○
- 445 Comparison of single trough-based area under the concentration-time curve versus trough concentration for the incidence of vancomycin-associated nephrotoxicity.. **2022**, ○ 1
- 444 Comparison of Outcomes of Patients With Acute Myocardial Infarction Between Chronic Total Occlusion Versus 90-99% Stenosis in Non-Culprit Arteries.. **2022**, ○
- 443 Probability of target attainment of oral antimicrobials for Escherichia coli and Klebsiella pneumoniae based on Monte Carlo simulations.. **2022**, 115662 ○
- 442 Albuminuria predicts worsening renal function after transcatheter aortic valve replacement.. **2021**, ○ 1
- 441 Hemoglobin concentration is associated with the hippocampal volume in community-dwelling adults.. **2022**, 101, 104668 ○
- 440 Coronary Microvascular Vasodilatory Function: Related Clinical Features and Differences According to the Different Coronary Arteries and Types of Coronary Spasm.. **2021**, 11, ○
- 439 Performance of creatinine-based equations for estimating glomerular filtration rate compared to endogenous creatinine clearance. **2021**, ○

- 438 The A Body Shape Index Might Be a Stronger Predictor of Chronic Kidney Disease Than BMI in a Senior Population.. **2021**, 18, 1
- 437 New longitudinal component of the RENAL nephrometry score for predicting the operative complexity in transperitoneal robot-assisted partial nephrectomy.. **2021**, 0
- 436 Cholesterol crystal embolism-related cerebral infarction: Magnetic resonance imaging and clinical characteristics.. **2021**, 25, 100388
- 435 Effect of sera from lupus patients on the glomerular endothelial fibrinolysis system.. **2021**, 64, e15099
- 434 Comparison of the usefulness of the cardio-ankle vascular index and augmentation index as an index of arteriosclerosis in patients with essential hypertension.. **2021**, 0
- 433 The relationship between day-to-day variability in home blood pressure measurement and multiple organ function.. **2021**, 0
- 432 Associations of Homocysteine with B Vitamins and Zinc in Serum Levels of Patients with Type 2 Diabetes Mellitus: A Cross-Sectional Study.. **2021**, 67, 417-423 1
- 431 Impact of citrus fruit intake on the mental health of patients with chronic heart failure.. **2021**, 1
- 430 Renal function in Japanese HIV-1-positive patients who switch to tenofovir alafenamide fumarate after long-term tenofovir disoproxil fumarate: a single-center observational study. **2021**, 18, 94 1
- 429 The Comparison of the Kidney Effects of Dipeptidyl Peptidase 4 Inhibitors and Glucagon-Like Peptide 1 Agonist-Administered Concomitant with Sodium-Glucose Cotransporter 2 Inhibitors in Japanese Patients with Type 2 Diabetes Mellitus and Chronic Kidney Disease.. **2021**, 2021, 6573369
- 428 Comparison of administration of single- and triple-course steroid pulse therapy combined with tonsillectomy for immunoglobulin A nephropathy.. **2021**, 100, e27778
- 427 Effects of mineralocorticoid receptor antagonists on sex hormones and body composition in patients with primary aldosteronism.. **2021**, 0
- 426 Management of bladder cancer in older patients.. **2021**, 2
- 425 Time-updated anion gap and cardiovascular events in advanced chronic kidney disease: a cohort study.. **2022**, 15, 929-936 1
- 424 Effect of Prophylactic Dose of Trimethoprim-Sulfamethoxazole on Serum Creatinine in Japanese Patients With Connective Tissue Diseases.. **2022**, 15, 11795441221085141 0
- 423 Comparison of two protocols for steroid pulse therapy in patients with IgA nephropathy: a retrospective observational study.. **2022**, 23, 153
- 422 The impact of the first announced state of emergency owing to coronavirus disease-2019 on stress and blood pressure levels among patients with type 2 diabetes mellitus in Japan.. **2022**,
- 421 High fibrosis-4 (FIB-4) index predicts new onset of ischemic heart disease during a 10-year period in a general population. 0

- 420 Clinical Characteristics Associated with the Development of Diabetic Ketoacidosis in Patients with Type 2 Diabetes.. **2022**, 61, 1125-1132 ○
- 419 Positive relationships between annual changes in salt intake and plasma B-type natriuretic peptide levels in the general population without hypertension and heart diseases.. **2022**, ○
- 418 Clinical features and predictors of outcome in patients with acute myocardial infarction complicated by out-of-hospital cardiac arrest.. **2022**, 22, 185 ○
- 417 Conventional medical therapy in heart failure patients eligible for the PARADIGM-HF, DAPA-HF, and SHIFT trials.. **2022**, ○
- 416 Relationship of body mass index to clinical outcomes after percutaneous coronary intervention.. **2022**, e13789 1
- 415 Severe hypokalemia in the emergency department: A retrospective, single-center study.. **2022**, 5, e594
- 414 The association of 5-year therapeutic responsiveness with long-term renal outcome in IgA nephropathy.. **2022**,
- 413 Assessing the Risk of Decrease in Kidney Function in Patients Prescribed Direct-Acting Antivirals for Hepatitis C Utilizing the MID-NET Medical Information Database Network in Japan.. **2022**, 1 ○
- 412 Development of a risk score model for 1-year graft loss after adult deceased donor liver transplantation in Japan based on a 20-year nationwide cohort. ○
- 411 Relationship between compliance with management target values and renal prognosis in multidisciplinary care for outpatients with chronic kidney disease.. **2022**, 1 ○
- 410 Data_Sheet_1.PDF. **2020**,
- 409 Table_1.DOCX. **2019**,
- 408 Clinical implications of severe sarcopenia in Japanese patients with acute heart failure.. **2022**, ○
- 407 Prevention of Contrast-Induced Nephropathy After Cardiovascular Catheterization and Intervention With High-Dose Strong Statin Therapy in Japan - The PREVENT CINC-J Study.. **2022**, 1
- 406 Association of serum fibroblast growth factor 21 with diabetic complications and insulin dose in patients with type 1 diabetes mellitus.. **2022**, 17, e0263774
- 405 Association between overweight and growth hormone secretion in patients with non-functioning pituitary tumors.. **2022**, 17, e0267324
- 404 Increased LDL cholesterol level is associated with deterioration of renal function in males. ○
- 403 Longitudinal relationship between long sleep duration and future kidney function decline.

- 402 Comparison of continuous subcutaneous insulin infusion treatment and multiple daily injection treatment on the progression of diabetic complications in Japanese patients with juvenile-onset type 1 diabetes mellitus.. **2022,**
- 401 Coronary Artery Bypass Grafting in Patients with Chronic Kidney Disease: Chronic Kidney Disease Has an Independent Adverse Effect on the Long-Term Outcome of Coronary Artery Bypass Grafting.. **2022, 2022, 4994970** ○
- 400 Changes in graft function after living donor kidney transplantation in children.. **2022, 1**
- 399 Difference between "Physical Fitness Age" Based on Physical Function and Chronological Age Is Associated with Obesity, Hyperglycemia, Depressive Symptoms, and Low Serum Albumin.. **2022, 26, 501-509** ○
- 398 Outcomes of Everolimus Plus Standard-Dose Tacrolimus Immunosuppression in De Novo Kidney Transplant: A Retrospective, Single-Center Study of 225 Transplants.. **2022, 20, 362-369**
- 397 Effects of antirejection therapies for early subclinical acute rejection in renal transplant protocol biopsies. **2022, 8,**
- 396 Early initiation of tolvaptan is associated with early discharge in patients with heart failure regardless of age.. **2022, 22, 202**
- 395 Antibody response to double SARS-CoV-2 mRNA vaccination in Japanese kidney transplant recipients.. **2022, 12, 6850** 1
- 394 Association between circulating CD34-positive cell count and height loss among older men.. **2022, 12, 7175** ○
- 393 A nationwide analysis of renal and patient outcomes for adults with lupus nephritis in Japan.. **2022,** ○
- 392 Risk factors for frailty in elderly Japanese people who received Ningen Dock: a cross-sectional study. **2022, 34,**
- 391 Relationship between physical function at discharge and hospital meal intake in elderly patients with heart failure.. **2022,**
- 390 Altered arginine vasopressin-cyclic AMP-aquaporin 2 pathway in patients with chronic kidney disease.. **2022, 1** ○
- 389 Association of Daily Physical Activity with Disability in Community-Dwelling Older Adults With/Without Chronic Kidney Disease.. **2022, 26, 521-528**
- 388 Impact of trimethoprim on serum creatinine, sodium, and potassium concentrations in patients taking trimethoprim-sulfamethoxazole without changes in glomerular filtration rate.. **2022,** 2
- 387 Early biomarkers for kidney injury in heat-related illness patients: a prospective observational study at Japanese Self-Defense Force Fuji Hospital.. **2022,**
- 386 Effects of different blood pressures and their long-term variability on the development of diabetic kidney disease in patients with type 2 diabetes mellitus.. **2022, 1-6**
- 385 Improved renal function in neurofibromatosis type 1 patients.

- 384 Kidney function at 3 months after acute kidney injury is an unreliable indicator of subsequent kidney dysfunction: the NARA-AKI Cohort Study.. **2022,** 0
- 383 Constipation Is a Frequent Problem Associated with Vascular Complications in Patients with Type 2 Diabetes: A Cross-sectional Study.. **2022,** 61, 1309-1317
- 382 Concomitant treatment with insulin and sodium-glucose cotransporter 2 inhibitors was associated with the renal composite outcome in Japanese patients with type 2 diabetes and chronic kidney disease: A propensity score-matched analysis.. **2022,** 0
- 381 Rationale, Design and Baseline Characteristics of the Effect of Canagliflozin in Type 2 Diabetic Patients with Microalbuminuria in Japanese Population (CANPIONE) study.. **2022,**
- 380 Diffusion-Weighted Imaging Hyperintensities in Acute and Subacute-Phase Intracerebral Hemorrhage.. **2022,** 31, 106549
- 379 Serum High-Sensitivity Cardiac Troponin T as an Independent Predictor for Incident Coronary Heart Disease in the Japanese General Population: The Circulatory Risk in Communities Study (CIRCS).. **2022,**
- 378 The Association Between Serum Uric Acid and Mortality in Patients with Acute Coronary Syndrome After Percutaneous Coronary Intervention.. **2022,** 63, 1
- 377 Transthyretin Amyloid Cardiomyopathy: Impact of Transthyretin Amyloid Deposition in Myocardium on Cardiac Morphology and Function. **2022,** 12, 792 0
- 376 Association of alcohol consumption with the incidence of proteinuria and chronic kidney disease: a retrospective cohort study in Japan.. **2022,** 21, 31 2
- 375 High fatty liver index is an independent predictor of ischemic heart disease during a 10-year period in a Japanese population.. **2022,** 0
- 374 Impact of hyperuricemia on coronary blood flow and in-hospital mortality in patients with acute myocardial infarction undergoing percutaneous coronary intervention.. **2022,** 0
- 373 Association between Vitamin Intake and Chronic Kidney Disease According to a Variant Located Upstream of the PTGS1 Gene: A Cross-Sectional Analysis of Shika Study. **2022,** 14, 2082 1
- 372 TG/HDL-C ratio as a predictor of stroke in the population with healthy BMI: The Jichi Medical School Cohort Study. **2022,** 0
- 371 Clinical and histopathological features related to time to complete remission in adult-onset minimal change nephrotic syndrome patients with corticosteroid treatment.. **2022,**
- 370 Potential association with malnutrition and allocation of combination medical therapies in hospitalized heart failure patients with reduced ejection fraction.. **2022,** 12, 8318 1
- 369 Impact of lenvatinib on renal function compared to sorafenib for unresectable hepatocellular carcinoma.. **2022,** 101, e29289
- 368 Association between proteinuria and retinal vein occlusion in individuals with preserved renal function: a retrospective cohort study.. **2022,** 0
- 367 Inverse Association of Skin Temperature With Ambulatory Blood Pressure and the Mediation of Skin Temperature in Blood Pressure Responses to Ambient Temperature.. **2022,** 101161HYPERTENSIONAHA12219190 0

- 366 Analysis of inflammatory cytokines and estimated glomerular filtration rate decline in Japanese patients with diabetic kidney disease: a pilot study.. **2022,**
- 365 Perioperative glycemic status is linked to postoperative complications in non-intensive care unit patients with type-2 diabetes: a retrospective study. **2022,** 13, 204201882210993 ○
- 364 Clinical, pathological, and genetic characteristics in patients with focal segmental glomerulosclerosis. 10.34067/KID.0000812022 ○
- 363 Glycated hemoglobin level on admission associated with progression to severe disease in hospitalized patients with non-severe coronavirus disease 2019. 1
- 362 Impact of activities of daily living on percutaneous coronary intervention and acute and long-term mortality in patients with acute myocardial infarction: Yamagata AMI registry. **2022,**
- 361 Effect of denosumab on renal function in women with osteoporosis evaluated using cystatin C. **2022,**
- 360 Structural modeling for Oxford histological classifications of immunoglobulin A nephropathy.
- 359 Metabolic dysfunction-associated fatty liver disease predicts new onset of chronic kidney disease better than does fatty liver or nonalcoholic fatty liver disease. 1
- 358 Prognosis and diastolic dysfunction predictors in patients with heart failure and recovered ejection fraction. **2022,** 12, ○
- 357 Effect of Using Concomitant Drugs on the Efficacy of Sodium Polystyrene Sulfonate. **2022,** 5, 33-38 1
- 356 Comparison of Long-Term Clinical Outcomes in Patients Stratified by a Novel Acute Myocardial Infarction Risk Stratification (nARS) System. **2022,** ○
- 355 Xanthine Oxidoreductase Activity in Marathon Runners: Potential Implications for Marathon-Induced Acute Kidney Injury. ○
- 354 Urinary free cortisol excretion is associated with lumbar bone density in patients with adrenal Cushing's syndrome.
- 353 Perioperative outcomes following robot-assisted partial nephrectomy for renal cell carcinoma according to surgeon generation. **2022,** 22,
- 352 Safety of anti-thrombotic therapy in acute aortic dissection; single center, observational study.
- 351 New Conversion Formula Between B-Type Natriuretic Peptide and N-Terminal-Pro-B-Type Natriuretic Peptide [Analysis From a Multicenter Study] **2022,** ○
- 350 Mortality and Cardiovascular Events in Patients With Chronic Kidney Disease and Sleep Apnea Syndrome. 9,
- 349 Dietary Fiber Intake Is Related to Skeletal Muscle Mass, Body Fat Mass, and Muscle-to-Fat Ratio Among People With Type 2 Diabetes: A Cross-Sectional Study. **2022,** 9, ○

- 348 A Comparative Study of Three Nutritional Risk/Screening Indices for Predicting Cardiac Events and Physical Functioning Among Patients with Acute Heart Failure. **2022**, 63, 541-549 1
- 347 Impact of Recovery from Acute Kidney Injury after Aortic Arch Repair. **2022**,
- 346 Cerebral and renal hemodynamics: similarities, differences, and associations with chronic kidney disease and aortic hemodynamics.
- 345 The correlation of atherosclerosis and triglyceride glucose index: a secondary analysis of a national cross-sectional study of Japanese. **2022**, 22, 1
- 344 Chronic kidney disease and clinical outcomes in patients with COVID-19 in Japan.
- 343 Association between Surgical Site Infection and Prophylactic Antimicrobial Cefmetazole Sodium Dose and Renal Function in Laparoscopic Colon Resection Surgery. **2021**, 36, 299-306
- 342 Different Clinical Courses of Nephronophthisis in Dizygotic Twins. **2022**,
- 341 Day-by-Day Home Blood Pressure Monitoring as a Biomarker in Diabetes. **2022**, 1-24
- 340 A Stratified Analysis of the Risk Associated With Low Body Mass Index for Patients After Percutaneous Coronary Intervention. **2022**,
- 339 Clinical and pathological characteristics of patients with acute kidney injury in Japan in whom kidney biopsy was performed: a cross-sectional analysis of the Japan Renal Biopsy Registry (J-RBR).
- 338 Elevated parathyroid hormone one year after kidney transplantation is an independent risk factor for graft loss even without hypercalcemia. **2022**, 23, 0
- 337 Chronic kidney disease is a key predictive factor for potential myocardial ischaemia and poor prognosis in asymptomatic patients with diabetes mellitus. **2022**, 12,
- 336 Insomnia in patients with acute heart failure: from the KCHF registry.
- 335 The Effect of Mild Renal Dysfunction on the Assessment of Plasma Amino Acid Concentration and Insulin Resistance in Patients with Type 2 Diabetes Mellitus. **2022**, 2022, 1-10
- 334 Blood pressure variability and the development of hypertensive organ damage in the general population.
- 333 Masticatory dysfunction in patients with diabetic neuropathy: A cross-sectional study. **2022**, 17, e0269594
- 332 Possible role of insulin resistance in activation of plasma xanthine oxidoreductase in health check-up examinees. **2022**, 12,
- 331 Body fluid regulation via chronic inhibition of sodium-glucose cotransporter-2 in patients with heart failure: a post hoc analysis of the CANDLE trial. 0

- 330 The renoprotective effect of once-weekly GLP -1 receptor agonist dulaglutide on progression of nephropathy in Japanese patients with type 2 diabetes and moderate to severe chronic kidney disease (JDDM67). ○
- 329 Development of a Laboratory Risk-Score Model to Predict One-Year Mortality in Acute Myocardial Infarction Survivors. **2022**, 11, 3497 ○
- 328 Predictors of early remission of proteinuria in adult patients with minimal change disease: a retrospective cohort study. **2022**, 12,
- 327 Kidney function deterioration is dependent on blood pressure levels: 11.2-year follow-up in diabetic patients.
- 326 A Long-term Estimated Glomerular Filtration Rate Plot Analysis Permits the Accurate Assessment of a Decline in the Renal Function by Minimizing the Influence of Estimated Glomerular Filtration Rate Fluctuations. **2022**, 61, 1823-1833
- 325 Switching to tenofovir alafenamide for nucleos(t)ide analogue-experienced patients with chronic hepatitis B: Week 144 results from a real-world, multicentre cohort study. ○
- 324 Relationships between cystatin C and creatinine-based eGFR with low tongue pressure in Japanese rural community-dwelling older adults.
- 323 Predictive ability of arterial stiffness parameters for renal function decline: a retrospective cohort study comparing cardio-ankle vascular index, pulse wave velocity and cardio-ankle vascular index0. Publish Ahead of Print, 1
- 322 The association between body mass index and all-cause mortality in Japanese patients with incident hemodialysis. **2022**, 17, e0269849 1
- 321 The Association of Controlling Nutritional Status Score and Prognostic Nutritional Index with Cardiovascular Diseases: the Fukuoka Kidney Disease Registry Study. **2022**,
- 320 Left Atrial Filling Time Index (LAFTI) Evaluated by 2D Speckle Tracking Echocardiography Predicts Poor Cardiac Prognosis in Patients with Heart Failure. **2022**,
- 319 Efficacy and Safety of Pemaibrate Versus Bezafibrate to Treat Patients with Hypertriglyceridemia: A Randomized Crossover Study. **2022**, ○
- 318 Streptococcal Infection-related Glomerulonephritis in an Elderly Diabetic Patient Complicated by Hemophagocytic Syndrome and Cytomegalovirus Nephritis. **2022**,
- 317 Association between Dietary Fat Intake and Hyperuricemia in Men with Chronic Kidney Disease. **2022**, 14, 2637 ○
- 316 Pharmacokinetic variability of vancomycin in patients with nosocomial meningitis. ○
- 315 Cerebral oxygenation and body mass index association with cognitive function in chronic kidney disease patients without dialysis: a longitudinal study. **2022**, 12, ○
- 314 Association of cardiovascular disease risk and changes in renin levels by mineralocorticoid receptor antagonists in patients with primary aldosteronism. 1
- 313 Impact of lifestyle and psychosocial factors on the onset of hypertension after the Great East Japan earthquake: a 7-year follow-up of the Fukushima Health Management Survey. ○

- 312 Initiating SGLT2 inhibitor therapy to improve renal outcomes for persons with diabetes eligible for an intensified glucose-lowering regimen: hypothetical intervention using parametric g-formula modeling. **2022**, 10, e002636
- 311 Combined ultrasound and magnetic resonance elastography predict hepatocellular carcinoma after hepatitis C virus eradication.
- 310 Urate Transporter ABCG2 Function and Asymptomatic Hyperuricemia: A Retrospective Cohort Study of CKD Progression. *American Journal of Kidney Diseases*, **2022**, 7-4 ○
- 309 The Association of Salt Intake and Non-alcoholic Fatty Liver Disease in People With Type 2 Diabetes: A Cross-Sectional Study. 9, 1
- 308 Association of serum total bilirubin levels with progressive renal decline and end-stage kidney disease: 10-year observational cohort study in Japanese patients with diabetes. **2022**, 17, e0271179 ○
- 307 Association of Increased Pulse Wave Velocity With Long-Term Clinical Outcomes in Patients With Preserved Ankle-Brachial Index After Acute Myocardial Infarction. **2022**,
- 306 Late Dialysis Modality Education Could Negatively Predict Peritoneal Dialysis Selection. **2022**, 11, 4042
- 305 Impact of Diabetes Duration on Clinical Outcome in Patients Receiving Rotational Atherectomy in Calcified Lesions in Korea Results from ROCK Registry. **2022**, 12, 993
- 304 Impact of Modified H&sub>2&/sub>FPEF Score on Chronic Limb-Threatening Ischemia in Patients With Lower Extremity Artery Disease Who Underwent Endovascular Therapy. **2022**,
- 303 Coronary artery lesion distribution in patients with chronic kidney disease undergoing percutaneous coronary intervention. **2022**, 44, 1098-1103
- 302 Exploratory analysis using machine learning of predictive factors for falls in type 2 diabetes. **2022**, 12, ○
- 301 Cystatin C-based eGFR predicts cardiovascular disease in patients with overweight/obesity and hyperglycemia.
- 300 Comparative proteomic analysis of glomerular proteins in primary and bucillamine-induced membranous nephropathy. **2022**, 19, ○
- 299 Association of ambulatory blood pressure with aortic valve and coronary artery calcification. **2022**, 40, 1344-1351
- 298 Factors linked to hepatocellular carcinoma development beyond 10 years after viral eradication in patients with hepatitis C virus. ○
- 297 Fibrosis-4 Index Is Closely Associated with Arterial Damage and Future Risk of Coronary Heart Disease in Type 2 Diabetes. **2022**, 2022, 1-8
- 296 Clinical significance of tumor markers in patients with type 2 diabetes: a retrospective observational study.
- 295 Clinical Benefit of Switching from Low-Dose to High-Dose Empagliflozin in Patients with Type 2 Diabetes. ○

- 294 Clinical impact of blood urea nitrogen, regardless of renal function, in heart failure with preserved ejection fraction. **2022**, 363, 94-101
- 293 A simplified prediction model for end-stage kidney disease in patients with diabetes. **2022**, 12,
- 292 Surgical and functional outcomes of robot-assisted laparoscopic partial nephrectomy for renal cell carcinoma in adolescents and young adults: a propensity score matching study.
- 291 Prognostic significance of cardiorenal dysfunction within 1 year after atrial fibrillation ablation in patients with systolic dysfunction.
- 290 Effect of canagliflozin on the decline of estimated glomerular filtration rate in chronic kidney disease patients with type 2 diabetes mellitus: A multicenter, randomized, double-blind, placebo-controlled, parallel-group, phase 3 study in Japan. ○
- 289 Renin-angiotensin system inhibitors may have an advantage over calcium channel blockers in reducing proteinuria in gastric cancer patients receiving ramucirumab. **2022**, 17,
- 288 Association between serum alkaline phosphatase and cardiovascular events in patients with atrial fibrillation.
- 287 The Importance of Continuous Glucose Monitoring-derived Metrics Beyond HbA1c for Optimal Individualized Glycemic Control. ○
- 286 Serum phosphate as an independent factor associated with cholesterol metabolism in patients undergoing hemodialysis: a cross-sectional analysis of the DREAM cohort.
- 285 Diverting ileostomy is a risk factor for renal impairment during CAPOX therapy. ○
- 284 Pharmacokinetics of Imeglimin in Caucasian and Japanese Healthy Subjects. ○
- 283 Detecting Sarcopenia Risk by Diabetes Clustering: A Japanese Prospective Cohort Study. ○
- 282 Clinical and pathological investigation of oligomeganephronia.
- 281 Percutaneous cryoablation combined with prior transcatheter arterial embolization for renal cell carcinomas of 3 cm or larger: a prospective study. ○
- 280 Characteristics of risk factors for acute kidney injury among inpatients administered sulfamethoxazole/trimethoprim: a retrospective observational study. **2022**, 8,
- 279 Early, Noninvasive Clinical Indicators of Kidney Prognosis in Primary Nephrotic Syndrome: A Retrospective Exploratory Study. **2022**, 2022, 1-7
- 278 Reduction of Severe Hypoglycemic Events Among Outpatients with Type 2 Diabetes Following Sodium-Glucose Cotransporter 2 Inhibitor Marketing in Japan.
- 277 Current antihypertensive treatment and treatment-resistant hypertension in Japanese patients with chronic kidney disease. 1

- 276 Low Insulin-Like Growth Factor-1 Levels Are Associated with High Comorbidity of Metabolic Diseases in Subjects with Severe Obesity.
- 275 Impact of Inflammation on Intra-individual Variation in Trough Voriconazole Concentration in Patients with Hematological Malignancies. **2022**, 45, 1084-1090
- 274 Clinical Impact of a Pharmacist-Driven Prospective Audit with Intervention and Feedback on the Treatment of Patients with Bloodstream Infection. **2022**, 11, 1144
- 273 Prognostic impact of urine cyclic AMP levels in patients with chronic kidney disease. ○
- 272 Robot-assisted laparoscopic versus open partial nephrectomy for renal cell carcinoma in patients with severe chronic kidney disease. ○
- 271 Prognosis and risk factors of chronic kidney disease progression in patients with diabetic kidney disease and non-diabetic kidney disease: a prospective cohort CKD-ROUTE study. **2022**, 44, 1309-1318 1
- 270 Complete Right Bundle Branch Block as a Predictor of Cardiovascular Events in Type 2 Diabetes. **2022**, 11, 4618
- 269 Household Income Is Related to Dietary Fiber Intake and Dietary Acid Load in People with Type 2 Diabetes: A Cross-Sectional Study. **2022**, 14, 3229
- 268 Analyses of operative time according to procedure phases during robot-assisted laparoscopic partial nephrectomy using the iPhone application My Intuitives□ ○
- 267 Long-term outcomes of percutaneous revascularization for internal mammary artery□left anterior descending artery bypass failure.
- 266 Correlation between albuminuria and interstitial injury marker reductions associated with SGLT2 inhibitor treatment in diabetic patients with renal dysfunction. **2022**, 27,
- 265 Rapid decrease in eGFR with concomitant use of tyrosine kinase inhibitors and renin□aldosterone□angiotensin system inhibitors in patients with chronic myelogenous leukemia.
- 264 Health-related behavioral changes and incidence of chronic kidney disease: The Japan Specific Health Checkups (J-SHC) Study.
- 263 Tacrolimus Use and Renal Function in Pregnancy with Lupus Nephritis: Analysis of Post-Marketing Surveillance Data in Japan.
- 262 The effect of preoperative sodium-glucose cotransporter 2 inhibitors on the incidence of perioperative metabolic acidosis: A retrospective cohort study. **2022**, 22, ○
- 261 Risk score-based substratification improves surveillance costs after transurethral resection of bladder tumor in patients with primary high-risk non-muscle-invasive bladder cancer. **2022**, 12,
- 260 Effect of Tolvaptan in Patients with Chronic Kidney Disease Stage G5, and Impact of Concomitant Use of Thiazide Diuretics: A Retrospective Cohort Study.
- 259 The moderate-intensity continuous exercise maintains renal blood flow and does not impair the renal function. **2022**, 10, 1

258	The significance of presepsin in chronic kidney disease with sepsis.	
257	Lower body mass index potentiates the association between skipping breakfast and prevalence of proteinuria. 13,	0
256	Randomized, double-blind, placebo-controlled phase 3 study of bardoxolone methyl in patients with diabetic kidney disease: Design and baseline characteristics of AYAME study.	1
255	Cardio-ankle vascular index is more closely associated than brachial-ankle pulse wave velocity with arterial damage and risk of cardiovascular disease in patients with diabetes. 2022, 22,	1
254	Population pharmacokinetics of everolimus in adult liver transplant patients: Comparison to tacrolimus disposition and extrapolation to pediatrics.	
253	Population kinetic-pharmacodynamic analysis of serum potassium in patients receiving sulfamethoxazole/trimethoprim.	
252	Plasma mid-regional pro-adrenomedullin: A biomarker of the ischemic penumbra in hyperacute stroke.	
251	Association between hemoglobin and chronic kidney disease progression: a secondary analysis of a prospective cohort study in Japanese patients. 2022, 23,	
250	Safety and effectiveness of empagliflozin and linagliptin fixed-dose combination therapy in Japanese patients with type 2 diabetes: final results of a one-year post-marketing surveillance study. 1-11	
249	The diversity and abundance of gut microbiota are associated with the pain sensation threshold in the Japanese population. 2022, 173, 105839	
248	Cystatin C-Based eGFR Predicts Post-Treatment Kidney Prognosis in Patients with Severe Obstructive Nephropathy. 2022, 2, 474-481	0
247	Prognostic Impact of Renal Dysfunction at 1-Year Follow-Up on Clinical Outcomes After Percutaneous Coronary Intervention. 2022,	0
246	Effect of Atrial Fibrillation on Renal Function: Time of Change-Point Analysis of the Estimated Glomerular Filtration Rate Before and after Development of Atrial Fibrillation.	0
245	Cutoff Values of Brachial-Ankle Pulse Wave Velocity for Atherosclerotic Risks by Age and Sex in the Japanese General Population. 2022,	0
244	Paradoxical Long-Term Impact Between Serum Apolipoprotein E and High-Density Lipoprotein Cholesterol in Patients Undergoing Percutaneous Coronary Intervention. 2022,	0
243	Percutaneous Thermal Ablation for Managing Small Renal Metastatic Tumors. 2022,	0
242	Heart Failure Re-admission and Mortality in Chronic Kidney Disease Patients after Transcatheter Aortic Valve Replacement in Asian Population. 2022, 4, 23-34	0
241	Relationship between Diabetes Mellitus and Serum Lathosterol and Campesterol Levels: The CACHE Study DM Analysis. 2022,	0

- 240 Differences in muscle strength, physical activity, and cardiometabolic risk factors between type 2 diabetic patients with and without benzodiazepines or antipsychotic medications. **2022**, 6, 167 ○
- 239 Association between the intake of plant and animal proteins and the serum fibroblast growth factor-23 level in patients with chronic kidney disease analyzed by the isocaloric substitution model. **2022**, ○
- 238 Association between Serum Magnesium Levels and Mortality in a Community-Based Population: The Yamagata (Takahata) Study. **2022**, 68, 270-275 ○
- 237 Long-term impact of renin-angiotensin system inhibitors for secondary prevention in patients with chronic kidney disease who underwent percutaneous coronary intervention. ○
- 236 Association between Time-Updated Eosinophil Counts and Progression of CKD. ○
- 235 Effects of Branched-Chain Amino Acids on Skeletal Muscle, Glycemic Control, and Neuropsychological Performance in Elderly Persons with Type 2 Diabetes Mellitus: An Exploratory Randomized Controlled Trial. **2022**, 14, 3917 ○
- 234 Efficacy and Safety of Esaxerenone in Hypertensive Patients with Diabetic Kidney Disease: A Multicenter, Open-Label, Prospective Study. ○
- 233 Post-Operative Kidney Function Using Deep Hypothermic Circulatory Arrest (DHCA) in Aortic Arch Operation. Volume 15, 239-252 ○
- 232 The Association Between Hypouricemia and Cardiometabolic Diseases. Publish Ahead of Print, ○
- 231 Associated factors and effects of comorbid atrial fibrillation in hypertensive patients due to primary aldosteronism. ○
- 230 Association between 3-Year Repetitive Isolated Hematuria and eGFR Deterioration in an Apparently Healthy Population: A Retrospective Cohort Study. **2022**, 19, 11466 ○
- 229 Reduced renal function is associated with prolonged hospitalization in frail older patients with non-severe pneumonia. 9, ○
- 228 Comparison of the blood pressure management between sodium-glucose cotransporter 2 inhibitors and glucagon-like peptide 1 receptor agonists. **2022**, 12, ○
- 227 Use of Anti-Thrombotic Drugs and In-Hospital Mortality in Acute Aortic Dissection Patients. **2022**, 12, 2322 ○
- 226 Potential progression biomarkers of diabetic kidney disease determined using comprehensive machine learning analysis of non-targeted metabolomics. **2022**, 12, ○
- 225 Tubulointerstitial B-cell infiltration and tertiary lymphoid tissue in adult-onset immunoglobulin A vasculitis with nephritis. ○
- 224 Association between Distigmine-induced Adverse Events and Serum Butyrylcholinesterase Level: A Multicenter Retrospective Study. **2022**, 142, 993-998 ○
- 223 Glomerular filtration in the aging population. 9, ○

- 222 Association of the estimated glomerular filtration rate (eGFR) and/or proteinuria to predict the risk of initiation of dialysis in people with and without diabetes. 0
- 221 Estimated Glomerular Filtration Rate and Hearing Impairment in Japan: A Longitudinal Analysis Using Large-Scale Occupational Health Check-Up Data. **2022**, 19, 12368 0
- 220 Comparison between the impact of tenofovir alafenamide and that of abacavir on rapid kidney function decline: A retrospective observational study. 0
- 219 Machine Learning-Based Model for Estimating Vancomycin Maintenance Dose to Target the Area under the Concentration Curve of 400-600 mg \cdot h/L in Japanese Patients. **2022**, 45, 1332-1339 1
- 218 Risk factors for pre-heart failure or symptomatic heart failure based on NT-proBNP. 0
- 217 Structural modeling for Oxford histological classifications of immunoglobulin A nephropathy. **2022**, 17, e0268731 0
- 216 The renoprotective effect of esaxerenone independent of blood pressure lowering: a post hoc mediation analysis of the ESAX-DN trial. 0
- 215 Liver steatosis and fibrosis markers association with cardiovascular and renal damage in Japanese adults: the TMM BirThree cohort study. **2022**, 100761 0
- 214 Subcutaneous Fat Thickness of the Lower Limb is Associated with Trunk Muscle Mass in Patients with Hepatocellular Carcinoma: A Simple Assessment for Sarcopenia Using Conventional Ultrasonography. **2022**, 0
- 213 Day-by-Day Home Blood Pressure Monitoring as a Biomarker in Diabetes. **2023**, 705-727 0
- 212 Impact of D-dimer for pathologic differentiation on transesophageal echocardiography in embolic stroke of undetermined source: a single-center experience. **2022**, 22, 0
- 211 Associations between nephron number and podometrics in human kidneys. **2022**, 0
- 210 Urinary A- and C-megalin predict progression of diabetic kidney disease: an exploratory retrospective cohort study. **2022**, 108312 0
- 209 Developing a common data model approach for DISCOVER CKD: A retrospective, global cohort of real-world patients with chronic kidney disease. **2022**, 17, e0274131 0
- 208 Health-related behavioral changes and incidence of chronic kidney disease: The Japan Specific Health Checkups (J-SHC) Study. **2022**, 12, 0
- 207 Effect of Chronic Kidney Disease or Anemia or Both on Cardiovascular Mortality in a 25-Year Follow-Up Study of Japanese General Population (From NIPPON DATA90). **2022**, 0
- 206 The association between hypothyroidism and proteinuria in patients with chronic kidney disease: a cross-sectional study. **2022**, 12, 1
- 205 Analysis of long-term renal function in patients with malignant solid tumors: Retrospective analysis using the estimated glomerular filtration rate. 0

- 204 MOONLIGHT study: the design of a comparative study of the effectiveness of belimumab in patients with a history of lupus nephritis from the post-Marketed effectiveness of belimumab cOhOrt and JapaN Lupus Natlonwide reGistry (LUNA) coHorT. **2022**, 9, e000746 ○
- 203 Potential link between high FIB-4 score and chronic kidney disease in metabolically healthy men. **2022**, 12, ○
- 202 Association between high-sensitivity cardiac troponin T levels and incident stroke in the elderly Japanese population: Results from the Tohoku Medical Megabank Community-based Cohort Study. **2022**, 22, 100212 ○
- 201 Non-recovery of vancomycin-associated nephrotoxicity is related to worsening survival outcomes: Combined retrospective analyses of two real-world databases. ○
- 200 The effect of the Kasuya CKD network on prevention of the progression of chronic kidney disease: successful collaboration of a public health service, primary care physicians and nephrologists community based cohort study. ○
- 199 Combined impacts of low apolipoprotein A-I levels and reduced renal function on long-term prognosis in patients with coronary artery disease undergoing percutaneous coronary intervention. **2022**, ○
- 198 Characterization of plasma daptomycin in patients with serum highly glycosylated albumin and obesity. **2022**, ○
- 197 Clues to mortality trends and their related factors in IgG4-related disease: A Japanese single-center retrospective study. ○
- 196 Dietary salt intake predicts future development of metabolic syndrome in the general population. ○
- 195 Use of the Visceral Adiposity Index as an Indicator of Chronic Kidney Disease in Older Adults: Comparison with Body Mass Index. **2022**, 11, 6297 1
- 194 The Impact of Myocardial Bridging on the Coronary Functional Test in Patients with Ischaemia with Non-Obstructive Coronary Artery Disease. **2022**, 12, 1560 ○
- 193 Impact of intravascular ultrasound-incomplete stent apposition on stent failure. ○
- 192 Change in fatty acid composition of plasma triglyceride caused by a 2 week comprehensive risk management for diabetes: A prospective observational study of type 2 diabetes patients with supercritical fluid chromatography/mass spectrometry-based semi-target lipidomic analysis. ○
- 191 Inadequate sleep duration may attenuate the anti-inflammatory effects of fish consumption in a healthy Japanese population: A cross-sectional study. 1-26 ○
- 190 Effects of one-hour discussion on the choice of dialysis modality at the outpatient clinic: A retrospective cohort study using propensity score matching. ○
- 189 Association of chronic kidney disease with total and site-specific cancer incidence in participants of the Japan Public Health Center-based Prospective Study. ○
- 188 Impact of kidney function and kidney volume on intracranial aneurysms in patients with autosomal dominant polycystic kidney disease. **2022**, 12, ○
- 187 Impact of Albuminuria on the Prognosis After Transcatheter Aortic Valve Implantation. **2022**, ○

- 186 Achievement of target blood pressure among community residents with hypertension and factors associated with therapeutic failure in the northern territory of Japan. **2022**, 50, 030006052211268 ○
- 185 CD44 as a pathological marker for the early detection of calcineurin inhibitor-induced nephrotoxicity post kidney transplantation. **2022**, 43, 181-186 ○
- 184 Prognostic Value of Novel Natriuretic Peptide Index After Percutaneous Coronary Intervention. **2022**, ○
- 183 Cardio-Ankle Vascular Index as an Arterial Stiffness Marker Improves the Prediction of Cardiovascular Events in Patients without Cardiovascular Diseases. **2022**, 9, 368 ○
- 182 Relationship between serum indirect bilirubin levels and skeletal muscle mass in older male and female patients with type 2 diabetes. **2022**, 17, e0276976 ○
- 181 Comparison of glomerular filtration rate estimating formulas among Japanese adults without kidney disease. **2022**, ○
- 180 Long term anti-vascular endothelial growth factor receptor treatment impairs renal function in renal cell carcinoma. ○
- 179 Association between prevalence of laxative use and history of bone fractures and cardiovascular diseases in patients with chronic kidney disease: the Fukuoka Kidney disease Registry (FKR) study. ○
- 178 Effects and safety of high-frequency rTMS in acute intracerebral hemorrhage patients: A pilot study. **2022**, 443, 120473 1
- 177 Blood pressure measurement in the elderly with atrial fibrillation: an observational study comparing different noninvasive sphygmomanometers. **2022**, 13, 204062232211370 ○
- 176 Association between circulating SerpinB1 levels and insulin sensitivity in Japanese with type 2 diabetes: A single-center, cross-sectional, observational study. **2022**, 17, e0276915 ○
- 175 Persistence and Safety of Golimumab in Elderly Patients with Rheumatoid Arthritis and Renal Dysfunction in a Real-World Setting. ○
- 174 Impact of warm ischemia on acute kidney injury after robotic partial nephrectomy stratified by baseline kidney function. ○
- 173 Improvement of physical activity significantly reduced serum hepatocyte growth factor levels in a general population: 10-year prospective study. ○
- 172 Long-term effects of primary aldosteronism treatment on patients with primary aldosteronism and chronic kidney disease. ○
- 171 Metabolomic Profiling of Plasma, Urine, and Saliva of Kidney Transplantation Recipients. **2022**, 23, 13938 ○
- 170 Effects of Smoking on the Gut Microbiota in Individuals with Type 2 Diabetes Mellitus. **2022**, 14, 4800 ○
- 169 Research on the relationship between serum indoxyl sulfate concentration and iron dynamics index in patients with end-stage kidney disease: a cross-sectional study. **2022**, 8, ○

- 168 Comparison of full-dose gemcitabine/cisplatin, dose-reduced gemcitabine/cisplatin, and gemcitabine/carboplatin in real-world patients with advanced urothelial carcinoma. **2022**, 22, ○
- 167 Association between microscopic hematuria and albuminuria in patients with chronic kidney disease caused by diabetes and hypertension: the Fukuoka Kidney disease Registry Study. ○
- 166 Association of Estimated Glomerular Filtration Rate With Progression of Albuminuria in Individuals With Type 2 Diabetes. ○
- 165 Removing Race and Body Surface Area Indexation for Estimated Kidney Function Based Drug Dosing: Aminoglycosides as Justification of These Principles. ○
- 164 Association between blood urea nitrogen to creatinine ratio and neurologically Favourable outcomes in out-of-hospital cardiac arrest in adults: A multicentre cohort study. **2022**, ○
- 163 Low intake of β -carotene and dietary fiber from vegetables and fruits in patients with chronic kidney disease. **2022**, 12, ○
- 162 Utility of Indices Obtained During Medical Checkups for Predicting Fatty Liver Disease in Non-obese People. **2023**, ○
- 161 Joint effects of per- and polyfluoroalkyl substance alternatives and heavy metals on renal health: A community-based population study in China. **2023**, 219, 115057 1
- 160 Impact of simple equation for estimating appendicular skeletal muscle mass in patients with stable coronary artery disease undergoing percutaneous coronary intervention. **2023**, 44, 101163 ○
- 159 Time Trend in Incidence of Sudden Cardiac Death After Percutaneous Coronary Intervention from 2009 to 2017 (from the Japanese Multicenter Registry). **2023**, 188, 44-51 ○
- 158 Exploring biomarkers of osteosarcopenia in older adults attending a frailty clinic. **2023**, 172, 112047 ○
- 157 Comparison of the prognostic value of lateral versus septal early mitral annulus velocity in patients with acute decompensated heart failure. ○
- 156 Association between COVID-19 vaccination and relapse of glomerulonephritis. ○
- 155 Comparison of robot-assisted partial nephrectomy with soft coagulation and double-layer technique for complex and non-complex tumors. ○
- 154 Cardiovascular vs. non-cardiovascular deaths after heart failure hospitalization in young, older, and very old patients. 1
- 153 Insulin-like growth factor-1 levels are associated with high comorbidity of metabolic disorders in obese subjects; a Japanese single-center, retrospective-study. **2022**, 12, ○
- 152 Dehydration and hospital-associated disability in acute hospitalized older adults. ○
- 151 Association between Dietary Potassium Intake Estimated from Multiple 24-hour Urine Collections and Serum Potassium in Patients with Chronic Kidney Disease. **2022**, ○

150	Differential effects of overweight/obesity depending on the severity of heart failure complicating acute myocardial infarction in Japan. 2022,	0
149	Prevalence and predictive factors for clinical outcomes of isolated functional tricuspid regurgitation. 2022,	0
148	The Relationship between Serum Adiponectin, Urinary Albumin/Creatinine Ratio and Type 2 Diabetes: A Population-Based Cross-Sectional Study. 2022, 11, 7232	0
147	Effects of External Radiation Exposure Resulting From the Fukushima Daiichi Nuclear Power Plant Accident on the Health of Residents in the Evacuation Zones: the Fukushima Health Management Survey. 2022, 32, S84-S94	4
146	Association of Serum Zinc Level with severity of chronic kidney disease in diabetic patients: a cross-sectional study. 2022, 23,	0
145	Positive correlation between organic anion transporter 1B function indicated by plasma concentration of coproporphyrin-I and blood concentration of cyclosporin A in real-world patients.	0
144	Nationwide cohort study identifies clinical outcomes of angioectasia in patients with acute hematochezia.	0
143	Long-term prognostic value of changes in left ventricular global longitudinal strain in patients with heart failure with preserved ejection fraction.	0
142	Time to normalisation of CRP and incidence of relapse in microscopic polyangiitis: A medical records review study in Japan.	0
141	Association between periodontitis and chronic kidney disease by functional atherosclerosis status among older Japanese individuals: A cross-sectional study.	0
140	Renal function outcome after selective bladder-preserving tetramodality therapy consisting of maximal transurethral resection, induction chemoradiotherapy and consolidative partial cystectomy in comparison with radical cystectomy for patients with muscle-invasive bladder cancer: a two-centre retrospective study.	0
139	Natural History of Leaflet Thrombosis After Transcatheter Aortic Valve Replacement: A 5-Year Follow-Up Study. 2022, 11,	2
138	Combination of Neutrophil-to-Lymphocyte and Platelet-to-Lymphocyte Ratios as a Novel Predictor of Cardiac Death in Patients With Acute Decompensated Heart Failure With Preserved Left Ventricular Ejection Fraction: A Multicenter Study.	0
137	Impact of body mass index on in-hospital mortality in older patients hospitalized for bacterial pneumonia with non-dialysis-dependent chronic kidney disease. 2022, 22,	0
136	Renoprotective effect of chronic treatment with sodium-glucose cotransporter 2 inhibitors and its associated factors in Japanese patients with chronic heart failure and diabetes. 2022, 43, 101152	0
135	Estimated excretion and clearance of uric acid as optimal surrogate indices for daily urinary uric acid excretion.	0
134	Utility of the Age Discrepancy between Frailty-Based Biological Age and Expected Life Age in Patients with Urological Cancers. 2022, 14, 6229	1
133	Association between Advanced Glycation End-Products, Carotenoids, and Severe Erectile Dysfunction. 41,	0

- 132 Population pharmacokinetics and limited sampling strategy for therapeutic drug monitoring of mycophenolate mofetil in Japanese patients with lupus nephritis. **2023**, 9, ○
- 131 Cardiovascular Risk Factor Burden and Treatment Control in Patients with Chronic Kidney Disease: A Cross-Sectional Study. **2022**, ○
- 130 Long-Term Tailor-Made Exercise Intervention Reduces the Risk of Developing Cardiovascular Diseases and All-Cause Mortality in Patients with Diabetic Kidney Disease. **2023**, 12, 691 ○
- 129 Development and validation of a prognostic dynamic nomogram for in-hospital mortality in patients with Stanford type B aortic dissection. 9, ○
- 128 Impact of glycemic control and CYP3A5 polymorphisms on tacrolimus trough concentrations after adult kidney transplantation. ○
- 127 Hyperuricemia predicts increased cardiovascular events in patients with chronic coronary syndrome after percutaneous coronary intervention: A nationwide cohort study from Japan. 9, ○
- 126 Optimal dose of rituximab in ABO-incompatible kidney transplantation in patients with low anti-A/B antibody titers: a single-center retrospective cohort study. ○
- 125 The body mass index change is associated with death or hemodialysis transfer in Japanese patients initiating peritoneal dialysis. **2023**, 45, ○
- 124 The cutoff value of presepsin for diagnosing sepsis increases with kidney dysfunction, a cross-sectional observational study. **2023**, 102, e32620 ○
- 123 Glomerular expression and urinary excretion of fatty acid-binding protein 4 in IgA nephropathy. ○
- 122 Late-onset Chronic Kidney Disease Over 2 Decades After Pediatric Liver Transplantation: A Single-center, Retrospective Study. Publish Ahead of Print, ○
- 121 Prophylactic tenofovir alafenamide for hepatitis B virus reactivation and reactivation-related hepatitis. **2023**, 95, ○
- 120 Kidney transplantation for a transgender male. ○
- 119 Association between Familial Hypercholesterolemia and Serum Levels of Cholesterol Synthesis and Absorption Markers: The CACHE Study FH Analysis. **2023**, ○
- 118 Significance of podocyte DNA damage and glomerular DNA methylation in CKD patients with proteinuria. ○
- 117 Impact of early initiation of renin-angiotensin blockade on renal function and prognosis in patients with hypertensive emergency: A retrospective cohort study. ○
- 116 Clinical Characteristics and Prognosis of Patients with Vasospastic Angina Subjected to the Spasm Provocation Test and the Unavoidable Use of Nitroglycerin. **2023**, 10, 16 ○
- 115 Association between trunk flexibility and renal flow pulsatility in middle-aged and older adults. **2023**, 172, 112060 ○

114	Usefulness of gallium-67 scintigraphy for evaluating the histopathological activity in interstitial nephritis.	0
113	Association of polypharmacy with incidence of CKD: a retrospective cohort study.	0
112	Incidence of acute kidney injury and decreased estimated glomerular filtration rate according to the site of cancer.	0
111	Is higher lymphocyte count a potential strategy for preventing chronic kidney disease in patients receiving long-term dasatinib treatment?. 2023 , 9,	0
110	Metabolic dysfunction-associated fatty liver disease is associated with an increase in systolic blood pressure over time: linear mixed-effects model analyses.	1
109	BMI and cardiometabolic traits in Japanese: a Mendelian randomization study. 2023 ,	0
108	Handgrip strength is inversely associated with augmentation index in patients with type 2 diabetes. 2023 , 13,	0
107	Serum Values of Cholesterol Absorption and Synthesis Biomarkers in Japanese Healthy Subjects: The CACHE Study HEALTHY Analysis. 2023 ,	0
106	Hyperkalemia by eplerenone or esaxerenone in the presence or absence of clarithromycin in hypertensive patients: a retrospective observational cohort study. Publish Ahead of Print,	0
105	The association of BP with cardiovascular outcomes in patients with dipstick proteinuria and preserved kidney function.	0
104	Sex Differences in Long-Term Outcomes in Patients With Chronic Coronary Syndrome After Percutaneous Coronary Intervention [Insights From a Japanese Real-World Database Using a Storage System] 2023 ,	0
103	Circulating Extracellular Vesicle-Propagated microRNA Signature as a Vascular Calcification Factor in Chronic Kidney Disease.	0
102	Use of Complementary and Alternative Medicine and Its Relationship with Health-Related Quality of Life (HRQOL) in Patients with Type 2 Diabetes Mellitus. 2023 , 4, 53-59	1
101	Associations Among Tooth Loss, Periodontitis, and Carotid Intima-Media Thickness: the Nagahama Study. 2023 ,	0
100	Association between Bone Mineral Density and Oral Frailty on Renal Function: Findings from the Shika Study. 2023 , 11, 314	0
99	Machine-learning-based Web system for the prediction of chronic kidney disease progression and mortality. 2023 , 2, e0000188	0
98	Periodic health checkups reduce the risk of hospitalization in patients with type 2 diabetes. 4,	0
97	Pharmacokinetic and dose-finding study of osimertinib in patients with impaired renal function and low body weight.	0

- 96 Population pharmacokinetic model and dosing optimization of vancomycin in hematologic malignancies with neutropenia and augmented renal clearance. **2023**, ○
- 95 Use of Japanese big data from electronic medical records to investigate risk factors and identify their high-risk combinations for linezolid-induced thrombocytopenia. ○
- 94 Serum and plasma levels of Ba, but not those of soluble C5b-9, might be affected by renal function in chronic kidney disease patients. **2023**, 24, ○
- 93 Subclassification of pT3 upper tract urothelial carcinoma: a multicenter retrospective study. ○
- 92 Clinical performance of a new intact FGF23 immunoassay in healthy individuals and patients with chronic hypophosphatemia. **2023**, 18, 101659 ○
- 91 Risk Factors for Hyperuricemia or Gout in Men and Women: The Circulatory Risk in Communities Study (CIRCS). **2023**, ○
- 90 Phenotyping of Elderly Patients With Heart Failure Focused on Noncardiac Conditions: A Latent Class Analysis From a Multicenter Registry of Patients Hospitalized With Heart Failure. **2023**, 12, ○
- 89 Association Between Endothelial Dysfunction and Left Ventricular Diastolic Stiffness [Subanalysis of the Flow-Mediated Dilation Japan (FMD-J) Study] **2023**, ○
- 88 Effect of pregnane X receptor and cytochrome P450 oxidoreductase gene polymorphisms on trough concentrations of rivaroxaban and edoxaban in patients with nonvalvular atrial fibrillation. ○
- 87 Changes in histopathology and heteroplasmy rates over 8 years and effectiveness of taurine supplementation in a patient with mitochondrial nephropathy caused by MT-TL1 mutation: A case report. **2023**, 9, e14923 ○
- 86 Skeletal Muscle Mass Is a Strong Predictor of Cardiorespiratory Fitness in Chinese Population with Obesity. **2023**, ○
- 85 Malnutrition according to the GLIM criteria with kidney dysfunction is associated with increased mortality in hospitalized patients with cardiovascular disease: A retrospective cohort study. **2023**, 55, 167-173 ○
- 84 Focal segmental glomerulosclerosis with a mutation in the mitochondrially encoded NADH dehydrogenase 5 gene: A case report. **2023**, 35, 100963 ○
- 83 Application of urinary biomarkers for diagnosing acute kidney injury in critically ill patients without baseline renal function data. **2023**, 77, 154312 ○
- 82 Fasting plasma glucose level-based formula for estimating starting daily dose in basal-bolus insulin therapy. **2023**, 13, ○
- 81 Changes in Kidney and Liver Volumes in Patients With Autosomal Dominant Polycystic Kidney Disease Before and After Dialysis Initiation. **2023**, 7, 69-80 ○
- 80 Use of Race in Kidney Function Estimation: Lessons Learned and the Path Toward Health Justice. **2023**, 74, 385-400 ○
- 79 A nutritional assessment tool, GNRI, predicts sarcopenia and its components in type 2 diabetes mellitus: A Japanese cross-sectional study. 10, ○

78	Prognostic Implications of Urinary Potassium to Creatinine Ratio in Patients With Predialysis Chronic Kidney Disease: A Cohort Study. 2023 ,	○
77	Relationship Between Retinal Microcirculation and Renal Function in Patients with Diabetes and Chronic Kidney Disease by Laser Speckle Flowgraphy. 2023 , 13, 424	1
76	Mean HbA 1c , HbA 1c Variability, and All-cause and Cardiovascular Disease related Mortality in Older Men with Type 2 Diabetes, a Retrospective cohort study: A Focus on Early Glycemic Control.	○
75	Excess risk of cardiovascular events in patients in the United States vs. Japan with chronic kidney disease is mediated mainly by left ventricular structure and function. 2023 , 103, 949-961	○
74	Antimicrobial Stewardship Program for Patients in the Hematological Department Receiving Carbapenem Therapy: A Single-Center and Interrupted Time Series Analysis. 2023 , 12, 302	○
73	Effects of olmesartan and amlodipine on blood pressure, endothelial function, and vascular inflammation.	○
72	Association of triglycerides to high-density lipoprotein cholesterol ratio with incident cardiovascular disease but not end-stage kidney disease among patients with biopsy-proven diabetic nephropathy.	○
71	Impact of urine cyclic AMP relative to plasma arginine vasopressin on response to tolvaptan in patients with chronic kidney disease and heart failure. 2023 , 27, 427-434	○
70	Differences in clinical outcomes between octogenarian and nonagenarian patients with acute heart failure. 2023 , 23, 227-233	○
69	Association between Urinary Creatinine Excretion and Hypothyroidism in Patients with Chronic Kidney Disease. 2023 , 13, 669	○
68	Sex differences in risk factors for end-stage kidney disease and death in type 2 diabetes: A retrospective cohort study. 2023 , 15, 246-254	○
67	Decreased thymic output predicts progression of chronic kidney disease. 2023 , 20,	○
66	Association between serum uric acid levels and achievement of target blood pressure among Japanese community residents with hypertension. 2023 , 25, 295-303	○
65	Serological and histopathological assessment of galactose-deficient immunoglobulin A1 deposition in kidney allografts: A multicenter prospective observational study. 2023 , 18, e0281945	○
64	Association between the extent of house collapse and urine sodium-to-potassium ratio of victims affected by the 2011 Great East Japan Earthquake and Tsunami: a cross-sectional study.	○
63	Non-dipping pulse rate and chronic changes of the kidney in patients with chronic kidney disease. 10,	○
62	Significant Correlates of Nocturnal Hypertension in Patients With Hypertension Who Are Treated With Antihypertensive Drugs.	○
61	Association of peak C-reactive protein with long-term clinical outcomes in patients with ST-segment elevation myocardial infarction.	○

- 60 Nephrolithiasis and/or nephrocalcinosis is significantly related to renal dysfunction in patients with primary Sjögren's syndrome. ○
- 59 Referral pattern to nephrologist and prognosis in diabetic kidney disease patients: Single center retrospective cohort study. **2023**, 18, e0282163 ○
- 58 Changes in urinary albumin as a surrogate for kidney disease progression in people with type 2 diabetes. **2023**, 27, 465-472 ○
- 57 Depression and health-related quality of life in adolescents and young adults with childhood-onset end-stage kidney disease: a multicenter study in Japan. **2023**, 27, 473-479 ○
- 56 Pharmacokinetics and Safety of Imeglimin in Japanese Patients With Impaired Renal Function. ○
- 55 Anti-human leukocyte antigen and anti-ABO antibodies after SARS-CoV-2 mRNA vaccination in kidney transplant recipients. ○
- 54 Mitochondrial stress and glycooxidation increase with decreased kidney function. **2023**, 72, 147-156 ○
- 53 Significance of Circulating Remnant Lipoprotein Cholesterol Levels Measured by Homogeneous Assay in Patients with Type 2 Diabetes. **2023**, 13, 468 ○
- 52 Serum sCD40L and IL-31 in Association with Early Phase of IgA Nephropathy. **2023**, 12, 2023 ○
- 51 Compared effectiveness of sodium zirconium cyclosilicate and calcium polystyrene sulfonate on hyperkalemia in patients with chronic kidney disease. 10, ○
- 50 Evaluation of a renal risk score for Japanese patients with ANCA-associated glomerulonephritis in a multi-center cohort study. 14, ○
- 49 Combination of Muscle Quantity and Quality Is Useful to Assess the Necessity of Surveillance after a 5-Year Cancer-Free Period in Patients Who Undergo Radical Cystectomy: A Multi-Institutional Retrospective Study. **2023**, 15, 1489 ○
- 48 Correlation between acylcarnitine/free carnitine ratio and cardiopulmonary exercise test parameters in patients with incident dialysis. 14, ○
- 47 Early effects of insulin therapy on cholesterol synthesis and absorption markers in patients with type 2 diabetes. **2023**, 48, 64-74 ○
- 46 Impact of C-Reactive Protein on Long-Term Cardiac Events in Stable Coronary Artery Disease Patients with Chronic Kidney Disease. **2023**, ○
- 45 Inflammation Related to Association of Low Uric Acid and Progression to Severe Disease in Patients Hospitalized for Non-Severe Coronavirus Disease 2019. **2023**, 11, 854 ○
- 44 Real-World Efficacy of Glucagon-like Peptide-1 (GLP-1) Receptor Agonist, Dulaglutide, on Metabolic Parameters in Japanese Patients with Type 2 Diabetes: A Retrospective Longitudinal Study. **2023**, 11, 869 ○
- 43 Height loss is associated with decreased kidney function: The Japan Specific Health Checkups (J-SHC) Study. **2023**, 23, 282-288 ○

- 42 Impact of lenvatinib-induced proteinuria and renal dysfunction in patients with thyroid cancer. 13, ○
- 41 Teicoplanin and vancomycin as treatment for glycopeptide-susceptible *Enterococcus faecium* bacteraemia: a propensity score-adjusted non-inferior comparative study. ○
- 40 The level of serum albumin is associated with renal prognosis and renal function decline in patients with chronic kidney disease. **2023**, 24, ○
- 39 Salt Reduction Using a Smartphone Application Based on an Artificial Intelligence System for Dietary Assessment in Patients with Chronic Kidney Disease: A Single-Center Retrospective Cohort Study. **2023**, 3, 139-151 ○
- 38 Low agreement between various eGFR formulae in pediatric and young adult ADPKD patients. ○
- 37 Impact of cisplatin-induced acute kidney injury on long-term renal function in patients with solid tumors. ○
- 36 Association between Systemic Factors and Vitreous Fluid Cytokines in Proliferative Diabetic Retinopathy. **2023**, 12, 2354 ○
- 35 D-Asparagine is an Ideal Endogenous Molecule for Measuring the Glomerular Filtration Rate. **2023**, ○
- 34 Alcohol Consumption and a Decline in Glomerular Filtration Rate: The Japan Specific Health Checkups Study. **2023**, 15, 1540 ○
- 33 Impact of early initiation of renin-angiotensin blockade on renal function and clinical outcomes in patients with hypertensive emergency: a retrospective cohort study. **2023**, 24, ○
- 32 Growth Differentiation Factor 15 and Clinical Outcomes in Japanese Patients With Heart Failure. **2023**, ○
- 31 Hypothetical intervention of targeted systolic blood pressure control of ≤ 120 mmHg on renal prognosis for persons with stage 3-4 chronic kidney disease: an application of parametric g-formula using health checkup data in Japan. ○
- 30 The effect of dapagliflozin on uric acid excretion and serum uric acid level in advanced CKD. **2023**, 13, ○
- 29 Severe lactic acidosis with euglycemic diabetic ketoacidosis due to metformin overdose. ○
- 28 The Effect of Medical Cooperation in the CKD Patients: 10-Year Multicenter Cohort Study. **2023**, 13, 582 ○
- 27 Development of a risk prediction score and equation for chronic kidney disease: a retrospective cohort study. **2023**, 13, ○
- 26 The cisplatin-induced acute kidney injury is a novel risk factor for postoperative complications in patients with esophageal cancer: a retrospective cohort study. **2023**, 23, ○
- 25 Higher risk of intraoperative bleeding in right-sided renal tumors compared to left-sided tumors in robot-assisted partial nephrectomy. ○

- 24 Intraoperative hemodialysis during open-heart surgery in patients with severe chronic kidney disease: a retrospective cohort study. **2023**, 24, ○
- 23 Association of blood pressure and hyperuricemia with proteinuria and reduced renal function in the general population. ○
- 22 Seasonal variations in renal biopsy numbers and primary glomerular disease features based on the Japan renal biopsy registry. **2023**, 13, ○
- 21 Association between glucose tolerance and mortality among Japanese community-dwelling older adults aged over 75 years: 12-year observation of the Tosa Longitudinal Aging Study. ○
- 20 Combination of Risks of BMI and Health-Related Lifestyles on Kidney Function in the Prediabetic Japanese Population: A Prospective Cohort Study. **2023**, 20, 5338 ○
- 19 Effectiveness and current status of multidisciplinary care for patients with chronic kidney disease in Japan: a nationwide multicenter cohort study. ○
- 18 Renal cryoablation combined with prior transcatheter arterial embolization in non-dialysis patients with stage 4 or 5 chronic kidney disease: a retrospective study. ○
- 17 Sex-specific association of body mass index and fatty liver index with prevalence of renal hyperfiltration: a cross sectional study using Japanese health check-up data. **2023**, 24, ○
- 16 Independent risk factors of rapid glomerular filtration rate decline in patients with type 2 diabetes with preserved kidney function and normoalbuminuria: A multicenter cohort study. ○
- 15 Dental care and oral conditions are associated with the prevalence of sarcopenia in people with type 2 diabetes: a cross-sectional study. **2023**, 23, ○
- 14 Effect of dapagliflozin on 24-hour glycemic variables in Japanese patients with type 2 diabetes mellitus receiving basal insulin supported oral therapy (DBOT): a multicenter, randomized, open-label, parallel-group study. **2023**, 11, e003302 ○
- 13 Plasma endothelin-1 may predict bevacizumab-induced proteinuria in patients with colorectal cancer. ○
- 12 Renal Function and Clinical Outcomes Among Elderly Patients With Nonvalvular Atrial Fibrillation From ANAFIE. **2023**, ○
- 11 Changes in renal function over time in outpatients with eGFR ≥ 30 mL/min/1.73 m²: implication for timing of renal function testing before contrast-enhanced CT imaging. ○
- 10 Age and sex differences in the risk of cardiovascular diseases by chronic kidney disease in a general Japanese population. ○
- 9 Prognostic value of the 7-year protocol biopsy of adult kidney allografts: impact of mesangiosclerosis and proteinuria. **2023**, 45, ○
- 8 Heart failure with preserved ejection fraction phenogroup classification using machine learning. ○
- 7 Left Renal Vein Division During Open Surgical Repair for Abdominal Aortic Aneurysm May Cause Long-Term Kidney Remodeling. **2023**, ○

