

CITATION REPORT

List of articles citing

Consensual predictions of potential distributional areas for invasive species: a case study of Argentine ants in the Iberian Peninsula

DOI: 10.1007/s10530-008-9313-3
Biological Invasions, 2009, 11, 1017-1031.

Source: <https://exaly.com/paper-pdf/46177585/citation-report.pdf>

Version: 2024-04-28

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
133	Partitioning and mapping uncertainties in ensembles of forecasts of species turnover under climate change. 2009 , 32, 897-906		409
132	Invasive and flexible: niche shift in the drosophilid <i>Zaprionus indianus</i> (Insecta, Diptera). <i>Biological Invasions</i> , 2010 , 12, 1231-1241	2.7	56
131	Ensemble habitat mapping of invasive plant species. 2010 , 30, 224-35		127
130	Predicting potential distributions of invasive species: where to go from here?. <i>Diversity and Distributions</i> , 2010 , 16, 331-342	5	227
129	Uncertainty in ensemble forecasting of species distribution. 2010 , 16, 1145-1157		437
128	Do habitat suitability models reliably predict the recovery areas of threatened species?. 2010 , 47, 421-430		66
127	Predicting invasive alien plant distributions: how geographical bias in occurrence records influences model performance. 2010 , 37, 1797-1810		35
126	Ensemble forecasting shifts in climatically suitable areas for <i>Tropidacris cristata</i> (Orthoptera: Acridoidea: Romaleidae). 2010 , 3, 213		36
125	Defying the curse of ignorance: perspectives in insect macroecology and conservation biogeography. 2010 , 3, 172		91
124	The three phases of the ensemble forecasting of niche models: geographic range and shifts in climatically suitable areas of <i>Utetheisa ornatrix</i> (Lepidoptera, Arctiidae). 2010 , 54, 339-349		23
123	The art of modelling range-shifting species. <i>Methods in Ecology and Evolution</i> , 2010 , 1, 330-342	7.7	1435
122	Spatial Resilience in Social-Ecological Systems. 2011 ,		87
121	Geographic distribution and relative abundance of the invasive glassy-winged sharpshooter: effects of temperature and egg parasitoids. 2011 , 40, 755-69		21
120	Predicting the invasion risk by the alien bee-hawking Yellow-legged hornet <i>Vespa velutina nigrithorax</i> across Europe and other continents with niche models. 2011 , 144, 2142-2150		113
119	Tracking a medically important spider: climate change, ecological niche modeling, and the brown recluse (<i>Loxosceles reclusa</i>). <i>PLoS ONE</i> , 2011 , 6, e17731	3.7	45
118	Modeling Environments: Interactive Causation and Adaptations to Environmental Conditions*. 2011 , 78, 448-471		15
117	Predicting the Potential Distribution and Conservation Needs of Travancore Flying Squirrel, <i>Petinomys fuscocapillus</i> , in Peninsular India and Sri Lanka, using GARP. 2011 , 4, 172-186		8

116	Habitat invasion risk assessment based on Landsat 5 data, exemplified by the shrub <i>Rosa rubiginosa</i> in southern Argentina. 2011 , 36, 870-880		15
115	Assessing the environmental requirements of invaders using ensembles of distribution models. <i>Diversity and Distributions</i> , 2011 , 17, 13-24	5	73
114	Predicting potential distribution of the jaguar (<i>Panthera onca</i>) in Mexico: identification of priority areas for conservation. <i>Diversity and Distributions</i> , 2011 , 17, 350-361	5	88
113	Current and potential distribution of <i>Senecio madagascariensis</i> Poir. (fireweed), an invasive alien plant in Japan. 2011 , 57, 150-157		10
112	Ensemble modelling of species distribution: the effects of geographical and environmental ranges. 2011 , 34, 9-17		197
111	Null models reveal preferential sampling, spatial autocorrelation and overfitting in habitat suitability modelling. 2011 , 222, 588-597		72
110	Modelling species distributions with penalised logistic regressions: A comparison with maximum entropy models. 2011 , 222, 2037-2041		46
109	A SVM-based model for predicting distribution of the invasive tree <i>Miconia calvescens</i> in tropical rainforests. 2011 , 222, 2631-2641		26
108	Assessing the invasive potential of the Mediterranean fruit fly in California and Italy. <i>Biological Invasions</i> , 2011 , 13, 2661-2676	2.7	30
107	The limits to prediction in ecological systems. 2011 , 2, art125		66
106	Ensemble-based analysis of regional climate change effects on the cabbage stem weevil (<i>Ceutorhynchus pallidactylus</i> (Mrsh.)) in winter oilseed rape (<i>Brassica napus</i> L.). 2012 , 150, 191-202		22
105	Ecological niche and species traits: key drivers of regional plant invader assemblages. <i>Biological Invasions</i> , 2012 , 14, 1963-1980	2.7	30
104	Spatial Factors and Management Associated with Livestock Predations by <i>Puma concolor</i> in Central Mexico. 2012 , 40, 631-638		29
103	Can species distribution modelling provide estimates of population densities? A case study with jaguars in the Neotropics. <i>Diversity and Distributions</i> , 2012 , 18, 615-627	5	91
102	Exploring consensus in 21st century projections of climatically suitable areas for African vertebrates. 2012 , 18, 1253-1269		121
101	Does adding multi-scale climatic variability improve our capacity to explain niche transferability in invasive species?. 2012 , 246, 60-67		9
100	Spatial Variation in Postfire Cheatgrass: Dinosaur National Monument, USA. 2012 , 8, 38-56		12
99	Predicting the impact of climate change on the invasive decapods of the Iberian inland waters: an assessment of reliability. <i>Biological Invasions</i> , 2012 , 14, 1737-1751	2.7	27

98	Vulnerability of baobab species to climate change and effectiveness of the protected area network in Madagascar: Towards new conservation priorities. 2013 , 166, 11-22		37
97	Global warming may freeze the invasion of big-headed ants. <i>Biological Invasions</i> , 2013 , 15, 1561-1572	2.7	16
96	Dispersal and species responses to climate change. 2013 , 122, 1532-1540		248
95	Climate change increases the risk of invasion by the Yellow-legged hornet. 2013 , 157, 4-10		70
94	The impact of climate change changes over time. 2013 , 167, 107-115		4
93	Predicting the geographical distribution of <i>Glycaspis brimblecombei</i> (Hemiptera: Psylloidea) in Brazil. 2013 , 52, 20-30		26
92	Probabilistic accounting of uncertainty in forecasts of species distributions under climate change. 2013 , 19, 3343-54		58
91	A straightforward conceptual approach for evaluating spatial conservation priorities under climate change. 2013 , 22, 483-495		53
90	Spatial model of livestock predation by jaguar and puma in Mexico: Conservation planning. 2013 , 159, 80-87		61
89	How does selection of climate variables affect predictions of species distributions? A case study of three new weeds in New Zealand. 2013 , 53, 259-268		24
88	A strict maximum likelihood explanation of MaxEnt, and some implications for distribution modelling. 2013 , 36, 1-132		49
87	Predicting habitat distribution of five heteropteran pest species in Iran. 2013 , 13, 116		15
86	Potential Distribution and Risk Assessment of an Invasive Plant Species: A Case Study of <i>Hymenachne amplexicaulis</i> in Australia. 2013 , 19, 53-79		5
85	Increase in quantity and quality of suitable areas for invasive species as climate changes. 2013 , 27, 1458-67		25
84	Climate change may boost the invasion of the Asian needle ant. <i>PLoS ONE</i> , 2013 , 8, e75438	3.7	28
83	Identification of priority conservation areas and potential corridors for jaguars in the Caatinga biome, Brazil. <i>PLoS ONE</i> , 2014 , 9, e92950	3.7	31
82	Identifying areas of deforestation risk for REDD+ using a species modeling tool. 2014 , 9, 10		17
81	Cross-Scale Assessment of Potential Habitat Shifts in a Rapidly Changing Climate. 2014 , 7, 491-502		5

80	Comparison of the exotic and native ant communities (Hymenoptera: Formicidae) in urban green areas at inland, coastal and insular sites in Spain. 2014 , 111, 421-428		5
79	Future ant invasions in France. 2014 , 41, 217-228		10
78	Spatial distribution and range expansion of the Tawny Coster butterfly, <i>Acraea terpsicore</i> (Linnaeus, 1758) (Lepidoptera: Nymphalidae), in South-East Asia and Australia. 2014 , 7, 132-143		13
77	Incorporating spatial constraints in different periods of the annual cycle improves species distribution model performance for a highly mobile bird species. <i>Diversity and Distributions</i> , 2014 , 20, 515-528	5	17
76	Landscape diversity slows the spread of an invasive forest pest species. 2014 , 37, 648-658		29
75	Modelling the meteorological forest fire niche in heterogeneous pyrologic conditions. <i>PLoS ONE</i> , 2015 , 10, e0116875	3.7	27
74	Adaptive invasive species distribution models: a framework for modeling incipient invasions. <i>Biological Invasions</i> , 2015 , 17, 2831-2850	2.7	44
73	Evidence of niche shift and global invasion potential of the Tawny Crazy ant, <i>Nylanderia fulva</i> . 2015 , 5, 4628-41		43
72	Caveats for correlative species distribution modeling. 2015 , 29, 6-15		146
71	Different behavioural strategies among seven highly invasive ant species. <i>Biological Invasions</i> , 2015 , 17, 2491-2503	2.7	23
70	Contrasting topoclimate, long-term macroclimatic averages, and habitat variables for modelling ant biodiversity at landscape scales. 2015 , 8, 43-53		3
69	Using pest monitoring data to inform the location and intensity of invasive-species control in New Zealand. 2015 , 191, 640-649		6
68	Potential distribution of invasive alien species in the upper Ili river basin: determination and mechanism of bioclimatic variables under climate change. 2015 , 73, 779-786		7
67	Uncertainty in ensemble modelling of large-scale species distribution: Effects from species characteristics and model techniques. 2015 , 306, 67-75		29
66	Using Global and Regional Species Distribution Models (SDM) to Infer the Invasive Stage of <i>Latrodectus geometricus</i> (Araneae: Theridiidae) in the Americas. 2016 , 45, 1379-1385		16
65	A multi-scale approach to identify invasion drivers and invaders' future dynamics. <i>Biological Invasions</i> , 2016 , 18, 411-426	2.7	35
64	Potential distribution of <i>Ursus americanus</i> in Mexico and its persistence: Implications for conservation. 2016 , 29, 62-68		8
63	It's not like everybody else: urbanization factors shaping spatial distribution of native and invasive ants are species-specific. 2017 , 20, 157-169		25

62	A performance based consensus approach for predicting spatial extent of the Chinese windmill palm (<i>Trachycarpus fortunei</i>) in New Zealand under climate change. 2017 , 39, 130-139	5
61	Effects of species prevalence on the performance of predictive models. 2017 , 354, 11-19	16
60	Joint ecological, geographical and cultural approach to identify territories of opportunity for large vertebrates conservation in Mexico. 2017 , 26, 1899-1918	7
59	Potential climate change effects on tree distributions in the Korean Peninsula: Understanding model & climate uncertainties. 2017 , 353, 17-27	18
58	The potential range of <i>Ailanthus altissima</i> (tree of heaven) in South Africa: the roles of climate, land use and disturbance. <i>Biological Invasions</i> , 2017 , 19, 3675-3690	2.7 23
57	Predicting Distributions of Invasive Species. 93-129	17
56	Integrating species distribution modelling into decision-making to inform conservation actions. 2017 , 26, 251-271	38
55	Selecting predictors to maximize the transferability of species distribution models: lessons from cross-continental plant invasions. 2017 , 26, 275-287	116
54	Species Distribution Modeling: Comparison of Fixed and Mixed Effects Models Using INLA. 2017 , 6, 391	10
53	Mapping the Potential Global Range of the Brown Marmorated Stink Bug, <i>Halyomorpha halys</i> , with Particular Reference to New Zealand. 2017 , 5, 75	5
52	Development and Comparison of Species Distribution Models for Forest Inventories. 2017 , 6, 176	9
51	What are the best predictors for invasive potential of weeds? Transferability evaluations of model predictions based on diverse environmental data sets for <i>Flaveria bidentis</i> . 2018 , 58, 141-149	13
50	Environmental factors associated with the spatial distribution of invasive plant pathogens in the Iberian Peninsula: The case of <i>Phytophthora cinnamomi</i> Rands. 2018 , 419-420, 101-109	17
49	Modeling potential habitat of <i>Astragalus verus</i> Olivier for conservation decisions: A comparison of three correlative models. 2018 , 242, 61-69	8
48	Contribution of consensus methods to resolve sources of uncertainty in suitability maps modeling: application in the Zahrez El Gharbi, Steppe of Algeria. 2018 , 11, 1	
47	Landscape delineation strategy and size of mapping units impact the performance of habitat suitability models. 2018 , 47, 55-60	1
46	Decreasing brown bear () habitat due to climate change in Central Asia and the Asian Highlands. 2018 , 8, 11887-11899	22
45	How much should one sample to accurately predict the distribution of species assemblages? A virtual community approach. 2018 , 48, 125-134	16

44	Temperature and hydrologic alteration predict the spread of invasive Largemouth Bass (<i>Micropterus salmoides</i>). 2018 , 639, 58-66		23
43	Global terrestrial biomes at risk of cacti invasion identified for four species using consensual modelling. <i>Journal of Arid Environments</i> , 2018 , 156, 77-86	2.5	5
42	Potential impact of climate change on the distribution of six invasive alien plants in Nepal. 2018 , 95, 99-107		48
41	Collinearity in ecological niche modeling: Confusions and challenges. 2019 , 9, 10365-10376		76
40	In search of Aliens: Factors influencing the distribution of <i>Chromolaena odorata</i> L. and <i>Mikania micrantha</i> Kunth in the Terai grasslands of Manas National Park, India. 2019 , 131, 16-26		9
39	Multi-Scenario Species Distribution Modeling. 2019 , 10,		8
38	Molecular and morphometric identification of pistachio psyllids with niche modeling of (Hemiptera: Aphalaridae). 2020 , 110, 259-269		3
37	Integrating multiple data sources and multi-scale land-cover data to model the distribution of a declining amphibian. 2020 , 241, 108374		6
36	Quantitative methods demonstrate that environment alone is an insufficient predictor of present-day language distributions in New Guinea. <i>PLoS ONE</i> , 2020 , 15, e0239359	3.7	3
35	Current and future spatial assessment of biological control as a mechanism to reduce economic losses and carbon emissions: the case of <i>Solanum sisymbriifolium</i> in Africa. <i>Pest Management Science</i> , 2020 , 76, 2395-2405	4.6	5
34	An evaluation of central Iran's protected areas under different climate change scenarios (A Case on Markazi and Hamedan provinces). 2020 , 17, 68-82		7
33	GCM compareR: A web application to assess differences and assist in the selection of general circulation models for climate change research. <i>Methods in Ecology and Evolution</i> , 2020 , 11, 656-663	7.7	19
32	Human-Black bear interactions in Northern Mexico. <i>Human Dimensions of Wildlife</i> , 2020 , 25, 438-451	1.6	0
31	Spatial characterization and distribution modelling of <i>Ensete ventricosum</i> (wild and cultivated) in Ethiopia. <i>Geocarto International</i> , 2021 , 36, 60-75	2.7	6
30	Hotspots of livestock depredation by pumas and jaguars in Brazil: a biome-scale analysis. <i>Animal Conservation</i> , 2021 , 24, 181-193	3.2	2
29	Evaluating the invasion risk of longhorn crazy ants (<i>Paratrechina longicornis</i>) in South Korea using spatial distribution model. <i>Journal of Asia-Pacific Entomology</i> , 2021 ,	1.4	3
28	Prediction of habitat suitability for the desert monitor (<i>Varanus griseus caspius</i>) under the influence of future climate change. <i>Journal of Arid Environments</i> , 2021 , 186, 104416	2.5	1
27	Update on the invasion status of the Argentine ant, <i>Linepithema humile</i> (Mayr, 1868), in Madrid, a large city in the interior of the Iberian Peninsula. <i>Journal of Hymenoptera Research</i> , 85, 161-177	0	0

26	Habitat Suitability Modeling to Inform Seascape Connectivity Conservation and Management. <i>Diversity</i> , 2021 , 13, 465	2.5	1
25	A global assessment of the potential distribution of naturalized and planted populations of the ornamental alien tree <i>Schinus molle</i> . <i>NeoBiota</i> , 68, 105-126	4.2	
24	Modelling invasive alien plant distribution: A literature review of concepts and bibliometric analysis. <i>Environmental Modelling and Software</i> , 2021 , 145, 105203	5.2	2
23	Projecting invasion risk of non-native watersnakes (<i>Nerodia fasciata</i> and <i>Nerodia sipedon</i>) in the western United States. <i>PLoS ONE</i> , 2014 , 9, e100277	3.7	8
22	Ecological niche transferability using invasive species as a case study. <i>PLoS ONE</i> , 2015 , 10, e0119891	3.7	34
21	Niche suitability affects development: skull asymmetry increases in less suitable areas. <i>PLoS ONE</i> , 2015 , 10, e0122412	3.7	10
20	Mapping Risk of Malaria Transmission in Mainland Portugal Using a Mathematical Modelling Approach. <i>PLoS ONE</i> , 2016 , 11, e0164788	3.7	10
19	Uncertainty and sensitivity analysis in quantitative pest risk assessments; practical rules for risk assessors. <i>NeoBiota</i> , 18, 157-171	4.2	9
18	Spatial Models in Ecology and Spatial Resilience. 2011 , 87-120		
17	Are trade-offs between flexibility and efficiency in systematic conservation planning avoidable ?		
16	Coyote (<i>Canis latrans</i>) in South America: potential routes of colonization. <i>Integrative Zoology</i> , 2020 , 15, 471-481	1.9	0
15	Challenges and opportunities of species distribution modelling of terrestrial arthropod predators. <i>Diversity and Distributions</i> , 2021 , 27, 2596	5	2
14	Pet distribution modelling: Untangling the invasive potential of <i>Trachemys dorbigni</i> (Emydidae) in the Americas. <i>PLoS ONE</i> , 2021 , 16, e0259626	3.7	
13	Fine-Tuned Ecological Niche Models Unveil Climatic Suitability and Association with Vegetation Groups for Selected <i>Chaetocnema</i> Species in South Africa (Coleoptera: Chrysomelidae). <i>Diversity</i> , 2022 , 14, 100	2.5	0
12	Can We Avoid Tacit Trade-Offs between Flexibility and Efficiency in Systematic Conservation Planning? The Mediterranean Sea as a Case Study. <i>Diversity</i> , 2022 , 14, 9	2.5	
11	Prediction of the potential distribution pattern of the great gerbil (<i>Rhombomys opimus</i>) under climate change based on ensemble modelling.. <i>Pest Management Science</i> , 2022 ,	4.6	0
10	Climate change on potential suitable areas for typical Pinaceae species on the Western Sichuan Plateau. <i>Geocarto International</i> , 1-19	2.7	
9	A Scoping Review of Species Distribution Modeling Methods for Tick Vectors. <i>Frontiers in Ecology and Evolution</i> , 10,	3.7	2

8	Evaluating invasion risk and population dynamics of the brown marmorated stink bug across the contiguous United States.	2
7	Understanding the summer roosting habitat selection of the greater mouse-tailed bat (<i>Rhinopoma microphyllum</i>) and the small mouse-tailed bat (<i>Rhinopoma muscatellum</i>) in Iran.	
6	Natural forest regeneration on anthropized landscapes could overcome climate change effects on the endangered maned sloth (<i>Bradypus torquatus</i> , Illiger 1811).	1
5	Ant Diversity Is Enhanced by Ecological Infrastructures in Agroecosystems: A Case Study in Irrigated Mediterranean Farmland. 2022 , 12, 2690	0
4	Land use system, invasive species and shrub diversity of the riparian ecological infrastructure determine the specific and functional richness of ant communities in Mediterranean river valleys. 2022 , 145, 109613	0
3	Spatial and climatic analyses for predicting potential distribution of an invasive ant, <i>Linepithema humile</i> (Hymenoptera: Formicidae). 2022 , 25,	0
2	Prediction of habitat suitability, connectivity, and corridors in the future to conserve roe deer (<i>Capreolus capreolus</i>) as a locally endangered species in northern Iran. 2023 , 71, 126313	0
1	Responses of the Distribution Pattern of the Suitable Habitat of <i>Juniperus tibetica</i> Komarov to Climate Change on the Qinghai-Tibet Plateau. 2023 , 14, 434	0