

CITATION REPORT

List of articles citing

In-hospital mortality and morbidity of elderly medical patients can be predicted at admission by the Modified Early Warning Score: a prospective study

DOI: 10.1111/j.1742-1241.2008.01986.x

International Journal of Clinical Practice, 2009, 63, 591-5.

Source: <https://exaly.com/paper-pdf/46359735/citation-report.pdf>

Version: 2024-04-28

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
664	Mortality in out-of-hours emergency medical admissions--more than just a weekend effect. 2010 , 40, 115-8		27
663	Ventilator-associated pneumonia caused by <i>Pseudomonas aeruginosa</i> and respiratory colonization by <i>Candida</i> spp. 2010 , 14, P87		78
662	Abstracts of the 30th International Symposium on Intensive Care and Emergency Medicine. Brussels, Belgium. March 9-12, 2010. 2010 , 14 Suppl 1, P1-602		
661	Sodium 4-phenylbutylate protects against myocardial ischemia-reperfusion injury by reducing unfolded protein response-mediated apoptosis in mice. 2010 , 14, P2		78
660	Time-dependent effects of intravenous H ₂ S during long-term, resuscitated porcine hemorrhagic shock. 2010 , 14, P3		1
659	An ovine intensive care model of septic shock. 2010 , 14, P4		1
658	Effects of temperature and H ₂ S inhalation on glucose metabolism in murine resuscitated septic shock. 2010 , 14, P5		78
657	Mitochondrial respiration and cytochrome c inhibition by sulfide in peritoneal macrophages in vitro: effects of temperature and pH. 2010 , 14, P6		2
656	Alterations of caspase 9 mRNA gene expression in survivors and nonsurvivors of severe sepsis. 2010 , 14, P7		78
655	The TLR4 antagonist CRX-526 reduces LPS-induced leukocyte activation and improves capillary perfusion of the rat intestine. 2010 , 14, P8		78
654	Effects of lipopolysaccharide on isolated muscle mitochondrial respiration are dose and time dependent. 2010 , 14, P9		78
653	Clinical severity and local inflammatory responses in animal models of sepsis. 2010 , 14, P10		78
652	Adenosine increases during human experimental endotoxemia, but does not influence the immune response and subsequent organ injury. 2010 , 14, P11		78
651	Bacterial load plays a crucial role for survival in experimental peritonitis and modulates immunoparalysis of monocytes. 2010 , 14, P12		78
650	Decreased whole blood TNF α production capacity after acute alcohol exposure and LPS stimulation ex vivo. 2010 , 14, P13		3
649	Leukocyte immunophenotyping: methodological aspects. 2010 , 14, P14		78
648	Phenotypical analysis of peripheral human T lymphocytes in early sepsis. 2010 , 14, P15		78

647	CD64, a marker of leucocyte activation kinetics after uncomplicated cardiac surgery. 2010 , 14, P16	78
646	Interferon-gamma reverses sepsis-induced immunoparalysis of monocytes in vitro. 2010 , 14, P17	78
645	Early hydrocortisone treatment counteracts circulatory derangement but not cytokine response in porcine endotoxemia. 2010 , 14, P18	78
644	CD40L is selectively expressed on platelets from thrombocytopenic septic patients. 2010 , 14, P19	1
643	Neuronal NOS-inhibition in the setting of septic cardiomyopathy. 2010 , 14, P20	78
642	Urinary albumin excretion is elevated in sepsis, but does not correlate with circulating VEGF-A levels. 2010 , 14, P21	1
641	High mobility group box protein-1 preconditions isolated rat hearts and protects human pulmonary and renal cells against hypoxic injury. 2010 , 14, P22	78
640	Intravenous immunoglobulins prevent breakdown of the blood-brain barrier in experimental sepsis. 2010 , 14, P24	1
639	Early use of immunoglobulin in septic shock. 2010 , 14, P25	1
638	Low levels of immunoglobulin G in patients with sepsis or septic shock: a signum mali ominis?. 2010 , 14, P26	2
637	Background factors in patients receiving immunoglobulin administration and changes in sepsis markers. 2010 , 14, P27	78
636	C1-esterase inhibitor (C1INH) implication in systemic inflammation in sepsis. 2010 , 14, P28	1
635	Prognostic value of B-type natriuretic peptide in critically ill patients with new onset of fever: preliminary study. 2010 , 14, P29	78
634	Microalbuminuria evaluated as a biomarker in patients with septic shock. 2010 , 14, P30	78
633	Evaluation of automated hematologic VCS parameters in severe sepsis and septic shock: a case-control study. 2010 , 14, P31	1
632	Use of CD64 for the diagnosis of sepsis: a case-control study. 2010 , 14, P32	1
631	Endotoxin assay with endotoxin scattering photometry is one of the diagnostic markers of sepsis. 2010 , 14, P33	78
630	Immature platelet fraction as predictive index of sepsis. 2010 , 14, P34	1

629	Serial estimation of cytokine markers and their correlation with mortality in elderly patients with sepsis. 2010 , 14, P35	78
628	Influence of TIMP-1/MMP-9 ratio on the severity and mortality in sepsis. 2010 , 14, P36	78
627	Cytokine response in severe sepsis: predicting and modelling the course of illness. 2010 , 14, P37	78
626	Time course of IL-6 and LBP, candidate biomarkers of sepsis in surgical critical care. 2010 , 14, P38	78
625	Prognostic value of melatonin in patients with sepsis: a comparison of survivors and nonsurvivors. 2010 , 14, P39	1
624	Serum C-reactive protein as a prognostic variable in elective surgery ICU patients: especially valuable following esophagectomy. 2010 , 14, P40	78
623	Role of procalcitonin as a diagnostic and prognostic marker of infectious diseases in the emergency department: preliminary data. 2010 , 14, P41	78
622	Evaluation of procalcitonin at James Paget Intensive Care Unit. 2010 , 14, P42	78
621	Procalcitonin facing the postoperative culture results after liver transplantation. 2010 , 14, P43	78
620	Efficiency diagnostic and advantages of procalcitonin and C-reactive protein in the early diagnosis of sepsis. 2010 , 14, P45	78
619	Biomarkers of sepsis in burn patients. 2010 , 14, P46	1
618	Age-related differences of outcomes in patients hospitalized with community-acquired pneumonia are not explained by differences in immune response. 2010 , 14, P47	78
617	A multicentre study of bacteraemia using a new commercial universal 16S rDNA PCR test. 2010 , 14, P49	78
616	Time to antibiotics in sepsis. 2010 , 14, P50	4
615	Utility of an antibiotic guideline in hospital-associated infections. 2010 , 14, P51	78
614	Assessment of antibiotic prescriptions in two French intensive care units. 2010 , 14, P52	78
613	Occurrence of alert pathogens in the clinical materials and consumption of antibiotics in the ICU, in the years 2007 and 2008. 2010 , 14, P53	78
612	Treatment cost of secondary peritonitis in Germany: a comparative study of medical cost incurred for tigecycline therapy and standard regimens. 2010 , 14, P54	78

611	Efficacy and safety of once daily dosing of colistin to critically ill patients. 2010 , 14, P56	78
610	A comparison of the effect on renal function of flucloxacillin and vancomycin antibiotic prophylaxis in cardiac surgery patients. 2010 , 14, P57	78
609	Microdialysis study of meropenem cerebral distribution in patients with acute brain injury. 2010 , 14, P58	78
608	Effects of two regimens of antibiotic prophylaxis on colonizing flora of the respiratory tract in patients with cardiovascular surgery. 2010 , 14, P59	78
607	Prevalence of infections among patients admitted to ICUs in Nordic countries and the Netherlands in comparison with Mediterranean countries: a report from the EPIC II study. 2010 , 14, P60	78
606	Risk factors for the development of carbapenem-resistant <i>Klebsiella pneumoniae</i> infections in critically ill patients. 2010 , 14, P61	78
605	Outcome of unimicrobial versus polymicrobial sepsis. 2010 , 14, P62	2
604	De-escalation practice pattern in an Indian intensive care unit. 2010 , 14, P63	1
603	Incidence, identification of multidrug-resistant pathogens and impact on the outcome after cardiac surgery. 2010 , 14, P64	2
602	Surveillance of ICU-acquired infections in Belgium: 2008 reference data. 2010 , 14, P65	78
601	Risk of acquiring drug-resistant (MDR) Gram-negative isolates with previous exposure to antibiotic. 2010 , 14, P66	78
600	Ozone fumigation successfully controlled and eradicated multidrug-resistant <i>Acinetobacter baumannii</i> from an intensive care unit. 2010 , 14, P67	0
599	Viral infections in the ICU: should we search for them?. 2010 , 14, P68	78
598	Oseltamivir dosing with haemofiltration. 2010 , 14, P69	6
597	Efficacy and utility of a protocol for pre-emptive antimycotic therapy. 2010 , 14, P70	2
596	Amphotericin B used as a continuous infusion is safer. 2010 , 14, P71	78
595	Anidulafungin compared with fluconazole therapy in critically ill patients. 2010 , 14, P72	5
594	Increase in systolic blood pressure and improvement in laboratory parameters following polymyxin B-immobilized fiber treatment in septic shock. 2010 , 14, P73	78

593	Selection of acute blood purification therapy according to lipid mediator adsorption and blood purification in patients with septic shock. 2010 , 14, P74	78
592	Prognostic assessment in community-acquired pneumonia by pneumonia severity scores and biomarkers. 2010 , 14, P75	78
591	Current progress in pneumonia research. 2010 , 14, P76	78
590	Bacterial colonization and infections of the lower respiratory tract in an interdisciplinary ICU. 2010 , 14, P77	78
589	Rehydration does not affect pulmonary immune responses to influenza or susceptibility to secondary bacterial pneumonia. 2010 , 14, P78	78
588	Bacteremic nosocomial pneumonia cases from the ATTAIN studies. 2010 , 14, P79	78
587	Late ventilator-associated pneumonia: analysis of baseline characteristics and clinical outcomes in the ATTAIN studies. 2010 , 14, P80	1
586	On the diagnosis of acute respiratory distress syndrome in nosocomial pneumonias. 2010 , 14, P81	78
585	Genetic variability at the surfactant proteins A and D in susceptibility and severity of pneumonia. 2010 , 14, P82	1
584	Successful strategy to reduce ventilator-associated pneumonia. 2010 , 14, P83	78
583	Ventilator-associated pneumonia rate and ventilator bundle compliance in a district general hospital. 2010 , 14, P84	78
582	Oral chlorhexidine to prevent nosocomial pneumonia in critically ill patients: a systematic review and meta-analysis. 2010 , 14, P85	78
581	Impact of routine multiple site decontamination in intubated patients on ICU-acquired infections. 2010 , 14, P86	78
580	Time course of RT-PCR positivity in H1N1-induced ARDS. 2010 , 14, P88	78
579	Novel swine influenza A: most frequent ultrasonographic lung patterns. 2010 , 14, P89	78
578	Outcome and prognostic factors in patients with HIV-negative pneumocystis pneumonia requiring mechanical ventilation. 2010 , 14, P90	78
577	Acute respiratory failure from pandemic influenza A (H1N1) in an intensive care unit in southern Brazil. 2010 , 14, P91	4
576	Extracorporeal membrane oxygenation for influenza A (H1N1): experience in a regional referral center. 2010 , 14, P92	78

575	Budd-Chiari syndrome complicated by abdominal compartment syndrome: evidence of central hypovolaemia?. 2010 , 14, P93	78
574	Adenosine signalling has a protective role in intestinal ischemia and reperfusion injury. 2010 , 14, P94	78
573	Threshold levels of extreme body surface area that may cause a misrepresentation of cardiac performance. 2010 , 14, P96	78
572	Cardiac output measurement is feasible in the presence of left-to-right shunt with ultrasound dilution method: a validation study in lambs. 2010 , 14, P97	2
571	The PAPIKAS trial: a comparative clinical trial of pulmonary catheter versus the PiCCO device during therapy of patients with acute heart failure and cardiogenic shock. 2010 , 14, P98	1
570	Bioreactance versus PICCOTD/PC in critically ill septic shock patients. 2010 , 14, P99	1
569	NICOM vs LiDCOplus during changes in cardiac output in critically ill patients. 2010 , 14, P100	78
568	PulseCO consistency: variation in calibration factor over 24 and 48 hours. 2010 , 14, P102	5
567	Comparison of cardiac output values; pulse contour methods against thermodilution technique. 2010 , 14, P105	78
566	Continuous cardiac output measurement: effect of time. 2010 , 14, P106	78
565	A comparison in cardiac output data: a random effects model for repeated measures. 2010 , 14, P107	78
564	In vivo validation of a new transpulmonary thermodilution method to assess global end-diastolic volume and extravascular lung water. 2010 , 14, P108	78
563	Cardiac index derived from arterial pressure waveform: uncalibrated analysis vs once-only calibrated analysis. 2010 , 14, P109	78
562	Cardiac output monitoring during abdominal aortic cross clamping: a comparison between Vigileo/FloTrac system and transoesophageal Doppler. 2010 , 14, P110	78
561	Stroke volume index assessment using two minimal invasive devices during hemodynamic postoperative optimization. 2010 , 14, P111	78
560	Non-invasive cardiac output monitoring in children: clinical validation. 2010 , 14, P112	2
559	Global end-diastolic volume and its correlation to cardiac index inside and outside normal values. 2010 , 14, P114	78
558	Relationship of stroke volume variation, pulse pressure variation and global end-diastolic volume in patients undergoing brain surgery. 2010 , 14, P115	78

557	The value of pulse pressure variation to predict volume response in patients ventilated with low VT. 2010 , 14, P116	1
556	Pre-ejection period is a reliable parameter to estimate cardiac preload in a hemorrhagic shock. 2010 , 14, P117	78
555	Predictive value of pulse pressure variation for fluid responsiveness after maneuver to change tidal volume in patients with protective ventilatory strategy. 2010 , 14, P118	1
554	Does stroke volume increase after a fluid challenge? A study on the management of patients undergoing major head and neck free flap surgery: preliminary data. 2010 , 14, P119	1
553	Optimizing stroke volume and oxygen delivery in elective abdominal aortic surgery. 2010 , 14, P120	78
552	Oxygen saturation determined from the mouth is an early indicator of central hypovolemia in humans. 2010 , 14, P121	78
551	Plethysmographic indices as predictors of fluid responsiveness: a meta-analysis. 2010 , 14, P122	1
550	Maintenance of cardiac index within normal range is associated with mortality reduction in patients undergoing major urological surgery. 2010 , 14, P123	2
549	Effect of a preload challenge on peripheral perfusion in critically ill patients. 2010 , 14, P124	78
548	Distensibility index of inferior vena cava diameter in ventilated septic and trauma patients with shock. 2010 , 14, P125	1
547	Aortic dP/dtmax accurately reflects left ventricular contractility when effective preload independence is achieved. 2010 , 14, P126	78
546	Cardiac cycle efficiency: a new index for cardiac work estimation tested during aortic valve plasty. 2010 , 14, P127	1
545	Nonocclusive mesenteric ischemia following cardiothoracic surgery. 2010 , 14, P128	78
544	Etiology, comorbidity and prognosis of hospitalized patients with congestive heart failure. 2010 , 14, P129	1
543	Factors associated with intensive care morbidity following surgical repair of tetralogy of fallot. 2010 , 14, P130	78
542	Sildenafil attenuates pulmonary vascular remodeling and upregulates Kv1.5 mRNA expression in pulmonary hypertension secondary to left-to-right shunt in rats. 2010 , 14, P131	78
541	Validation of echocardiographic indices of right ventricular afterload: an experimental, open pericardium pig model. 2010 , 14, P132	78
540	Activation of apoptotic pathways in experimental acute afterload-induced right ventricular failure. 2010 , 14, P133	78

539	High incidence of severely prolonged QT interval after cardiac surgery. 2010 , 14, P134	2
538	Endothelial protein C receptor expression after cardiopulmonary bypass in adult cardiac surgical patients. 2010 , 14, P135	78
537	Urapidyl for hypertension control in severe pre-eclampsia: a comparative study with nicardipine. 2010 , 14, P136	2
536	Comparison of the effects of furosemide, captopril and lorazepam on noncomplicated hypertensive patients in the emergency department. 2010 , 14, P137	78
535	Incidence, risk factors and outcome of hypertensive crises in critically ill patients: a retrospective survey. 2010 , 14, P138	78
534	Comparative study to evaluate blood pressure levels measured by an invasive method versus a non-invasive method in patients in an intensive care unit. 2010 , 14, P139	78
533	Chronic atrial fibrillation in intensive care unit treated with electrical cardioversion: is it appropriate?. 2010 , 14, P140	78
532	Applanation tonometry application in ICU patients with acute cardiogenic edema. 2010 , 14, P141	78
531	A better prognosis for septic patients with left ventricular dysfunction?. 2010 , 14, P142	78
530	Microvascular effects of pulsatile versus nonpulsatile perfusion during cardiopulmonary bypass. 2010 , 14, P143	78
529	Microcirculation and intravascular coagulopathy in patients with severe sepsis and septic shock. 2010 , 14, P144	78
528	Prevalence and prognostic significance of cardiac abnormalities in the ICU: an echocardiographic study. 2010 , 14, P145	78
527	BNP and NT-proBNP in patients with acute myocardial infarction complicated by cardiogenic shock: results from the IABP Shock trial. 2010 , 14, P146	6
526	Understanding vital-sign abnormalities in critical care patients. 2010 , 14, P147	78
525	Tissue microdialysis in critically ill septic patients: associations with sepsis severity and mortality. 2010 , 14, P148	1
524	Delayed increase in inflammation and thenar VO ₂ in marathon runners. 2010 , 14, P149	78
523	Value of peripheral tissue oxygen tension in animal experimental models of haemorrhagic and septic shock. 2010 , 14, P150	78
522	Incidence of a low tissue oxygen saturation in a mixed population of critically ill patients. 2010 , 14, P151	2

521	Use of near-infrared spectroscopy during a vascular occlusion test to assess the microcirculatory response during fluid responsiveness. 2010, 14, P152	1
520	Influence of mechanical ventilation on intestinal microcirculation and inflammation in septic rats. 2010, 14, P153	78
519	ScvO ₂ and Pcv-aCO ₂ as complementary tools for goal-directed therapy during high-risk surgery. 2010, 14, P154	3
518	Prognostic value of central venous-arterial pCO ₂ difference in severe sepsis and septic shock patients. 2010, 14, P155	78
517	Veno-arterial carbon dioxide gradient at the early stage of septic shock. 2010, 14, P156	2
516	Venous-arterial PCO ₂ difference as an early predictor of organ dysfunction in the ICU. 2010, 14, P157	78
515	Lactate and ScvO ₂ combinations do not predict mortality in ICU patients. 2010, 14, P158	78
514	Early changes in lactate and central venous saturation combinations do not predict subsequent changes in organ performance. 2010, 14, P159	78
513	Regional lactate and oxygen saturation gradients: preliminary experimental data. 2010, 14, P160	78
512	Metformin increases platelet lactate production by inhibiting mitochondrial function. 2010, 14, P162	1
511	Artifactual lactic acidosis from point of care analysis: mind the gap. 2010, 14, P163	78
510	Physicochemical acid-base parameters and mortality in 1,070 ICU patients: a retrospective cohort study. 2010, 14, P165	78
509	Accuracy of predicting arterial blood gases following mechanical ventilator adjustment. 2010, 14, P166	78
508	Use of capnography and the availability of airway equipment on UK intensive care units. 2010, 14, P167	78
507	Combined oximetry and transcutaneous capnography is a reliable non-invasive monitoring in patients after cardiac surgery. 2010, 14, P168	1
506	Computerized lung sound monitoring to assess effectiveness of physiotherapy and secretion removal: a feasibility study. 2010, 14, P169	3
505	Analysis of sound waves recorded with the VRI pulmonary acoustic monitoring system in critically ill mechanically ventilated patients. 2010, 14, P170	78
504	Lung tissue properties obtained by optical alveolar elastometry. 2010, 14, P171	1

503	Combined monitoring of functional residual capacity and compliance may avoid hyperinflation and cardiac depression in ARDS. 2010 , 14, P172	78
502	Electrical impedance tomography at two thoracic levels provides detailed information about ventilation distribution in the cranial to caudal direction. 2010 , 14, P173	78
501	Changes in regional ventilation determined by volume and pressure-controlled ventilation as assessed by electrical impedance tomography in an experimental model of acid-induced acute lung injury. 2010 , 14,	78
500	Changes in lung tidal distribution assessed by electrical impedance tomography in a model of hemorrhagic shock with endotoxemia and resuscitation. 2010 , 14,	78
499	Estimating the intratidal compliance profile with the new gliding-SLICE method. 2010 , 14, P176	1
498	Atelectasis and overdistension are detected by heartbeat-induced disturbances in the pressure-volume loop. 2010 , 14, P177	78
497	Transpulmonary pressure and ventilation distribution measured with EIT during a PEEP trial in porcine acute lung injury. 2010 , 14, P178	1
496	A prospective pilot study of the effect of neutrophil elastase on the pulmonary vascular permeability in patients with pneumonia. 2010 , 14, P179	78
495	Evaluating the fibroproliferative response to ventilator-induced lung injury. 2010 , 14, P180	78
494	Effects of severe hemorrhage on pulmonary mechanics in ventilated pigs with ARDS. 2010 , 14, P181	78
493	Higher versus lower positive end-expiratory pressure in acute lung injury and acute respiratory distress syndrome: systematic review and individual patient data meta-analysis. 2010 , 14, P182	1
492	Effect of different recruitment maneuvers on bacterial translocation. 2010 , 14, P183	78
491	Nonlinear recruitment model with viscoelastic component fit respiratory mechanics in ARDS. 2010 , 14, P184	2
490	Lung recruitment and PEEP response in ARDS-related H1N1 virus patients. 2010 , 14, P185	78
489	Maximal recruitment strategy minimizes tidal recruitment in severe ARDS: a CT scan study. 2010 , 14, P186	78
488	Mechanical ventilation with different alveolar pressures improves alveolar recruitment: a model study. 2010 , 14, P187	78
487	Decreasing tidal volume from 6 to 4 ml/kg: feasibility and effects on repeated opening and closing. 2010 , 14, P188	78
486	Barriers to providing lung-protective ventilation to patients with ALI/ARDS. 2010 , 14, P189	1

485	Influence of aetiology of acute respiratory distress syndrome and early differences in oxygenation on outcome. 2010 , 14, P190	78
484	Acute lung injury in TBI patients with SIRS: role of vascular endothelial damage. 2010 , 14, P191	1
483	Mechanical ventilation aggravates transfusion-related acute lung injury induced by MHC class I antibodies. 2010 , 14, P192	78
482	Adaptive support ventilation may be inappropriate for patients with ALI/ARDS after recruitment: an observational study. 2010 , 14, P193	78
481	Receptor for advanced glycation end products is associated with systemic and organ-specific severity of acute lung injury and acute respiratory distress syndrome. 2010 , 14, P194	1
480	Mild hypothermia was protective in a physiological model of ventilator-induced lung injury by reducing inflammation but not by reducing the respiratory rate. 2010 , 14, P195	78
479	Effect of induced hypothermia on respiratory parameters in mechanically ventilated patients. 2010 , 14, P196	78
478	Extracorporeal life support service in a regional referral center: the Florence experience. 2010 , 14, P197	78
477	Interhospital ground transportation of severe acute respiratory distress syndrome patients on extracorporeal membrane oxygenation: Monza's experience. 2010 , 14, P198	78
476	Effects of hypertonic saline on a pig model of acute lung injury induced by hydrochloric acid instillation. 2010 , 14,	78
475	Conventional mechanical ventilation can injury intact lungs in severe trauma patients. 2010 , 14, P200	1
474	Protective effect of methylprednisolone on ventilator-induced diaphragm dysfunction is dose dependent. 2010 , 14, P201	78
473	Alveolar recruitment with non-invasive mechanical ventilation (C-PAP) in patients with nonobstructive respiratory failure. 2010 , 14, P202	78
472	N-acetylcysteine attenuates ventilator-induced diaphragm dysfunction in rats. 2010 , 14, P203	78
471	In vitro muscle contraction force measurements on isolated and entire rat diaphragms. 2010 , 14, P204	1
470	Useful of implementation of neurally adjusted ventilatory assist in critically ill patients. 2010 , 14, P205	1
469	Twenty-four-hour neurally adjusted ventilatory assist and volume cycled ventilation have similar effects on respiratory neuromuscular function in a porcine model of fecal peritonitis. 2010 , 14, P206	78
468	Effects of conventional and closed-loop neurally adjusted ventilatory assist in intubated spinal cord injured patients. 2010 , 14, P207	78

467	A level of neurally adjusted ventilatory assist corresponds to low levels of continuous positive airway pressure and pressure support ventilation in patients preparing to be extubated from mechanical ventilation. 2010 , 14, P208	78
466	Towards partially automated ventilation: adapting decision-making according to medical preferences. 2010 , 14, P209	78
465	Postoperative pulmonary complications in adult elective surgery patients in the US: severity, outcomes and resources use. 2010 , 14, P210	10
464	Increased duration of mechanical ventilation is associated with decreased diaphragmatic force. 2010 , 14, P211	1
463	Haemodynamics and volhaemic stage in patients with respiratory disorders to thermal injury. 2010 , 14, P212	78
462	Invasive mechanical ventilation in cancer patients: prior non-invasive ventilation is a poor prognostic factor. 2010 , 14, P213	78
461	Laryngeal angioedema: do we need a new treatment algorithm in the ER?. 2010 , 14, P214	78
460	Comparison of two insertion techniques of the ProSeal laryngeal mask airway: standard versus 90° rotation in pediatric patients. 2010 , 14, P215	78
459	Endotracheal intubation in the critically ill: complication rate and use of intubation aids. 2010 , 14, P216	78
458	Intensity and adequacy of laryngoscope light in seven intensive care units. 2010 , 14, P217	78
457	Airtraq: new device in patients at increased risk for difficult tracheal intubation. 2010 , 14, P218	78
456	Evaluation of a new tracheostomy technique for morbidly obese patients on an intensive care unit. 2010 , 14, P219	2
455	Our percutaneous tracheostomy experiences using the rotation-dilatation screw method. 2010 , 14, P220	78
454	Single-step vs balloon dilatation tracheostomy: a pilot study on 20 tracheostomies. 2010 , 14, P221	78
453	Recruitment maneuver for prevention of alevolar derecruitment during percutaneous dilation tracheostomy. 2010 , 14, P222	78
452	An audit of timing of tracheostomy formation in neurosurgical patients: how long should we wait?. 2010 , 14, P223	78
451	Percutaneous dilational tracheotomy in adult burn patients. 2010 , 14, P224	2
450	Safety of percutaneous tracheostomy in trauma patients. 2010 , 14, P225	3

449	Safety of balloon dilatational tracheostomy with the Ciaglia Blue Dolphin device. 2010 , 14, P226	78
448	Influence of the cuff pressure on the swallowing reflex in tracheostomized ICU patients. 2010 , 14, P227	78
447	Microaspirations during mechanical ventilation: polyurethane versus polyvinyl cuffed endotracheal tubes. 2010 , 14, P228	78
446	Do endotracheal tubes prevent microaspiration?. 2010 , 14, P229	4
445	A cuff -leak pressure test is a simple method for assessing severe laryngeal edema in postoperative patients. 2010 , 14, P230	78
444	Code critical airway teams improves patient safety. 2010 , 14, P231	3
443	A prospective evaluation of hemodynamic indexes to predict weaning from mechanical ventilation in patients after cardiac surgery. 2010 , 14, P232	78
442	Rapid shallow breathing index predicts readiness to wean?. 2010 , 14, P233	78
441	Increasing pressure support during vibrocompression is useful during respiratory therapy?. 2010 , 14, P234	78
440	Predicting success in weaning from mechanical ventilation in patients with and without cardiac disease. 2010 , 14, P235	78
439	Implementation of a standardised weaning protocol in patients with prolonged mechanical ventilation in a post-acute care ICU. 2010 , 14, P236	0
438	Comparison of pressure control and pressure support modes for non-invasive mechanical ventilation in acute hypercapnic respiratory failure. 2010 , 14, P237	1
437	Non-invasive CPAP with face mask: comparison between different devices. 2010 , 14, P238	1
436	Effects of non-invasive mechanical ventilation on hemodynamic and gas exchanges parameters after cardiac surgery in patients with $300 >PaO_2/FiO_2 >150$. 2010 , 14, P239	78
435	Non-invasive positive pressure ventilation in burns. 2010 , 14, P240	78
434	Non-invasive mechanical ventilation in the weaning process: continuous hemodynamic monitoring. 2010 , 14, P241	78
433	Influence of severe obesity on non-invasive ventilation strategies and responses in patients with acute hypercapnic respiratory failure in ICU. 2010 , 14, P242	1
432	A simple predictive scoring system for prolonged mechanical ventilation in severe sepsis and septic shock. 2010 , 14, P243	78

431	Simplified approach to ICU severity scoring with MPM and EUROSCORE. 2010, 14, P244	78
430	Validation of a computerised system to calculate the sequential organ failure assessment score. 2010, 14, P245	3
429	Sequential Organ Failure Assessment scores and mortality. 2010, 14, P246	78
428	Bacteremia, APACHE II and need for renal replacement therapy are independent predictors of mortality in critically ill transplanted patients with cirrhosis. 2010, 14, P247	78
427	External validation of six mortality prediction systems for intensive care unit patients. 2010, 14, P248	78
426	Prognostic accuracy of severity score for prolonged ICU length of stay >72 hours in general surgical ICU: a prospective study. 2010, 14, P249	78
425	Perioperative scores to predict mortality in surgical oncologic patients: a review of 1,362 cases. 2010, 14,	78
424	Scoring patients with a suspected infection in the emergency department (ED): comparison of the ED-specific MEDS score with APACHE II and SOFA score. 2010, 14, P251	1
423	Validation of the new proposed Emergency Trauma Score (EMTRAS). 2010, 14, P252	2
422	A new severity score for community-acquired pneumonia: PARB score. 2010, 14, P253	2
421	Modified Early Warning Score and identification of patients with severe sepsis. 2010, 14, P254	78
420	Impact of a systematic MEWS introduction on preoperative and postoperative evaluation in urgent/emergency surgery. 2010, 14, P255	2
419	Clinical predictors of physiological deterioration. 2010, 14, P256	78
418	Modified Early Warning Scores: inaccurate summation or inaccurate assignment of score?. 2010, 14, P257	8
417	Analysis of Modified Early Warning System scores and intraoperative factors on the incidence of sepsis and septic shock after elective major surgery. 2010, 14, P258	2
416	Temporal variability of medical emergency team calls. 2010, 14, P259	78
415	Medical emergency team use and outcomes in an Australian private hospital. 2010, 14, P260	1
414	Continuously improving patient safety by a rapid response system. 2010, 14, P261	78

413	Role of critical care outreach to provide the intention to treat of hematological disorders: a pilot study. 2010 , 14, P262	78
412	Score to door time: a benchmarking tool for rapid response systems. 2010 , 14, P263	1
411	A comparison of injected lidocaine 2% and no analgesia for peripheral intravenous cannula insertion in the emergency department. 2010 , 14, P264	78
410	High-sensitive troponin T measurement for patients with acute chest pain: improvement of diagnostics?. 2010 , 14, P265	78
409	Correlation of lung comet score with decreased gas exchange in patients with acute pulmonary edema: a pilot study. 2010 , 14, P267	78
408	Magnitude of first medical contact in the long-term prognosis of patients treated with STEMI. 2010 , 14, P268	78
407	Ultrasound-guided hematoma block and fracture reduction: a new way to go forward. 2010 , 14, P269	2
406	Pulmonary edema in the prehospital setting: the importance of early treatment. 2010 , 14, P270	78
405	Efficacy of bedside ultrasonography in early diagnosis of patients admitted to the emergency department with dyspnea. 2010 , 14, P271	78
404	Auscultation in flight: comparison of amplified and traditional stethoscopes. 2010 , 14, P272	1
403	Performance of ventilators at simulated altitude: study of fraction of inspired oxygen. 2010 , 14, P273	78
402	Aeromedical evacuation of the French Army: survey of the past 10 years. 2010 , 14, P274	1
401	Comparison of two disaster drills' management performed by trained and not-trained students: key times evaluation. 2010 , 14, P275	78
400	Acute severe asthma: performance of ventilator at simulated altitude. 2010 , 14, P276	78
399	Validation of a new French-language triage algorithm: the ELISA scale. 2010 , 14, P277	1
398	Examining the suitability of medical admissions to the emergency short stay ward of a large UK hospital. 2010 , 14, P278	3
397	A study on reliability of a new triage emergency method among nursing students. 2010 , 14, P279	78
396	Aggression and violence against staff of ambulance services. 2010 , 14, P280	78

395	Analysis of the disease spectrum of treated patients after Wenchuan earthquake. 2010 , 14, P281	78
394	Large-scale implementation of a critical care surge capacity management program. 2010 , 14, P282	78
393	Standard X-rays for the victims of severe trauma: time for a change. 2010 , 14, P283	78
392	Focused assessment with sonography for trauma in the prehospital setting. 2010 , 14, P284	0
391	Clinicopathogenetic aspects of systemic inflammatory response development in polytrauma. 2010 , 14, P285	78
390	Diagnostic and prognostic significance of serum apoptosis markers in the patients with severe trauma injury. 2010 , 14, P286	78
389	Predictors of mechanical ventilation after burn injury. 2010 , 14, P287	78
388	Impact of regional imaging guidelines for head injury and unconscious polytrauma patients. 2010 , 14, P288	78
387	Blunt vascular injuries of the carotid and vertebral arteries: should we screen the asymptomatic trauma patients at high risk? Preliminary results of a prospective cohort study. 2010 , 14, P289	78
386	Effect of 8.4% sodium bicarbonate on raised intracranial pressure after traumatic brain injury. 2010 , 14, P290	78
385	Hypernatremia in pediatric patients with severe traumatic brain injury. 2010 , 14, P291	78
384	Use of transcranial Doppler in patients with severe traumatic brain injury. 2010 , 14, P292	2
383	Effect of hyperbaric oxygen on cerebral oxygen saturation in patients with sub-acute traumatic brain injury. 2010 , 14, P294	78
382	Brain biomarkers and brain tissue oxygenation: changes and correlations following severe head injury. 2010 , 14, P295	78
381	Cortex and deep grey matter have different sensitivities to hypoxia-hypotension and traumatic brain injury: a combined PtIO ₂ and microdialysis study in rats. 2010 , 14, P296	78
380	Diagnosis of carotid and vertebral artery injury in major trauma with head injury. 2010 , 14, P297	78
379	Decompressive craniectomy in severe intracranial hypertension after brain injury: early or late?. 2010 , 14, P298	78
378	Optic nerve ultrasound for detecting intracranial hypertension. 2010 , 14, P300	1

377	Measurement of neuronal specific enolase as a predictor of outcome in patients suffering from severe traumatic brain injury. 2010 , 14, P301	78
376	Hyperphosphorylated neurofilament NF-H, S100 β and NSE as indicators of severity of brain injury in children. 2010 , 14, P302	78
375	A simple hospital discharge score predicts Glasgow Outcome Scale at 12 months in patients with traumatic brain injury. 2010 , 14, P303	78
374	Outcome of aggressive treatment for blunt and penetrating traumatic cardiac arrest. 2010 , 14, P304	78
373	Using flow-rate limiting bag-valve-mask device without training. 2010 , 14, P305	78
372	Out-of-hospital cardiac arrest and resource utilisation in the ICU. 2010 , 14, P306	78
371	Comparison of external cardiac compression on the floor and in a dental chair. 2010 , 14, P307	78
370	Time factors rather than type and/or nature of CPR modify the prognosis of out-of-hospital cardiac arrests witnessed by bystanders. 2010 , 14, P309	78
369	Use of the laryngeal tube by paramedics during prehospital resuscitation: an observational study. 2010 , 14, P310	78
368	The i-gel supraglottic airway as an alternative in the preclinical emergency medical aid? A comparative study on a manikin. 2010 , 14, P311	78
367	Retention of skills during emergency airway management training: a 12-month follow-up. 2010 , 14, P312	2
366	Factors correlated with the advanced life support simulation performance of senior medical students. 2010 , 14, P313	78
365	Aging augments unwillingness to the initial three links in chains of survival. 2010 , 14, P314	78
364	Prediction of the success of cardiac resuscitation: a pattern classification approach based on ECG spectral and temporal features. 2010 , 14, P315	78
363	Dispatcher-assisted telephone cardiopulmonary resuscitation using a French compression-only protocol: performance of volunteers with or without prior life support training. 2010 , 14, P316	78
362	CPR team members' attitudes towards family presence during cardiopulmonary resuscitation. 2010 , 14, P317	78
361	Prehospital therapeutic hypothermia induced with cold infusions improves haemodynamic stability in nonshockable cardiac arrest patients. 2010 , 14, P318	1
360	Trans-nasal cooling during CPR: a single-center experience. 2010 , 14, P319	1

359	Intra-arrest cooling using a novel intra-nasal cooling method for immediate induction of therapeutic hypothermia in Germany. 2010 , 14, P320	78
358	Therapeutic hypothermia after cardiac arrest: implementation in UK intensive care units. 2010 , 14, P321	78
357	Hospital-wide therapeutic hypothermia protocol significantly reduces the time to induce hypothermia but does not improve outcomes. 2010 , 14, P322	78
356	Induced hypothermia after cardiac arrest for 12 hours: single-centre experience. 2010 , 14, P323	78
355	Survival following adult cardiac arrest in intensive care units: a 5-year retrospective analysis. 2010 , 14, P324	78
354	Effects of bed height and time on the quality of chest compressions during cardiopulmonary resuscitation. 2010 , 14, P325	78
353	New results about early therapeutic hypothermia in sudden cardiac death and following return of spontaneous circulation. 2010 , 14, P326	78
352	Infection rates among cooled patients post cardiac arrest. 2010 , 14, P327	78
351	Strategies to implement a therapeutic hypothermia program. 2010 , 14, P328	78
350	A comparison of core and tympanic temperature measurement in the critically ill. 2010 , 14, P329	1
349	Can we predict neurological prognoses with computed tomography just after resuscitation?. 2010 , 14, P330	78
348	Unintended consequences of technology improvement. 2010 , 14, P331	78
347	Bispectral Index Scale alterations in brain death patients. 2010 , 14, P332	78
346	Prognosis in patients presenting with nontraumatic coma. 2010 , 14, P333	2
345	Dynamic changes in cerebral oxygenation by two methods during cardiac surgery and postoperative cognitive decline. 2010 , 14, P334	1
344	Acetazolamide-induced cerebrovascular reactivity is impaired in sepsis-associated encephalopathy. 2010 , 14, P335	1
343	Epidemiological analysis of patients with cerebral aneurysms submitted to embolization at São Josãdo Ava Hospital. 2010 , 14, P336	78
342	Frequency/prevalence analysis of risk factors on aneurysmal subarachnoid hemorrhage. 2010 , 14, P337	1

341	Monitoring regional brain oxygen saturation in patients with clipped aneurysms of the anterior circle of Willis. 2010 , 14, P338	78
340	Early prognosis value of the first 24 hours mean plasma S100 β protein level after subarachnoid aneurysmal hemorrhage. 2010 , 14, P339	78
339	Altered aortic stiffness associated with stress cardiomyopathy in patients with aneurysmal subarachnoid haemorrhage. 2010 , 14, P340	78
338	Effect of packed red blood cell transfusion on cerebral oxygenation and metabolism after subarachnoid hemorrhage. 2010 , 14, P341	1
337	Comparison of surgical or endovascular treated aneurysmal SAH patients with a special emphasis on cerebral vasospasm. 2010 , 14, P342	78
336	Prediction of cerebral vasospasm in patients with aneurysmal subarachnoid hemorrhage using jugular bulb oximetry monitoring: preliminary results. 2010 , 14, P343	0
335	Magnesium use on prophylaxis of vasospasm morbidity and mortality rate in subarachnoid hemorrhage. 2010 , 14, P344	78
334	Effects of simvastatin in prevention of vasospasm in nontraumatic subarachnoid hemorrhage: preliminary data. 2010 , 14, P345	78
333	Addition of new criteria to Sequential Organ Failure Assessment for patients with subarachnoid hemorrhage. 2010 , 14, P346	78
332	Cardiac output augmentation with fluid resuscitation improves brain tissue oxygenation after severe brain injury. 2010 , 14, P347	78
331	Multicentre experience of using ICM+ for investigations of cerebrovascular dynamics with near-infrared spectroscopy. 2010 , 14, P348	2
330	Cerebral oxygen saturation and cerebral blood flow are relatively stable during single-lung ventilation, if normocapnia is maintained. 2010 , 14, P349	78
329	Factors affecting the transient hyperaemic response test of the cerebral autoregulation. 2010 , 14, P350	78
328	Infection associated with the external ventricular derivation catheter: preliminary data. 2010 , 14, P351	78
327	fMRI default mode connectivity in vegetative state and locked-in syndrome patients. 2010 , 14, P352	78
326	Incidence of cerebral salt wasting syndrome: comparison in subarachnoid haemorrhagic compared with head trauma. 2010 , 14, P353	78
325	Sodium dysbalances in neurointensive care: results of a 5-year prospective study. 2010 , 14, P354	78
324	Impact of hypernatremia on patients with severe traumatic brain injury. 2010 , 14, P355	5

323	Prevalence and clinical setting of SIADH in neurosurgery: comparison between subarachnoid haemorrhage and head trauma. 2010 , 14, P356	78
322	Risk factors for venous thromboembolism in Japanese patients with severe blunt trauma. 2010 , 14, P357	78
321	Safety of heparin prophylaxis for venous thromboembolism in patients with neurotrauma. 2010 , 14, P358	78
320	Prevalence and incidence of proximal deep vein thrombosis in critically ill patients. 2010 , 14, P359	78
319	Trauma patients with prophylactic IVC filter and chemical prophylaxis: a descriptive study. 2010 , 14, P360	78
318	Clinical signs and risk factors of acute deep vein thrombosis of lower extremities in patients after abdominal surgery for cancer: efficiency and safety of different types of anticoagulant therapy. 2010 , 14, P361	78
317	Multidrug sedation as a risk factor for deep vein thrombosis in the intensive care unit. 2010 , 14, P362	78
316	New method diagnostics for coagulation disorders after surgery. 2010 , 14, P363	78
315	Thrombocytopenia in surgical ICU patients. 2010 , 14, P364	78
314	Platelet count as a prognostic marker in intensive care. 2010 , 14, P365	1
313	66 Evaluation of the particle immunofiltration anti-platelet factor 4 rapid assay in MICU patients with thrombocytopenia. 2010 , 14, P366	78
312	Heparin-induced thrombocytopenia incidence in the ICU: preliminary results. 2010 , 14, P367	78
311	Treatment of the coagulation deficit during extracorporeal circulation in a porcine model by a prothrombin complex concentrate. 2010 , 14, P368	78
310	Fibrinogen in dilutional coagulopathy: a dose study in pigs. 2010 , 14, P369	78
309	Tranexamic acid reduces blood loss in postpartum haemorrhage. 2010 , 14, P370	6
308	Regional variation of the off-licence use of rFVIIa for patients with uncontrolled haemorrhage in England. 2010 , 14, P371	78
307	Early use of factor VIIa in cardiac surgery is associated with lower rates of blood transfusions without impact on outcome. 2010 , 14,	78
306	Low-frequency hemoviscoelastography in monitoring coagulation disorders after abdominal surgery for cancer. 2010 , 14, P373	78

305	Reliability of the HemoCue [®] hemoglobinometer in critically ill patients. 2010 , 14, P374	78
304	Effect of transfusion, leukocyte-depleted blood product on onset of new septic shock and mortality in septic shock. 2010 , 14, P375	78
303	Transfusion Requirements After Cardiac Surgery (TRACS) study: preliminary data of a randomized controlled clinical trial. 2010 , 14,	3
302	Analysis of red cell transfusion practices in patients without active haemorrhage over a 12-month period in a UK intensive care unit. 2010 , 14, P377	78
301	Effects of red blood cell transfusion in central venous saturation and lactate levels in patients with septic shock. 2010 , 14, P378	78
300	Massive transfusion in paediatric trauma: a single-centre experience. 2010 , 14, P379	78
299	Cytokine levels evaluation during acute isovolemic anemia. 2010 , 14,	78
298	Haemodynamic alterations in community-acquired sepsis at early stage. 2010 , 14, P381	3
297	Epidemiology of severe sepsis in India: an update. 2010 , 14, P382	18
296	Systemic inflammatory response syndrome is reduced by inline filtration in intensive care patients. 2010 , 14, P383	78
295	Review of central venous catheter-related infections in neurointensive care patients in a tertiary referral centre. 2010 , 14, P384	78
294	Clinical impact of diagnosis methods for catheter-related bloodstream infection in ICU patients. 2010 , 14, P385	78
293	Catheter-related bacteremia in a jugular site with tracheostomy and in a femoral site. 2010 , 14, P386	78
292	A comparative study between conventional and antiseptic impregnated central venous catheters in neurocritical patients. 2010 , 14, P387	78
291	Comparison between peripherally inserted central venous catheters and short-term central venous catheter in patients discharged from the ICU. 2010 , 14, P388	1
290	Vacuum-assisted closure of laparostomy, an efficient strategy for complicated intra-abdominal infection. 2010 , 14, P389	78
289	Comparison between left ventricular tissue Doppler strain imaging and the sequential organ failure assessment score (SOFA score) in patients with severe sepsis or septic shock. 2010 , 14, P390	78
288	Systolic and diastolic function and survival in severe sepsis and septic shock: clinical, echocardiographic, biochemical correlation. 2010 , 14, P391	78

287	Frequency and clinical spectrum of myocardial dysfunction in severe sepsis and septic shock. 2010 , 14, P392	78
286	Effects of nitroglycerin on intestinal microcirculatory blood flow in patients with abdominal sepsis. 2010 , 14, P393	78
285	Diastolic dysfunction in septic shock is an independent predictor of elevated B-type natriuretic peptide and hospital mortality. 2010 , 14, P394	1
284	Cardiac-specific troponin I and cardiovascular dysfunction in post-traumatic sepsis. 2010 , 14, P395	78
283	Both systolic and diastolic functions are impaired in critically ill patients with community-acquired sepsis. 2010 , 14, P396	78
282	Selective V1a agonism reduces vascular leakage and cardiopulmonary dysfunction in methicillin-resistant <i>Staphylococcus aureus</i> sepsis. 2010 , 14, P397	2
281	Sepsis resuscitation bundle compliance in septic shock: improves survival and yet better late than never. 2010 , 14, P398	78
280	Effect of myocardial preservation of early goal-directed therapy on severe sepsis/septic shock patients in the ICU. 2010 , 14, P399	1
279	Impact of the sepsis bundle strategy on the outcomes of the patients suffering from severe sepsis and septic shock. 2010 , 14, P400	78
278	Impact of Surviving Sepsis Campaign implementation in Brazil. 2010 , 14, P401	78
277	Effect of synthetic vasopressin in septic shock using esophageal Doppler. 2010 , 14, P402	78
276	Effects of sepsis resuscitation bundle in cirrhotic patients with septic shock. 2010 , 14, P403	78
275	Outcomes and incidence of bleeding events associated with drotrecogin alfa: a single-centre experience of 440 patients. 2010 , 14, P404	78
274	Activated protein C: cost matters!. 2010 , 14, P405	78
273	Outcome of patients receiving two or more infusions of activated protein C: a single-centre experience. 2010 , 14, P406	78
272	Drotrecogin alfa (activated): a single-centre experience of 110 patients. 2010 , 14, P407	78
271	Efficacy of recombinant human soluble thrombomodulin in disseminated intravascular coagulation. 2010 , 14, P408	78
270	Clinical experience with lipopolysaccharide adsorber in cancer patients with severe sepsis and septic shock. 2010 , 14, P409	78

269	Apheresis of activated leukocytes with an immobilized polymyxin B filter. 2010 , 14, P410	0
268	Therapeutic blood purification using functionalized core/shell nanomagnets. 2010 , 14, P411	78
267	Coupled plasma filtration and adsorption in septic shock: a multicentric experience. 2010 , 14, P412	2
266	Patients admitted to the ICU for extra-abdominal disease and operated on for emergency laparotomy have significant survival. 2010 , 14, P413	78
265	Can ICU admission be predicted?. 2010 , 14, P414	78
264	Unsuitable for ICU: what happens next?. 2010 , 14, P415	78
263	Admission pattern and outcome in a UK weaning centre. 2010 , 14, P416	78
262	Referral to intensive care: who and when?. 2010 , 14, P417	78
261	Acute drug overdose in an ICU: 15 years experience. 2010 , 14, P418	3
260	Prognostic factors for very short-term mortality in severe sepsis. 2010 , 14, P419	78
259	Is sepsis mortality rate driven by differences in Brazilian healthcare systems?. 2010 , 14, P420	78
258	Prognoses indicators of HIV patients admitted to the ICU. 2010 , 14, P421	78
257	Outcomes of cancer patients admitted to intensive care units with severe acute kidney injury. 2010 , 14, P422	78
256	Is intensive care beneficial for patients with haematologic malignancies?. 2010 , 14, P423	78
255	Postoperative mortality in the Netherlands: identifying high-risk surgery. 2010 , 14, P424	0
254	Outcomes of 1,000 patients with cancer admitted to the intensive care unit: a Brazilian prospective study. 2010 , 14,	1
253	Trend in outcome of critically ill patients with co-morbidity: a 15-year study. 2010 , 14, P426	78
252	Clinical profile, maternal and fetal mortality analysis of severe gestosis admitted to the ICU. 2010 , 14, P427	78

251	Obstetric critical care in developing countries: a comparison between two intensive care units. 2010 , 14, P428	78
250	Co-morbidity, demographic and social factors as predictors of outcome from intensive care: an observational cohort study. 2010 , 14, P429	78
249	C-Reactive protein and albumin as prognostic markers in intensive care. 2010 , 14, P430	1
248	Statin therapy on admission to intensive care and outcome: an observational cohort study. 2010 , 14, P431	78
247	How did you sleep in the ICU?. 2010 , 14, P432	1
246	Factors associated with longer stay on a surgical intensive care unit. 2010 , 14, P433	78
245	Social support: a significant factor for health-related quality of life after critical illness?. 2010 , 14, P434	78
244	Quality of the evaluations of clinical studies in adult critical care and emergency medicine on a specialized Internet site. 2010 , 14, P435	0
243	Functional status and quality of life in patients suffering severe cranioencephalic trauma at the time of discharge from the intensive care unit and 1 year after. 2010 , 14, P436	2
242	Long-term (>6 years) quality of life after surgical intensive care admission. 2010 , 14, P437	78
241	Children's self-reported health-related quality of life after intensive care treatment. 2010 , 14, P438	4
240	Activities of daily living (Barthel Index) at discharge from the intensive care unit. 2010 , 14, P439	4
239	A comparison of Short Form 36 and Hospital Anxiety and Depression Scale based on patients 1 month after discharge from the intensive care unit. 2010 , 14, P440	78
238	Post-traumatic stress disorder after surgical ICU admission. 2010 , 14, P441	78
237	Health-related quality of life before intensive care. 2010 , 14, P442	78
236	Nursing staff development: strategies for success in a unique critical care unit in a developing country. 2010 , 14, P443	78
235	Implementing the Leapfrog standard in a developing country. 2010 , 14, P444	78
234	Impact of thiopental use on cerebral perfusion pressure in severe brain trauma: an example of caregivers' adaptation. 2010 , 14, P445	78

233	Knowledge of ICU nurses regarding selected care areas commonly guided by protocols. 2010 , 14, P446	78
232	Pharmacist clinical interventions in the ICU. 2010 , 14, P447	5
231	Adverse drug events in the critical care unit. 2010 , 14, P448	78
230	Evaluation of appropriateness of intensive care unit admissions: the GIVITI's StART approach. 2010 , 14, P449	1
229	Noise in the critical care environment. 2010 , 14, P450	1
228	Allergy documentation and transfer within critical care. 2010 , 14, P451	78
227	Use of an anonymous medication incident reporting system on a critical care unit. 2010 , 14, P452	1
226	Three years experience of a quality improvement programme. 2010 , 14, P453	78
225	Improving team work during bedside rounds: using daily goals and best practices. 2010 , 14, P454	78
224	Stress induced by simulation-based training in anesthesiologist residents. 2010 , 14, P455	78
223	Impact of an education program in respiratory therapy quality of care. 2010 , 14, P456	78
222	Use of the FAST HUG checklist in Mexican critically ill patients. 2010 , 14, P457	78
221	Bay rooms vs single-bed rooms in intensive care unit nosocomial infections: a case-control study. 2010 , 14, P458	2
220	Impact of quality improvement process on healthcare-associated infection in the ICU in a tertiary care hospital in India. 2010 , 14, P459	78
219	Teaching airway resuscitation to novices: a randomized controlled simulator mannequin study. 2010 , 14, P460	78
218	Impact of ultrasound guidance in central venous catheterization by inexperienced trainees: a prospective observational study. 2010 , 14, P461	78
217	Performance analysis of an ICU according to GiViTI's StART approach. 2010 , 14, P462	78
216	Simulation training at the point of care to decrease the incidence of airway complications on the intensive care unit. 2010 , 14, P463	78

215	Management of needle stick injuries from incapacitated patients. 2010 , 14, P464	78
214	Stress and fatigue among anaesthesia and intensive care doctors in Lithuania. 2010 , 14, P465	78
213	Implementation of daily goals in the ICU reduces length of ICU stay and errors of omission in patient care. 2010 , 14, P466	1
212	Use of single patient rooms to decrease acquisition of antibiotic-resistant bacteria in the ICU. 2010 , 14, P467	78
211	Time to first review of new admissions to critical care by the consultant intensivist. 2010 , 14, P468	78
210	Analysis of critical care referrals. 2010 , 14, P469	2
209	Communication with intensive care patients. 2010 , 14, P470	78
208	Quality of transfers of critically ill patients within the hospital. 2010 , 14, P471	78
207	Are bariatric patients putting a strain on intensive care beds?. 2010 , 14, P472	78
206	Five-year evolution of drug prescribing in a university adult intensive care unit. 2010 , 14, P473	78
205	A semiquantitative assessment of the effects of delayed discharge from the intensive care unit of survivors of critical illness. 2010 , 14, P474	78
204	Robotics telepresence in acute care facilities during influenza AH1N1 outbreak in Mexico. 2010 , 14, P475	78
203	Association between out of hours discharge from the ICU and subsequent readmission. 2010 , 14, P476	3
202	Sequential Organ Failure Assessment in pandemic planning. 2010 , 14, P477	2
201	Failure mode and effect analysis in interhospital transport of critically ill patients: a tool to design checklists for handovers in the mobile intensive care unit. 2010 , 14, P478	78
200	Selective neurogenic blockades and perioperative immune reactivity in patients undergoing lung resection. 2010 , 14, P479	78
199	Adverse effects of beta adrenergic blockers during spinal anesthesia with 0.5% bupivacaine. 2010 , 14, P480	2
198	Remifentanyl for postoperative analgo-sedation: effects on glycaemia and HOMA score. 2010 , 14, P481	78

197	Adding ultralow-dose oral naltrexone to morphine: a double-blind, randomized clinical trial. 2010 , 14, P482	78
196	Four-day sedation using sevoflurane: fluoride and renal function. 2010 , 14, P483	78
195	Beneficial effect of sevoflurane on endotoxin-induced pulmonary hypertension. 2010 , 14, P484	1
194	Does etomidate affect prognosis in septic shock patients treated with hydrocortisone?. 2010 , 14, P485	78
193	Use of methyl-naltrexone for the treatment of opioid-induced constipation in critical care patients. 2010 , 14, P486	78
192	Safety and efficacy of a nurse-driven sedation protocol with daily sedation interruption in the ICU. 2010 , 14, P487	78
191	Effects of propofol and midazolam on indocyanine green elimination assessed with LiMON for early sepsis patients. 2010 , 14, P488	78
190	Assessment of correlation between Bispectral Index monitorization and sedation scales used in the intensive care unit. 2010 , 14, P489	0
189	Sedation and analgesia: early results from a web-based survey of UK practice. 2010 , 14, P490	4
188	Severe alcohol withdrawal syndrome in ICU: is propofol a safe option?. 2010 , 14, P491	1
187	Agitation and pain during physiotherapy in intensive care. 2010 , 14, P492	1
186	Pain evaluation in the intensive care unit: listen to the patient, a new approach!. 2010 , 14, P493	0
185	Pharmacokinetics of orally administered melatonin in critically ill patients. 2010 , 14, P494	78
184	Manifestations and clinical course of delirium tremens in the ICU. 2010 , 14, P495	1
183	Comparison of two methods for diagnosing delirium conducted in a mixed medical ICU and the concordance between two independent nurse assessors. 2010 , 14, P496	78
182	Quetiapine in prolonged ICU delirium. 2010 , 14, P497	2
181	PREDICT, Prediction of Delirium in ICU Patients: development and validation of a prediction model. 2010 , 14, P498	2
180	Hospital anxiety depression scale in our surgical ICU. 2010 , 14, P499	1

179	Fluid therapy change from synthetic colloids to only crystalloids in patients with severe sepsis. 2010 , 14, P500	78
178	Effects of balanced and unbalanced colloid and crystalloid solutions on renal microvascular perfusion in endotoxemic rats. 2010 , 14, P501	78
177	Effects of balanced and unbalanced colloid and crystalloid solutions on renal oxygenation in a rat model of hemorrhagic shock and resuscitation. 2010 , 14, P502	78
176	Effect of rapidly infused crystalloids on acid-base status of dehydrated patients in the emergency department. 2010 , 14, P503	78
175	Impact of saline-based tetrastarch vs balanced tetrastarch and crystalloid solutions on hyperchloraemic acidosis in endotoxaemic sheep. 2010 , 14, P504	78
174	Are different hydroxyethyl starch solutions an independent risk factor for acute kidney injury in surgical critically ill patients? A retrospective analysis of 3,591 patients. 2010 , 14, P505	78
173	Role of early continuous venovenous hemodiafiltration in septic shock and multiorgan failure. 2010 , 14, P506	78
172	Effect of intravenous H2S on porcine aortic occlusion-induced systemic inflammation and kidney ischemia/reperfusion injury. 2010 , 14, P507	78
171	Survival improvement in pigs with liver failure and superimposed sepsis by a new liver support system (Hepa Wash [®]). 2010 , 14, P508	1
170	Increased glomerular filtration in the critically ill patient receiving anti-infective treatment. 2010 , 14, P509	4
169	Therapeutic plasma exchange in children with acute liver failure: assessment of laboratory parameters. 2010 , 14, P510	2
168	New markers of inflammation-induced renal injury subside when endotoxin tolerance develops in humans as measured by urine proteomics. 2010 , 14, P511	78
167	Extraction and purification of 1-hydroxymidazolam glucuronide from ultrafiltrate. 2010 , 14, P512	78
166	Continuous renal replacement therapy: are we getting the dose right?. 2010 , 14, P513	78
165	Use of CRRT-CVVH in a hemato-oncological ICU in patients treated with defibrotide for veno-occlusive disease in patients after allogeneic hematopoietic stem cell transplantation: single-center experience. 2010 , 14, P514	1
164	Comparison of the efficacy and safety of two regional citrate anticoagulation protocols using acid citrate dextrose A or Prismocitrate 10/2, in patients with acute renal failure undergoing continuous venovenous haemodiafiltration. 2010 , 14, P515	2
163	Addition of prostacyclin to heparin does not prolong circuit life during continuous haemofiltration. 2010 , 14, P516	78
162	Recurrent early filter clotting in regional citrate anticoagulated continuous venovenous hemodialysis due to undetected antibodies to heparin-platelet factor 4 complexes. 2010 , 14, P517	1

161	Pilot study on a regional citrate anticoagulated continuous venovenous hemodiafiltration protocol with variable treatment dose. 2010 , 14, P518	1
160	Characteristics and outcome of patients treated with renal replacement therapy: single-center, 2-year experience. 2010 , 14, P519	78
159	Renal function in living related kidney donors above the age of 70. 2010 , 14, P520	78
158	Augmented renal clearance in traumatic brain injury. 2010 , 14, P521	2
157	Pressure-controlled hemorrhagic shock in mice: a new model of acute kidney injury. 2010 , 14, P522	78
156	Continuous urinary electrolyte measurement in a swine model of mechanical ventilation. 2010 , 14, P523	78
155	Impact of positive fluid balance on survival in critically ill cancer patients. 2010 , 14, P524	78
154	Fluid accumulation has an adverse effect on outcome of ICU patients with acute kidney injury. 2010 , 14, P525	78
153	Significant microalbuminuria, on admission to the ICU, may indicate AKI in patients with sepsis. 2010 , 14, P526	78
152	Neutrophil gelatinase-associated lipocalin as an early indicator for postoperative renal failure. 2010 , 14, P527	78
151	Neutrophil gelatinase-associated lipocalin as an early indicator for mortality in adult critically ill patients with acute kidney injury. 2010 , 14, P528	1
150	Urinary and plasma NGAL levels reflect the severity of acute kidney injury. 2010 , 14, P529	78
149	Economic impact and cost-effectiveness of urine neutrophil gelatinase-associated lipocalin following cardiac surgery. 2010 , 14, P530	2
148	Neutrophil gelatinase-associated lipocalin predicts acute kidney injury in critically ill patients. 2010 , 14, P531	78
147	Study of serum vancomycin levels in different modalities of hemodialysis in Indian patients. 2010 , 14, P532	78
146	RIFLE criteria for acute kidney dysfunction following liver transplantation: incidence and risk factors. 2010 , 14, P533	78
145	Prophylactic use of laxative for constipation in critically ill patients. 2010 , 14, P534	1
144	Study of intra-abdominal hypertension in the medical intensive care unit. 2010 , 14, P535	78

143	Intra-abdominal pressure values of emergency department intensive care unit patients and clinical outcomes. 2010 , 14, P536	78
142	Advanced haemodynamic assessment of patients with liver disease and abdominal compartment syndrome. 2010 , 14, P537	78
141	Molecular adsorbent recirculating system in hepatic failure patients with a ventricular assist device. 2010 , 14, P538	78
140	Molecular adsorbent recirculating system in patients with acute liver failure: prospective follow-up. 2010 , 14, P539	78
139	A study of patients with cirrhosis admitted to nontransplant general intensive care in the UK: prevalence, case mix, outcomes and evaluation of critical illness and disease-specific scoring systems. 2010 , 14, P540	1
138	Acute liver failure: a European perspective. 2010 , 14, P541	78
137	Intracranial pressure monitoring in acute liver failure: a review of 10 years experience. 2010 , 14, P542	8
136	Microdialysis catheters in liver transplants detect vascular complications and rejection. 2010 , 14, P543	78
135	Drug-induced hepatic cholestasis in the ICU: a case-control study. 2010 , 14, P544	78
134	Diagnosis of acute cholecystitis in septic patients. 2010 , 14, P545	78
133	Bleeding and pneumonia in intensive care unit patients given PPIs and H2RAs for prevention of stress ulcer: a meta analysis. 2010 , 14, P546	78
132	Influence of septic shock on energy expenditure in critically ill ICU patients, estimated with indirect calorimetry. 2010 , 14, P547	78
131	Evaluation of a novel motility capsule for gastric emptying in a porcine model of acute lung injury. 2010 , 14, P548	78
130	Preliminary results on the validation of a novel method for measuring intra-abdominal pressure and gastric residual volume. 2010 , 14, P549	78
129	Monitoring enteral nutrition efficacy in ICU patients: is there a place for the Brix value-derived fraction of enteral formula in gastric residual volume?. 2010 , 14, P550	0
128	Initial gastric volume, nasogastric tube and pneumonia incidence in trauma patients. 2010 , 14, P551	78
127	Achieving nutritional targets in the ICU. 2010 , 14, P552	78
126	Evaluation of studies of nutritional support team: 2 years' experience. 2010 , 14, P553	78

125	Are we feeding our critically ill patients appropriately?. 2010 , 14, P554	78
124	Effect of increased enteral nutrition on ICU-acquired infections: a multicenter observational study. 2010 , 14, P555	78
123	Impact of early enteral nutrition up to goal on the prognosis of patients with different severity of illness in the ICU. 2010 , 14, P556	78
122	Benefit of supplemental parenteral nutrition in the critically ill patient? Results of a multicenter observational study. 2010 , 14, P557	2
121	Role of fiber diet formula in surgical critically ill septic patients. 2010 , 14, P558	78
120	Effect of oral supplementation with immunonutrients on radiation-induced hematologic toxicity in cancer patients. 2010 , 14, P559	1
119	Impact of multichamber bag parenteral nutrition (OliClinomel) upon inflammatory markers (EPICOS Study). 2010 , 14, P560	78
118	Effect of n-3 fatty acids on cerebral markers and the inflammatory response in sepsis. 2010 , 14, P561	78
117	Effects of an olive oil-based lipid emulsion (ClinOleic) versus LCT/MCT-based lipid emulsions upon the inflammatory markers of critically ill patients (EPICOS Study). 2010 , 14, P562	78
116	Does a fish-oil-containing lipid emulsion improve liver function in comparison with a soybean oil lipid emulsion?. 2010 , 14, P565	78
115	Therapy of compartmentalization of the proinflammatory and anti-inflammatory cytokines in sepsis. 2010 , 14, P566	1
114	Intensive insulin therapy and brief hypoglycemia do not increase neurological injury markers in critically ill children. 2010 , 14, P567	78
113	A cluster-controlled implementation project of intensive insulin therapy: effects on blood glucose control and incidence of severe hypoglycemia. 2010 , 14, P568	78
112	Incidence of hypoglycemia in NICE sugar not so nice after consideration of the treatment period. 2010 , 14, P569	78
111	Comparison of a planned versus emergent approach implementation of tight glycemic control therapy on an ICU: an experimental design. 2010 , 14, P570	78
110	Coagulation profile in hyperglycemic septic shock patients. 2010 , 14, P571	78
109	Impact of glycemic control on survival in critical care patients with sepsis. 2010 , 14, P572	1
108	Therapeutic impact on intravenous insulin dosing with three new point-of-care glucose meters in an adult ICU. 2010 , 14, P573	78

107	Significance of blood glucose variability in the early phase for predicting survival in acutely ill patients with glucose intolerance. 2010, 14, P574	78
106	Blood glucose amplitude variation: effects of intensive insulin therapy and relative association with mortality. 2010, 14, P575	78
105	Prevention of fatal and disabling neuroglycopenia: a comparison of arterial blood sampling methods. 2010, 14, P576	78
104	Impact of real-time continuous glucose monitoring on glucose variability in critically ill patients. 2010, 14, P577	78
103	Automated blood glucose monitor is insensitive to hematocrit variation and common saccharide interferents. 2010, 14, P578	78
102	Performance of an automated blood glucose monitor in the OR/ICU. 2010, 14, P579	78
101	ICU glucose monitoring measured in plasma using mid-infrared spectroscopy. 2010, 14, P580	1
100	Validation of HbA1c as a diagnostic marker for diabetes in the critically ill. 2010, 14, P581	1
99	Effect of intravenous glucocorticoids on the efficiency of tight glycaemic control. 2010, 14, P582	78
98	Chronic critical illness: high levels of estrogens and mortality. 2010, 14, P583	78
97	Estrogen decreases inflammation and apoptotic signaling in the heart following severe burn injury. 2010, 14, P584	78
96	Dexamethasone treatment during cardiac surgery and postoperative lactate levels. 2010, 14, P585	78
95	Risk factors for hypertriglyceridemia in the intensive care unit: an exploratory study. 2010, 14, P586	78
94	Septic patients show significantly lower 1,25-dihydroxy vitamin D levels than trauma patients. 2010, 14, P587	1
93	High prevalence of severe vitamin D deficiency in intensive care patients. 2010, 14, P588	3
92	Vitamin D, parathyroid hormone and serum calcium levels and their association with hospital mortality in critically ill patients. 2010, 14, P589	5
91	Effect of restoration of 25OHD3 status in prolonged critical illness on serum innate immunity parameters LL-37 and sCD163. 2010, 14, P590	10
90	Alterations in adipose tissue during critical illness: an adaptive and protective response?. 2010, 14, P591	1

89	Impact of collection method and sample handling on measured levels of circulating ACTH. 2010 , 14, P592	1
88	Prognostic value of serum zinc levels in critically ill patients. 2010 , 14, P593	1
87	High stress during admission predicts the impact of a PICU follow-up clinic on parents' psychological outcome. 2010 , 14, P594	1
86	Patient's contact person in the intensive care unit: is it really important?. 2010 , 14, P595	78
85	Evaluating the quality of communication with patients' relatives in critical care. 2010 , 14, P596	78
84	Family satisfaction on the intensive care unit. 2010 , 14, P597	78
83	Survey into bereavement of family members of patients who died in the intensive care unit. 2010 , 14, P598	78
82	Exploration of the needs and experiences of family members of a patient being admitted to a neurosurgery intensive care unit in Shiraz, Iran. 2010 , 14, P599	78
81	Deciding the end of life for patients with disorders of consciousness: a European survey. 2010 , 14, P600	78
80	Evaluation of a new limitations and withdrawal of care framework order form in the intensive care unit. 2010 , 14, P601	78
79	Examination of the utility of serum lactate and base deficit in hemorrhagic shock. 2010 , 14, P161	78
78	Response of NICOM stroke volume to passive leg raising to predict fluid responsiveness in critically ill patients with spontaneous breathing activity. 2010 , 14, P113	2
77	Vital signs as determinants of immediate and longer term mortality outcome of patients admitted from nursing homes. 2011 , 23, 309-15	6
76	Glucose levels at admission and mortality in admitted medical patients: a not so strict association. 2011 , 22, e151	2
75	Emergency hospital admissions from care-homes: who, why and what happens? A cross-sectional study. 2011 , 57, 115-20	14
74	Is Walking Speed a Vital Sign? Absolutely!. 2012 , 28, 67-76	24
73	Examination of Patients in the Intensive Care Unit. 2012 , 645-650	
72	Epidemiology of sepsis in surgical patients. 2012 , 92, 1425-43	25

71	Organizzazione dell'ospedale per intensità di cure: gli errori da evitare. <i>Italian Journal of Medicine</i> , 2012 , 6, 1-13	0.5	3
70	Hospital organization based on intensity of care: potential errors to avoid. <i>Italian Journal of Medicine</i> , 2012 , 1-13	0.5	4
69	Simple clinical score is associated with mortality and length of stay of acute general medical admissions to an Australian hospital. 2012 , 42, 160-5		5
68	A simple scoring system for the diagnosis of palmo-plantar pigmented skin lesions by digital dermoscopy analysis. 2013 , 27, e312-9		5
67	Abnormal level of arousal as a predictor of delirium and inattention: an exploratory study. 2013 , 21, 1244-53		76
66	Association between admission delay and adverse outcome of emergency medical patients. <i>Emergency Medicine Journal</i> , 2013 , 30, 320-3	1.5	9
65	The hospital Internal Medicine specialist today: a literature review and strength, weaknesses, opportunity, threats (SWOT) analysis to develop a working proposal. <i>Italian Journal of Medicine</i> , 2013 , 7, 278	0.5	4
64	Early warning scores generated in developed healthcare settings are not sufficient at predicting early mortality in Blantyre, Malawi: a prospective cohort study. <i>PLoS ONE</i> , 2013 , 8, e59830	3.7	41
63	Medical emergency team activation: performance of conventional dichotomised criteria versus national early warning score. <i>Acta Anaesthesiologica Scandinavica</i> , 2014 , 58, 411-9	1.9	33
62	The DSM-5 criteria, level of arousal and delirium diagnosis: inclusiveness is safer. <i>BMC Medicine</i> , 2014 , 12, 141	11.4	231
61	Use of the modified early warning score and serum lactate to prevent cardiopulmonary arrest in hematology-oncology patients: a quality improvement study. <i>American Journal of Medical Quality</i> , 2014 , 29, 530-7	1.1	11
60	Development of a six-month prognostic index in patients with advanced chronic medical conditions: the PALIAR score. <i>Journal of Pain and Symptom Management</i> , 2014 , 47, 551-65	4.8	38
59	Improving recognition of patients at risk in a Portuguese general hospital: results from a preliminary study on the early warning score. <i>International Journal of Emergency Medicine</i> , 2014 , 7, 22	3.9	9
58	The impact of the use of the Early Warning Score (EWS) on patient outcomes: a systematic review. <i>Resuscitation</i> , 2014 , 85, 587-94	4	234
57	Modified early warning scoring (MEWS): evaluating the evidence for tool inclusion of sepsis screening criteria and impact on mortality and failure to rescue. <i>Journal of Clinical Nursing</i> , 2015 , 24, 3343-54	3.2	24
56	Modified Early Warning Score Changes Prior to Cardiac Arrest in General Wards. <i>PLoS ONE</i> , 2015 , 10, e0130523	3.7	23
55	An Early Warning Score Predicts Risk of Death after In-hospital Cardiopulmonary Arrest in Surgical Patients. <i>American Surgeon</i> , 2015 , 81, 916-921	0.8	6
54	Aggregate National Early Warning Score (NEWS) values are more important than high scores for a single vital signs parameter for discriminating the risk of adverse outcomes. <i>Resuscitation</i> , 2015 , 87, 75-80	4	40

53	A single-centre observational cohort study of admission National Early Warning Score (NEWS). <i>Resuscitation</i> , 2015 , 92, 89-93	4	30
52	A prognostic index for 1-year mortality can also predict in-hospital mortality of elderly medical patients. <i>Internal and Emergency Medicine</i> , 2015 , 10, 575-9	3.7	4
51	A combination of clinical parameters and blood-gas analysis identifies patients at risk of transfer to intensive care upon arrival to the Emergency Department. <i>European Journal of Emergency Medicine</i> , 2016 , 23, 305-310	2.3	2
50	Emergency department Modified Early Warning Score association with admission, admission disposition, mortality, and length of stay. <i>Journal of Community Hospital Internal Medicine Perspectives</i> , 2016 , 6, 31456	1.1	13
49	Modified Early Warning Score and VitalPac Early Warning Score in geriatric patients admitted to emergency department. <i>European Journal of Emergency Medicine</i> , 2016 , 23, 406-412	2.3	29
48	Respiratory rates measured by a standardised clinical approach, ward staff, and a wireless device. <i>Acta Anaesthesiologica Scandinavica</i> , 2016 , 60, 1444-1452	1.9	22
47	Can prehospital Modified Early Warning Score identify non-trauma patients requiring life-saving intervention in the emergency department?. <i>EMA - Emergency Medicine Australasia</i> , 2016 , 28, 84-9	1.5	10
46	Early warning score predicts acute mortality in stroke patients. <i>Acta Neurologica Scandinavica</i> , 2016 , 133, 261-7	3.8	13
45	Identifying High-Risk Children in the Emergency Department. <i>Journal of Intensive Care Medicine</i> , 2016 , 31, 660-666	3.3	8
44	Predicting mortality in acutely hospitalized older patients: a retrospective cohort study. <i>Internal and Emergency Medicine</i> , 2016 , 11, 587-94	3.7	15
43	The Effect of an Impaired Arousal on Short- and Long-Term Mortality of Elderly Patients Admitted to an Acute Geriatric Unit. <i>Journal of the American Medical Directors Association</i> , 2016 , 17, 214-9	5.9	9
42	Quick SOFA Scores Predict Mortality in Adult Emergency Department Patients With and Without Suspected Infection. <i>Annals of Emergency Medicine</i> , 2017 , 69, 475-479	2.1	83
41	Respiratory Compromise as a New Paradigm for the Care of Vulnerable Hospitalized Patients. <i>Respiratory Care</i> , 2017 , 62, 497-512	2.1	20
40	Application of the National Early Warning Score (NEWS) as a stratification tool on admission in an Italian acute medical ward: A perspective study. <i>International Journal of Clinical Practice</i> , 2017 , 71, e12934 ⁹	2.9	18
39	Strengths and limitations of early warning scores: A systematic review and narrative synthesis. <i>International Journal of Nursing Studies</i> , 2017 , 76, 106-119	5.8	115
38	Early warning systems and rapid response to the deteriorating patient in hospital: A systematic realist review. <i>Journal of Advanced Nursing</i> , 2017 , 73, 2877-2891	3.1	39
37	Development of an intelligent e-healthcare system for the domestic care industry. <i>Industrial Management and Data Systems</i> , 2017 , 117, 1426-1445	3.6	17
36	Scoping review: The use of early warning systems for the identification of in-hospital patients at risk of deterioration. <i>Australian Critical Care</i> , 2017 , 30, 211-218	2.9	25

35	Clinical study of a new Modified Early Warning System scoring system for rapidly evaluating shock in adults. <i>Journal of Critical Care</i> , 2017 , 37, 50-55	4	5
34	Barriers and facilitators related to the implementation of a physiological track and trigger system: A systematic review of the qualitative evidence. <i>International Journal for Quality in Health Care</i> , 2017 , 29, 973-980	1.9	12
33	Deaths among the elderly with ICU infections. <i>Revista Brasileira De Enfermagem</i> , 2017 , 70, 733-739	0.9	6
32	The most commonly used disease severity scores are inappropriate for risk stratification of older emergency department sepsis patients: an observational multi-centre study. <i>Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine</i> , 2017 , 25, 91	3.6	43
31	The Niguarda MEWS, a new and refined tool to determine criticality and instability in Internal Medicine Ward and Emergency Medicine Unit. <i>Italian Journal of Medicine</i> , 2017 , 11, 310	0.5	
30	Norton scale score on admission and mortality of patients hospitalized in internal medicine departments. <i>Revista Clinica Espanola</i> , 2018 , 218, 177-184	0.5	
29	Norton scale score on admission and mortality of patients hospitalised in Internal Medicine departments. <i>Revista Clinica Espanola</i> , 2018 , 218, 177-184	0.7	6
28	Hospital at Home care for older patients with cognitive impairment: a protocol for a randomised controlled feasibility trial. <i>BMJ Open</i> , 2018 , 8, e020332	3	4
27	Early prediction of hospital admission for emergency department patients: a comparison between patients younger or older than 70 years. <i>Emergency Medicine Journal</i> , 2018 , 35, 18-27	1.5	24
26	Nursing clinical competence in area medica. <i>Italian Journal of Medicine</i> , 2018 , 1-83	0.5	
25	Are Early Warning Scores Useful Predictors for Mortality and Morbidity in Hospitalised Acutely Unwell Older Patients? A Systematic Review. <i>Journal of Clinical Medicine</i> , 2018 , 7,	5.1	13
24	Using modified early warning score to predict need of lifesaving intervention in adult non-trauma patients in a tertiary state hospital. <i>Hong Kong Journal of Emergency Medicine</i> , 2018 , 25, 146-151	0.4	0
23	Characterising variation in composition and activation criteria of rapid response and cardiac arrest teams: a survey of Medicare participating hospitals in five American states. <i>BMJ Open</i> , 2019 , 9, e024548 ³		7
22	Comparison of the National Early Warning Score (NEWS) and the Modified Early Warning Score (MEWS) for predicting admission and in-hospital mortality in elderly patients in the pre-hospital setting and in the emergency department. <i>PeerJ</i> , 2019 , 7, e6947	3.1	31
21	Hospital morbidity among elderly in the region of Sfax, Tunisia: Epidemiological profile and chronological trends between 2003 and 2015. <i>Clinical Epidemiology and Global Health</i> , 2019 , 7, 131-135	1.8	1
20	Early warning scores for detecting deterioration in adult hospital patients: systematic review and critical appraisal of methodology. <i>BMJ, The</i> , 2020 , 369, m1501	5.9	64
19	Predicting cardiac arrest in the emergency department. <i>Journal of the American College of Emergency Physicians Open</i> , 2020 , 1, 321-326	1.6	2
18	Sepsis Screening: Combining Early Warning Scores and SIRS Criteria. <i>Clinical Nursing Research</i> , 2021 , 30, 42-49	1.7	5

17	Initial MEWS score to predict ICU admission or transfer of hospitalized patients with COVID-19: A retrospective study. <i>Journal of Infection</i> , 2021 , 82, 282-327	18.9	7
16	LifeSenior ̄A Health Monitoring IoT System Based on Deep Learning Architecture. <i>Lecture Notes in Computer Science</i> , 2021 , 293-306	0.9	0
15	Comparison of within 7 Day All-Cause Mortality among HDU Patients with Modified Early Warning Score of B with those with Score of . <i>Pakistan Journal of Medical Sciences</i> , 2021 , 37, 515-519	2	
14	Modified early warning score and risk of mortality after acute stroke. <i>Clinical Neurology and Neurosurgery</i> , 2021 , 202, 106547	2	0
13	Performance of universal early warning scores in different patient subgroups and clinical settings: a systematic review. <i>BMJ Open</i> , 2021 , 11, e045849	3	0
12	The effect of COVID-19 epidemic on vital signs in hospitalized patients: a pre-post heat-map study from a large teaching hospital. <i>Journal of Clinical Monitoring and Computing</i> , 2021 , 1	2	1
11	Service evaluation comparing Acute Care at Home for older people service and conventional service within an acute hospital care of elderly ward. <i>Journal of Clinical Nursing</i> , 2021 , 30, 2978-2989	3.2	1
10	The Universal Form of Treatment Options (UFTO) as an alternative to Do Not Attempt Cardiopulmonary Resuscitation (DNACPR) orders: a mixed methods evaluation of the effects on clinical practice and patient care. <i>PLoS ONE</i> , 2013 , 8, e70977	3.7	40
9	Can a Patient's In-Hospital Length of Stay and Mortality Be Explained by Early-Risk Assessments?. <i>PLoS ONE</i> , 2016 , 11, e0162976	3.7	8
8	Cascade iatrogenesis: a case-control study to detect postoperative respiratory failure in hospitalized older adults. <i>Research in Gerontological Nursing</i> , 2014 , 7, 66-77	1.6	10
7	Geriatrics-focused indicators predict mortality more than age in older adults hospitalized with COVID-19. <i>BMC Geriatrics</i> , 2021 , 21, 554	4.1	2
6	Examination of Patients in the Intensive Care Unit. 2018 , 645-652.e2		
5	Older people and urgent care. 2020 , 539-547		
4	A quick modified early warning score for triaging medical patients at admission.. <i>European Journal of Emergency Medicine</i> , 2022 , 29, 80-81	2.3	0
3	Prognostic value of National Early Warning Score and Modified Early Warning Score on intensive care unit readmission and mortality: A prospective observational study. 9,		1
2	Hospital practices and clinical outcomes associated with behavioral symptoms in persons with dementia.		0
1	A Retrospective Cohort Study Evaluating the Use of the Modified Early Warning Score to Improve Outcome Prediction in Neurosurgical Patients. 2022 ,		0