

CITATION REPORT

List of articles citing

The overlap of storage, voiding and postmicturition symptoms and implications for treatment seeking in the USA, UK and Sweden: EpiLUTS

DOI: 10.1111/j.1464-410x.2009.08369.x
BJU International, 2009, 103 Suppl 3, 12-23.

Source: <https://exaly.com/paper-pdf/46394131/citation-report.pdf>

Version: 2024-04-29

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
154	The burden of lower urinary tract symptoms: evaluating the effect of LUTS on health-related quality of life, anxiety and depression: EpiLUTS. <i>BJU International</i> , 2009 , 103 Suppl 3, 4-11	5.6	293
153	Implications of recent epidemiology studies for the clinical management of lower urinary tract symptoms. <i>BJU International</i> , 2009 , 103 Suppl 3, 48-57	5.6	23
152	Lower urinary tract symptoms: new perspectives on prevalence, burden and comorbidities. Introduction. <i>BJU International</i> , 2009 , 103 Suppl 3, 1-3	5.6	
151	Antimuscarinics in men with lower urinary tract symptoms suggestive of bladder outlet obstruction due to benign prostatic hyperplasia. 2010 , 20, 43-8		32
150	Análisis del síndrome de vejiga hiperactiva y de la incontinencia urinaria en varones en la franja de edad de los 50 a los 65 años. Estudio EPICC. <i>Actas Urológicas Españolas</i> , 2010 , 34, 543-548	0.7	2
149	Análisis del síndrome de vejiga hiperactiva y de la incontinencia urinaria en mujeres laboralmente activas entre 25 y 64 años. Estudio EPICC. <i>Actas Urológicas Españolas</i> , 2010 , 34, 618-624	0.7	4
148	Role of voiding and storage symptoms for the quality of life before and after treatment in men with voiding dysfunction. 2010 , 28, 3-8		25
147	Bladder dysfunction after acute urinary retention in the rats: a novel over active bladder model. 2010 , 333, 109-14		28
146	A placebo-controlled study investigating the efficacy and safety of the phosphodiesterase type 5 inhibitor UK-369,003 for the treatment of men with lower urinary tract symptoms associated with clinical benign prostatic hyperplasia. <i>BJU International</i> , 2010 , 106, 674-80	5.6	35
145	A placebo-controlled exploratory study investigating the efficacy and safety of the phosphodiesterase type 5 inhibitor UK-369,003 for the treatment of men with storage lower urinary tract symptoms associated with a clinical diagnosis of overactive bladder. <i>BJU International</i> , 2010 , 106, 666-73	5.6	19
144	Analysis of overactive bladder and urinary incontinence in males in the age range between 50 and 65 years. EPICC study. <i>Actas Urológicas Españolas (English Edition)</i> , 2010 , 34, 543-548	0.1	1
143	Analysis of overactive bladder and urinary incontinence in working women aged between 25 and 64 years. EPICC study. <i>Actas Urológicas Españolas (English Edition)</i> , 2010 , 34, 618-624	0.1	4
142	Efficacy and safety of tolterodine extended release and dutasteride in male overactive bladder patients with prostates >30 grams. <i>Urology</i> , 2010 , 75, 1144-8	1.6	35
141	Race and ethnic differences in health beliefs about lower urinary tract symptoms. 2011 , 60, 165-72		23
140	Self-management after prostate cancer treatment: evaluating the feasibility of providing a cognitive and behavioural programme for lower urinary tract symptoms. <i>BJU International</i> , 2011 , 107, 783-790	5.6	26
139	Prevalence of post-micturition symptoms in association with lower urinary tract symptoms and health-related quality of life in men and women. <i>BJU International</i> , 2011 , 108, 1452-8	5.6	23
138	The effects of dutasteride or tamsulosin alone and in combination on storage and voiding symptoms in men with lower urinary tract symptoms (LUTS) and benign prostatic hyperplasia (BPH): 4-year data from the Combination of Avodart and Tamsulosin (CombAT) study. <i>BJU International</i> , 2011 , 107, 1426-31	5.6	47

137	Management of lower urinary tract symptoms and bladder outlet obstruction. <i>Medicine</i> , 2011 , 39, 378-383	36	1
136	Patients' experiences of seeking health care for lower urinary tract symptoms. 2011 , 34, 496-507		15
135	Efficacy of propiverine ER with or without β -blockers related to maximum urinary flow rate in adult men with OAB: results of a 12-week, multicenter, non-interventional study. 2011 , 29, 217-23		17
134	Overactive bladder and outlet obstruction in men. <i>Current Urology Reports</i> , 2011 , 12, 77-85	2.9	15
133	Epidemiology of stress urinary incontinence in women. <i>Current Urology Reports</i> , 2011 , 12, 370-6	2.9	72
132	The prevalence of lower urinary tract symptoms and treatment-seeking behaviour in males over 40 years in Singapore: a community-based study. <i>Prostate Cancer and Prostatic Diseases</i> , 2012 , 15, 273-7	6.2	22
131	Drug development for LUTS – The challenge for industry. 2012 , 9, e5-e14		0
130	[Update of the usefulness of β -blockers in the treatment of different urologic pathologies]. 2012 , 139, 582-7		1
129	An overview of prostate diseases and their characteristics specific to Asian men. 2012 , 14, 458-64		30
128	Correlates for nocturia: a review of epidemiological studies. 2012 , 19, 317-29		51
127	A phase II dose-ranging study of mirabegron in patients with overactive bladder. <i>International Urogynecology Journal</i> , 2013 , 24, 1447-58	2	97
126	Overactive bladder in elderly men: epidemiology, evaluation, clinical effects, and management. <i>Current Urology Reports</i> , 2013 , 14, 418-25	2.9	22
125	Current State of the Art in Non-Invasive Urodynamics. 2013 , 8, 83-91		4
124	Current Understanding of the Interplay between Oab & Bph. 2013 , 8, 128-133		1
123	Medical management of lower urinary tract symptoms in men with benign prostatic enlargement. 2013 , 30, 309-19		9
122	Healthcare-seeking prevalence of lower urinary tract symptoms among national health insurance enrollees in Taiwan, 2000-2009. <i>Urology</i> , 2013 , 81, 61-5	1.6	17
121	Solifenacin plus tamsulosin combination treatment in men with lower urinary tract symptoms and bladder outlet obstruction: a randomized controlled trial. <i>European Urology</i> , 2013 , 63, 158-65	10.2	64
120	Impaired detrusor contractility in a rat model of chronic bladder ischemia. <i>Urology</i> , 2013 , 81, 1379.e9-14	1.6	22

119	Efficacy and safety of solifenacin plus tamsulosin OCAS in men with voiding and storage lower urinary tract symptoms: results from a phase 2, dose-finding study (SATURN). <i>European Urology</i> , 2013 , 64, 398-407	10.2	52
118	A proof-of-concept study: mirabegron, a new therapy for overactive bladder. <i>Neurourology and Urodynamics</i> , 2013 , 32, 1116-22	2.3	70
117	Efficacy of Additional Dutasteride on Lower Urinary Tract Symptoms in Patients with Alpha-1 Blocker-Resistant Benign Prostatic Hyperplasia. <i>LUTS: Lower Urinary Tract Symptoms</i> , 2013 , 5, 159-63	1.9	1
116	Erectile dysfunction and lower urinary tract symptoms: a consensus on the importance of co-diagnosis. <i>International Journal of Clinical Practice</i> , 2013 , 67, 606-18	2.9	48
115	Stress Urinary Incontinence in the Elderly: Evaluation, Surgical Treatment, and Management of Postoperative Voiding Dysfunction. 2014 , 9, 379-388		2
114	Improved irritative voiding symptoms 3 years after stereotactic body radiation therapy for prostate cancer. 2014 , 4, 290		12
113	Early effect of dutasteride added to alpha-1 blocker therapy for patients with lower urinary tract symptoms associated with benign prostatic hyperplasia. 2014 , 21, 815-9		1
112	Clinical implications of a feeling of incomplete emptying with little post-void residue in men with lower urinary tract symptoms. <i>Neurourology and Urodynamics</i> , 2014 , 33, 1123-7	2.3	7
111	Current medical treatment of lower urinary tract symptoms/BPH: do we have a standard?. 2014 , 24, 21-8		39
110	Antimuscarinics. 2014 , 100-112		
109	Postmenopausal overactive bladder. 2014 , 13, 313-29		6
108	Lower urinary tract symptom: still too much focus on the prostate?. 2014 , 24, 3-9		11
107	What is the most bothersome lower urinary tract symptom? Individual- and population-level perspectives for both men and women. <i>European Urology</i> , 2014 , 65, 1211-7	10.2	132
106	Systematic review and meta-analysis of candidate gene association studies of lower urinary tract symptoms in men. <i>European Urology</i> , 2014 , 66, 752-68	10.2	21
105	Future Pharmacotherapies for Male Lower Urinary Tract Symptoms. 2014 , 9, 134-141		
104	The incontinence impact questionnaire-7 (IIQ-7) can be applicable to Chinese males and females with lower urinary tract symptoms. 2014 , 7, 403-11		14
103	Management of male lower urinary tract symptoms suggestive of benign prostatic hyperplasia by general practitioners in Jakarta. 2014 , 2, 97-103		1
102	Combination therapy with tamsulosin and solifenacin for male lower urinary tract symptoms with predominant filling symptoms: a new approach to an old problem. 2015 , 31, 1963-5		

101	Efficacy and safety of muscarinic antagonists as add-on therapy for male lower urinary tract symptoms. <i>Scientific Reports</i> , 2014 , 4, 3948	4.9	6
100	Hypovitaminosis D is associated with lower urinary tract symptoms and benign prostate hyperplasia in type 2 diabetes. 2015 , 3, 1062-7		9
99	Overactive bladder. 2015 , 4,		9
98	Solifenacin/tamsulosin fixed-dose combination therapy to treat lower urinary tract symptoms in patients with benign prostatic hyperplasia. 2015 , 9, 1707-16		7
97	Evaluation of the Effectiveness of Nurse-Led Continence Care Treatments for Chinese Primary Care Patients with Lower Urinary Tract Symptoms. <i>PLoS ONE</i> , 2015 , 10, e0129875	3.7	7
96	A synopsis of drugs currently in preclinical and early clinical development for the treatment of benign prostatic hyperplasia. 2015 , 24, 1059-73		7
95	Progression and treatment of incident lower urinary tract symptoms (LUTS) among men in the California Men's Health Study. <i>BJU International</i> , 2015 , 115, 127-33	5.6	10
94	Change in storage symptoms following laser prostatectomy: comparison between photoselective vaporization of the prostate (PVP) and holmium laser enucleation of the prostate (HoLEP). 2015 , 33, 1173-80		13
93	Initial management of lower urinary tract symptoms and bladder outlet obstruction. <i>Medicine</i> , 2015 , 43, 431-434	0.6	
92	Postoperative Lower Urinary Tract Storage Symptoms: Does Prostate Enucleation Differ from Prostate Vaporization for Treatment of Symptomatic Benign Prostatic Hyperplasia?. <i>Journal of Endourology</i> , 2015 , 29, 1159-65	2.7	10
91	Urodynamic evaluation of the efficacy of mirabegron on storage and voiding functions in women with overactive bladder. <i>Urology</i> , 2015 , 85, 786-90	1.6	28
90	Open-label, 9-month extension study investigating the uro-selective alpha-blocker silodosin in men with LUTS associated with BPH. 2015 , 33, 697-706		9
89	Efficacy and safety of solifenacin plus tamsulosin oral controlled absorption system in men with lower urinary tract symptoms: a meta-analysis. 2015 , 17, 124-34		4
88	Cost-effectiveness of a fixed-dose combination of solifenacin and oral controlled adsorption system formulation of tamsulosin in men with lower urinary tract symptoms associated with benign prostatic hyperplasia. <i>BMC Urology</i> , 2015 , 15, 41	2.2	5
87	Elucidation of the Pattern of the Onset of Male Lower Urinary Tract Symptoms Using Cluster Analysis: Efficacy of Tamsulosin in Each Symptom Group. <i>Urology</i> , 2015 , 86, 349-53	1.6	5
86	Drug treatment patterns for the management of men with lower urinary tract symptoms associated with benign prostatic hyperplasia who have both storage and voiding symptoms: a study using the health improvement network UK primary care data. 2015 , 31, 43-50		18
85	Long-term safety and efficacy of single-tablet combinations of solifenacin and tamsulosin oral controlled absorption system in men with storage and voiding lower urinary tract symptoms: results from the NEPTUNE Study and NEPTUNE II open-label extension. <i>European Urology</i> , 2015 , 67, 262-70	10.2	47
84	Men's preferences for the treatment of lower urinary tract symptoms associated with benign prostatic hyperplasia: a discrete choice experiment. 2016 , 10, 2407-2417		6

83	New therapeutic strategies for the treatment of male lower urinary tract symptoms. 2016 , 8, 51-9		7
82	Prevalence of storage lower urinary tract symptoms in male patients attending Spanish urology office. Urinary urgency as predictor of quality of life. <i>Actas Urológicas Españolas (English Edition)</i> , 2016 , 40, 621-627	0.1	0
81	The impact of surgical treatments for lower urinary tract symptoms/benign prostatic hyperplasia on male erectile function: A systematic review and network meta-analysis. 2016 , 95, e3862		29
80	Silodosina en el tratamiento de los síntomas del tracto urinario inferior en el varón por hiperplasia benigna de próstata: enfoque estándar y uso en la «vida real». 2016 , 76, 352-359		1
79	Statin Medications and Development and Progression of Benign Prostatic Hyperplasia and Lower Urinary Tract Symptoms. 2016 , 11, 146-152		
78	Antimuscarinics, β Agonists, and Phosphodiesterase Inhibitors in the Treatment of Male Lower Urinary Tract Symptoms: An Evolving Paradigm. 2016 , 43, 337-49		3
77	Responder and health-related quality of life analyses in men with lower urinary tract symptoms treated with a fixed-dose combination of solifenacin and tamsulosin oral-controlled absorption system: results from the NEPTUNE study. <i>BJU International</i> , 2016 , 117, 165-72	5.6	11
76	Prevalence of storage lower urinary tract symptoms in male patients attending Spanish urology office. Urinary urgency as predictor of quality of life. <i>Actas Urológicas Españolas</i> , 2016 , 40, 621-627	0.7	10
75	Factors affecting help-seeking behaviour of women with urinary incontinence; a commentary providing insights for osteopaths. 2016 , 21, 51-57		1
74	The diagnosis of benign prostatic obstruction: Development of a clinical nomogram. <i>Neurourology and Urodynamics</i> , 2016 , 35, 235-40	2.3	13
73	Cardiovascular safety in refractory incontinent patients with overactive bladder receiving add-on mirabegron therapy to solifenacin (BESIDE). <i>International Journal of Clinical Practice</i> , 2017 , 71, e12944	2.9	15
72	Underactive bladder, detrusor underactivity, definition, symptoms, epidemiology, etiopathogenesis, and risk factors. 2017 , 27, 293-299		31
71	Predictors of Improvement in Storage Symptoms at Three Years After 120W GreenLight High Performance System Laser Treatment for Benign Prostate Hyperplasia. <i>Journal of Endourology</i> , 2017 , 31, 666-673	2.7	2
70	Mirabegron for the treatment of overactive bladder: a review of efficacy, safety and tolerability with a focus on male, elderly and antimuscarinic poor-responder populations, and patients with OAB in Asia. 2017 , 10, 131-151		27
69	Efficacy and safety of daily mirabegron 50 mg in male patients with overactive bladder: a critical analysis of five phase III studies. <i>Therapeutic Advances in Urology</i> , 2017 , 9, 137-154	3.2	29
68	Improvement of severe baseline lower urinary tract symptoms following robotic-assisted laparoscopic prostatectomy. <i>Neurourology and Urodynamics</i> , 2017 , 36, 1382-1386	2.3	1
67	Best practice in the management of storage symptoms in male lower urinary tract symptoms: a review of the evidence base. <i>Therapeutic Advances in Urology</i> , 2018 , 10, 79-92	3.2	18
66	Prevalence and Characteristics of Urinary Incontinence in a Treatment Seeking Male Prospective Cohort: Results from the LURN Study. <i>Journal of Urology</i> , 2018 , 200, 397-404	2.5	11

65	Diagnosis and treatment of male patients with lower urinary tract symptoms in Spain - The MERCURY Study. Do urologists follow the recommendations of the European guidelines?. <i>Actas Urológicas Españolas</i> , 2018 , 42, 323-330	0.7	
64	Non-invasive evaluation of lower urinary tract symptoms (LUTS) in men. 2018 , 5, 42-47		7
63	Healthcare resource utilization and cost among males with lower urinary tract symptoms with a predominant storage component in Spain: The epidemiological, cross-sectional MERCURY study. <i>Neurourology and Urodynamics</i> , 2018 , 37, 307-315	2.3	6
62	Underactive bladder: Pathophysiology and clinical significance. 2018 , 5, 17-21		16
61	The prevalence of lower urinary tract symptoms (LUTS) in Brazil: Results from the epidemiology of LUTS (Brazil LUTS) study. <i>Neurourology and Urodynamics</i> , 2018 , 37, 1356-1364	2.3	28
60	Reasons for Seeking Clinical Care for Lower Urinary Tract Symptoms: A Mixed Methods Study. <i>Journal of Urology</i> , 2018 , 199, 528-535	2.5	11
59	[Epidemiology of benign prostatic hyperplasia]. <i>Progres En Urologie</i> , 2018 , 28, 803-812	0.9	12
58	Postmicturition Dribble Is Associated with Erectile Dysfunction in Middle-Aged and Older Men with Lower Urinary Tract Symptoms. 2018 , 36, 263-270		5
57	The Prevention of Lower Urinary Tract Symptoms (PLUS) in girls and women: Developing a conceptual framework for a prevention research agenda. <i>Neurourology and Urodynamics</i> , 2018 , 37, 2951-2964	2.3	25
56	A Novel Research Definition of Bladder Health in Women and Girls: Implications for Research and Public Health Promotion. 2018 , 27, 974-981		74
55	Urological symptoms among 23,240 men in the general danish population - concerns about symptoms, their persistence and influence on primary care contacts. 2018 , 36, 227-236		9
54	Medical Aspects of the Treatment of LUTS/BPH: Combination Therapies. 2018 , 217-239		
53	Diagnosis and treatment of male patients with lower urinary tract symptoms in Spain □The MERCURY Study. Do urologists follow the recommendations of the European guidelines?. <i>Actas Urológicas Españolas (English Edition)</i> , 2018 , 42, 323-330	0.1	
52	Initial management of lower urinary tract symptoms and bladder outlet obstruction. <i>Medicine</i> , 2019 , 47, 541-545	0.6	
51	Vaporization of Prostate by 160W GreenLight Laser on Postoperative Erectile Function-A Single Center Report. <i>Urology</i> , 2019 , 132, 164-169	1.6	2
50	Effect of tadalafil 5 mg on post-micturition dribble in men with lower urinary tract symptoms: a multicentre, double-blind, randomized, placebo-controlled trial. <i>BJU International</i> , 2019 , 124, 862-869	5.6	4
49	Friction underwear for ease of pulling down in elderly patients with overactive bladder: A prospective randomized control trial. <i>Investigative and Clinical Urology</i> , 2019 , 60, 216-221	1.9	
48	A current perspective on post-micturition dribble in males. <i>Investigative and Clinical Urology</i> , 2019 , 60, 142-147	1.9	3

47	Impact of LUTS on treatment-related behaviors and quality of life: A population-based study in Brazil. <i>Neurourology and Urodynamics</i> , 2019 , 38, 1579-1587	2.3	8
46	Sexual function and urinary incontinence complaints and other urinary tract symptoms of perimenopausal Turkish women. <i>Psychology, Health and Medicine</i> , 2019 , 24, 1111-1122	2.1	3
45	Vesomni improves the quality of life in men with lower urinary tract symptoms in routine clinical practice in Europe. <i>Neurourology and Urodynamics</i> , 2019 , 38, 981-989	2.3	4
44	Is the belief that urinary incontinence is normal for ageing related to older Canadian women's experience of urinary incontinence?. <i>International Urogynecology Journal</i> , 2019 , 30, 2157-2160	2	9
43	Efficacy and safety of fexapotide trifluate in outpatient medical treatment of male lower urinary tract symptoms associated with benign prostatic hyperplasia. <i>Therapeutic Advances in Urology</i> , 2019 , 11, 1756287218820807	3.2	3
42	Comparison in the efficacy of fesoterodine or mirabegron add-on therapy to silodosin for patients with benign prostatic hyperplasia complicated by overactive bladder: A randomized, prospective trial using urodynamic studies. <i>Neurourology and Urodynamics</i> , 2019 , 38, 941-949	2.3	6
41	A Survey on Voiding Complaints in Women Presenting at a Pelvic Care Center. <i>Current Urology</i> , 2019 , 13, 31-36	1.7	1
40	Impact of lower urinary tract symptoms on mortality: a 21-year follow-up among middle-aged and elderly Finnish men. <i>Prostate Cancer and Prostatic Diseases</i> , 2019 , 22, 317-323	6.2	5
39	Effect of lower urinary tract symptoms on the quality of life and sexual function of males in China, Taiwan, and South Korea: Subgroup analysis of a cross-sectional, population-based study. <i>LUTS: Lower Urinary Tract Symptoms</i> , 2019 , 11, O78-O84	1.9	8
38	Feeling of incomplete emptying with little post-void residual in patients with lower urinary tract symptoms: Clinical implications and treatment outcomes. <i>LUTS: Lower Urinary Tract Symptoms</i> , 2019 , 11, O59-O64	1.9	4
37	The Role of Urinary Microbiota in Lower Urinary Tract Dysfunction: A Systematic Review. <i>European Urology Focus</i> , 2020 , 6, 361-369	5.1	16
36	Prevalence of Lower Urinary Tract Symptoms in Pregnant Adolescents and the Influencing Factors. <i>Journal of Pediatric and Adolescent Gynecology</i> , 2020 , 33, 160-166	2	6
35	Prevalence and bother of lower urinary tract symptoms and overactive bladder in Poland, an Eastern European Study. <i>Scientific Reports</i> , 2020 , 10, 19819	4.9	14
34	A retrospective analysis comparing persistence and adherence to treatment with free- vs fixed-dose combination of an alpha blocker and an antimuscarinic agent in men with LUTS in Spain. <i>International Journal of Clinical Practice</i> , 2020 , 74, e13616	2.9	2
33	Prevalence, Bother and Treatment Behavior Related to Lower Urinary Tract Symptoms and Overactive Bladder among Cardiology Patients. <i>Journal of Clinical Medicine</i> , 2020 , 9,	5.1	2
32	Lower urinary tract symptoms and related factors in female university students. <i>LUTS: Lower Urinary Tract Symptoms</i> , 2020 , 12, 211-217	1.9	1
31	Diagnostic Potential of Imaging Modalities in the Assessment of Lower Urinary Tract Dysfunctions. 2020 ,		
30	Sex Differences in Prevalence and Patient Behavior Regarding Lower Urinary Tract Symptoms Among Japanese Medical Checkup Examinees. <i>Urology</i> , 2021 , 151, 24-30	1.6	0

29	Improving management of the genitourinary syndrome of menopause: evaluation of a health system-based, cluster-randomized intervention. <i>American Journal of Obstetrics and Gynecology</i> , 2021 , 224, 62.e1-62.e13	6.4	2
28	Incidence of Pelvic Floor Disorders in US Army Female Soldiers. <i>Urology</i> , 2021 , 150, 158-164	1.6	0
27	Treatment patterns for lower urinary tract symptoms and overactive bladder in an Eastern European country: a nationwide population-representative survey. <i>Central European Journal of Urology</i> , 2021 , 74, 382-387	0.9	
26	Male Overactive Bladder: Underappreciated, Under-researched. More Please?. <i>European Urology</i> , 2021 , 79, 505-506	10.2	0
25	Impacts of nocturia on quality of life, mental health, work limitation, and health care seeking in China, Taiwan and South Korea (LUTS Asia): Results from a cross-sectional, population-based study. <i>Journal of the Formosan Medical Association</i> , 2021 , 121, 285-285	3.2	3
24	Proctored Step by Step Training Program for GreenLight Laser Anatomic Photovaporization of the Prostate: A Single Surgeon's Experience. <i>Frontiers in Surgery</i> , 2021 , 8, 705105	2.3	0
23	Is there concordance between overactive bladder and detrusor overactivity in men with predominant storage urinary symptoms referred to Functional Urology and Urodynamics Units?. <i>Actas Urológicas Españolas</i> , 2021 ,	0.7	
22	The characteristics and risk factors of healthcare-seeking men with lower urinary tract symptoms in China: Initial report from the POInT group. <i>Neurourology and Urodynamics</i> , 2021 , 40, 1740-1753	2.3	0
21	Adverse events associated with mirabegron 50mg versus placebo: A systematic review and meta-analysis. <i>Progres En Urologie</i> , 2021 , 31, 627-633	0.9	1
20	The use of mono- and combination drug therapy in men and women with lower urinary tract symptoms (LUTS) in the UK: a retrospective observational study. <i>BMC Urology</i> , 2021 , 21, 119	2.2	0
19	Physiopathology of Underactive Bladder. <i>Urodynamics, Neurourology and Pelvic Floor Dysfunctions</i> , 2021 , 97-104	0.1	
18	Relationship between Lower Urinary Tract Symptoms and Treatment-Related Behavior in an Eastern European Country: Findings from the LUTS POLAND Study. <i>International Journal of Environmental Research and Public Health</i> , 2021 , 18,	4.6	4
17	Overactive Bladder. 2012 , 1947-1957.e2		4
16	A New Brief Clinical Assessment of Lower Urinary Tract Symptoms for Women and Men: LURN SI-10. <i>Journal of Urology</i> , 2020 , 203, 164-170	2.5	6
15	Add-on treatment with mirabegron may improve quality of life in patients with benign prostatic hyperplasia complaining of persistent storage symptoms after tamsulosin monotherapy. <i>Therapeutic Advances in Urology</i> , 2020 , 12, 1756287220974130	3.2	3
14	Comparative effectiveness of oral drug therapies for lower urinary tract symptoms due to benign prostatic hyperplasia: a systematic review and network meta-analysis. <i>PLoS ONE</i> , 2014 , 9, e107593	3.7	33
13	Relationship between lifestyle and health factors and severe lower urinary tract symptoms (LUTS) in 106,435 middle-aged and older Australian men: population-based study. <i>PLoS ONE</i> , 2014 , 9, e109278	3.7	34
12	Systematic review of therapy for men with overactive bladder. <i>Canadian Urological Association Journal</i> , 2011 , 5, S143-5	1.2	10

11 Introduction. **2012**, 1-19

10	PREVALENCE OF CHILDHOOD OVERWEIGHT AND OBESITY IN RURAL PUNE. <i>Indian Journal of Child Health</i> , 2016 , 03, 301-304	0.1	4
9	The impact of adding solifenacin to tamsulosin therapy for treatment of storage lower urinary tract symptoms owing to benign prostatic hyperplasia. <i>African Journal of Urology</i> , 2020 , 26,	1	
8	Overactive Bladder and Outlet Obstruction in Men. <i>Current Urology Reports</i> ,	2.9	
7	The Role of Apoptosis in Detrusor Contractility. <i>International Neurourology Journal</i> , 2020 , 24, 67-71	2.6	0
6	Complementary and alternative therapies for urinary symptoms: use in a diverse population sample qualitative study. <i>Urologic Nursing</i> , 2012 , 32, 149-57	1.3	3
5	Is there concordance between overactive bladder and detrusor overactivity in men with predominant storage urinary symptoms referred to Functional Urology and Urodynamics Units?. <i>Actas Urológicas Españolas (English Edition)</i> , 2021 , 46, 41-41	0.1	
4	Change in postoperative storage symptoms and de novo urge-incontinence after Thulium: YAG laser enucleation of the prostate: results from a prospective, multicenter study.. <i>Journal of Endourology</i> , 2022 ,	2.7	
3	A pilot trial of movement-based pelvic floor physical therapy to address pelvic floor myofascial pain and lower urinary tract symptoms.		0
2	Impact of Female Stress Urinary Incontinence on Quality of Life, Mental Health, Work Limitation, and Healthcare Seeking in China, Taiwan, and South Korea (LUTS Asia): Results from a Cross-Sectional, Population-Based Study. Volume 14, 1871-1880		0
1	Lower Urinary Tract Symptoms and Overactive Bladder in a Large Cohort of Older Poles. Representative Tele-Survey. 2023 , 12, 2859		0