

Association of C-reactive protein and metabolic syndrome

Clinical Biochemistry

42, 976-983

DOI: [10.1016/j.clinbiochem.2009.03.026](https://doi.org/10.1016/j.clinbiochem.2009.03.026)

Citation Report

#	ARTICLE	IF	CITATIONS
1	Abdominal obesity and other risk factors largely explain the high CRP in Indigenous Australians relative to the general population, but not gender differences: a cross-sectional study. BMC Public Health, 2010, 10, 700.	1.2	24
2	High-sensitivity c-reactive protein and gamma-glutamyl transferase levels are synergistically associated with metabolic syndrome in community-dwelling persons. Cardiovascular Diabetology, 2010, 9, 87.	2.7	33
3	The New Unified International Diabetes Federation/American Heart Association/National Heart, Lung, and Blood Institute Metabolic Syndrome Definition: Does it Correlate Better with C-Reactive Protein in Chinese Patients Diagnosed with Type 2 Diabetes?. Journal of International Medical Research, 2010, 38, 1923-1932.	0.4	7
4	Relationship of body composition and C-reactive protein with pulmonary function. Respiratory Medicine, 2010, 104, 1197-1203.	1.3	17
5	Characteristics of Obesity and Its Related Disorders in China. Biomedical and Environmental Sciences, 2010, 23, 4-11.	0.2	33
6	Role of C-reactive protein in cerebrovascular disease: a critical review. Expert Review of Cardiovascular Therapy, 2011, 9, 1565-1584.	0.6	65
7	Association of inflammation and endothelial dysfunction with metabolic syndrome, prediabetes and diabetes in adults from Inner Mongolia, China. BMC Endocrine Disorders, 2011, 11, 16.	0.9	19
8	Central obesity and smoking are key modifiable risk factors for elevated C-reactive protein in Asian individuals who are not eligible for statin therapy. Nutrition and Diabetes, 2011, 1, e8-e8.	1.5	5
9	Relationship of C-reactive protein with components of the metabolic syndrome in a Tunisian population. European Journal of Internal Medicine, 2012, 23, e5-e9.	1.0	17
10	Inflammatory markers in metabolic syndrome. International Journal of Diabetes in Developing Countries, 2012, 32, 131-137.	0.3	4
11	Body composition, C-reactive protein, carotid artery remodeling and subclinical atherosclerosis in a general Taiwanese population. Journal of Thrombosis and Thrombolysis, 2012, 33, 185-192.	1.0	8
12	Metabolic syndrome, C-reactive protein and microalbuminuria in a rural Chinese population: a cross-sectional study. BMC Nephrology, 2013, 14, 118.	0.8	12
13	Association of Ideal Cardiovascular Metrics and Serum High-Sensitivity C-Reactive Protein in Hypertensive Population. PLoS ONE, 2013, 8, e81597.	1.1	27
14	Ideal cardiovascular health behaviors and factors and high sensitivity C-reactive protein: the Kailuan cross-sectional study in Chinese. Clinical Chemistry and Laboratory Medicine, 2014, 52, 1379-86.	1.4	10
15	Leptin, Insulin and Lipid Profiles in Obese Subjects with and without Metabolic Syndrome in the Region of Cap-Bon: Tunisia. Endocrinology & Metabolic Syndrome: Current Research, 2015, 04, .	0.3	3
16	Comparisons of Different Metabolic Syndrome Definitions and Associations with Coronary Heart Disease, Stroke, and Peripheral Arterial Disease in a Rural Chinese Population. PLoS ONE, 2015, 10, e0126832.	1.1	28
17	High-sensitive C-reactive protein and risk of incident type 2 diabetes: a case-control study nested within the Singapore Chinese Health Study. BMC Endocrine Disorders, 2017, 17, 8.	0.9	25
18	Mast cell specific immunological biomarkers and metabolic syndrome among middle-aged and older Chinese adults. Endocrine Journal, 2017, 64, 245-253.	0.7	5

#	ARTICLE	IF	CITATIONS
19	<p>Correlation Between Ideal Cardiovascular Health Metrics and Plasma hs-CRP Levels in a North China Population: One Four-Year Follow-Up Study</p>. International Journal of General Medicine, 2020, Volume 13, 617-625.	0.8	3
20	<p>High-Sensitivity C-Reactive Protein Leads to Increased Incident Metabolic Syndrome in Women but Not in Men: A Five-Year Follow-Up Study in a Chinese Population</p>. Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy, 2020, Volume 13, 581-590.	1.1	21
21	Risk scores for predicting incident chronic kidney disease among rural Chinese people: a village-based cohort study. BMC Nephrology, 2020, 21, 120.	0.8	10
22	High sensitivity C-reactive protein associated with different health predictors in middle-aged and oldest old Chinese. Biomedical and Environmental Sciences, 2012, 25, 257-66.	0.2	8