

# CITATION REPORT

List of articles citing

Relaxant effects of sildenafil on the human isolated bladder neck

DOI: 10.1016/j.urology.2008.06.060  
Urology, 2009, 73, 427-30.

**Source:** <https://exaly.com/paper-pdf/46704161/citation-report.pdf>

**Version:** 2024-04-26

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
26	Current world literature. <i>Current Opinion in Urology</i> , <b>2010</b> , 20, 98-103	2.8	
25	[Treatment of the lower urinary tract symptoms secondary to benign prostatic hyperplasia by phosphodiesterase type 5 inhibitors. Review article]. <i>Progres En Urologie</i> , <b>2010</b> , 20, 616-26	0.9	2
24	Effects of tadalafil on lower urinary tract symptoms secondary to benign prostatic hyperplasia in men with or without erectile dysfunction. <i>Urology</i> , <b>2010</b> , 75, 1452-8	1.6	51
23	Phosphodiesterase Type 5 Inhibitors for Lower Urinary Tract Symptoms Associated With Benign Prostatic Hyperplasia. <i>Urological Science</i> , <b>2010</b> , 21, 2-7	0.3	
22	Phosphodiesterase (PDE) inhibitors in the treatment of lower urinary tract dysfunction. <i>British Journal of Clinical Pharmacology</i> , <b>2011</b> , 72, 197-204	3.8	43
21	Current benign prostatic hyperplasia treatment: impact on sexual function and management of related sexual adverse events. <i>International Journal of Clinical Practice</i> , <b>2011</b> , 65, 1005-13	2.9	34
20	Phosphodiesterase type 5 inhibitors in the management of non-neurogenic male lower urinary tract symptoms: critical analysis of current evidence. <i>European Urology</i> , <b>2011</b> , 60, 527-35	10.2	42
19	Tadalafil for the treatment of lower urinary tract symptoms secondary to benign prostatic hyperplasia: pathophysiology and mechanism(s) of action. <i>Neurourology and Urodynamics</i> , <b>2011</b> , 30, 292-301	3.1	162
18	Sildenafil effect on the human bladder involves the L-cysteine/hydrogen sulfide pathway: a novel mechanism of action of phosphodiesterase type 5 inhibitors. <i>European Urology</i> , <b>2012</b> , 62, 1174-80	10.2	58
17	Phosphodiesterase inhibitors for lower urinary tract symptoms consistent with benign prostatic hyperplasia. <i>The Cochrane Library</i> , <b>2012</b> ,	5.2	1
16	Les inhibiteurs de la phosphodiesterase de type 5 : une révolution dans le traitement des symptômes du bas appareil urinaire?. <i>Basic and Clinical Andrology</i> , <b>2012</b> , 22, 80-91	2.8	1
15	Prophylactic effect of tadalafil on bladder function in a rat model of chronic bladder ischemia. <i>Journal of Urology</i> , <b>2013</b> , 189, 754-61	2.5	50
14	The mechanism of action of phosphodiesterase type 5 inhibitors in the treatment of lower urinary tract symptoms related to benign prostatic hyperplasia. <i>European Urology</i> , <b>2013</b> , 63, 506-16	10.2	103
13	The role of phosphodiesterases in bladder pathophysiology. <i>Nature Reviews Urology</i> , <b>2013</b> , 10, 414-24	5.5	22
12	Nitric Oxide Synthase is Necessary for Normal Urogenital Development. <b>2013</b> , 2, 108		1
11	A comprehensive review of urologic complications in patients with diabetes. <i>SpringerPlus</i> , <b>2014</b> , 3, 549		38
10	Efficacy of daily low-dose sildenafil for treating interstitial cystitis: results of a randomized, double-blind, placebo-controlled trial--treatment of interstitial cystitis/painful bladder syndrome with low-dose sildenafil. <i>Urology</i> , <b>2014</b> , 84, 51-6	1.6	26

9	Nightly sildenafil use after radical prostatectomy has adverse effects on urinary convalescence: Results from a randomized trial of nightly vs on-demand dosing regimens. <i>Canadian Urological Association Journal</i> , <b>2015</b> , 9, 414-9	1.2	4
8	Tadalafil for lower urinary tract symptoms secondary to benign prostatic hyperplasia: a review of clinical data in Asian men and an update on the mechanism of action. <i>Therapeutic Advances in Urology</i> , <b>2015</b> , 7, 249-64	3.2	17
7	Current and emerging drugs for interstitial cystitis/bladder pain syndrome (IC/BPS). <i>Expert Opinion on Emerging Drugs</i> , <b>2015</b> , 20, 555-70	3.7	21
6	Impact of postoperative phosphodiesterase type 5 inhibitor treatment on lower urinary tract symptoms after robot-assisted radical prostatectomy: a longitudinal study. <i>Scandinavian Journal of Urology</i> , <b>2017</b> , 51, 33-37	1.6	5
5	Phosphodiesterase inhibitors for lower urinary tract symptoms consistent with benign prostatic hyperplasia. <i>The Cochrane Library</i> , <b>2018</b> , 11, CD010060	5.2	8
4	Objective impacts of tadalafil on storage and voiding function in male patients with benign prostatic hyperplasia: 1-year outcomes from a prospective urodynamic study. <i>World Journal of Urology</i> , <b>2019</b> , 37, 867-872	4	12
3	Mirodenafil prevents bladder dysfunction induced by chronic bladder ischemia in rats. <i>International Neurourology Journal</i> , <b>2015</b> , 19, 19-26	2.6	4
2	Effect of sildenafil citrate in testosterone induced benign prostate hyperplasia rat model. <i>Turkish Journal of Urology</i> , <b>2017</b> , 43, 434-438	1.3	
1	Efficacy of Tadalafil on Ureteral Stent Symptoms: A Randomized Controlled Trial. <i>Nephro-Urology Monthly</i> , <b>2019</b> , In Press,	0.4	