Clinical and dosimetric predictors of late rectal toxicity localized prostate cancer: Results of a large multicenter

Radiotherapy and Oncology 93, 197-202 DOI: 10.1016/j.radonc.2009.09.004

Citation Report

CITATION REDORT

| # | Article | IF | CITATIONS |
|----|--|------|-----------|
| 1 | Dose–volume effects for normal tissues in external radiotherapy: Pelvis. Radiotherapy and Oncology, 2009, 93, 153-167. | 0.6 | 249 |
| 2 | Gastrointestinal toxicity associated to radiation therapy. Clinical and Translational Oncology, 2010, 12, 554-561. | 2.4 | 30 |
| 3 | Early clinical experience of radiotherapy of prostate cancer with volumetric modulated arc therapy. Radiation Oncology, 2010, 5, 54. | 2.7 | 35 |
| 4 | Treating prostate cancer with radiotherapy. Nature Reviews Clinical Oncology, 2010, 7, 583-589. | 27.6 | 17 |
| 6 | A comparison of dose–volume constraints derived using peak and longitudinal definitions of late rectal toxicity. Radiotherapy and Oncology, 2010, 94, 241-247. | 0.6 | 37 |
| 7 | Deformable image registration for contour propagation from CT to cone-beam CT scans in radiotherapy of prostate cancer. Acta OncolÃ ³ gica, 2011, 50, 918-925. | 1.8 | 118 |
| 8 | Faecal incontinence following radiotherapy for prostate cancer: A systematic review. Radiotherapy and Oncology, 2011, 98, 145-153. | 0.6 | 41 |
| 9 | Does the planning dose–volume histogram represent treatment doses in image-guided prostate radiation therapy? Assessment with cone-beam computerised tomography scans. Radiotherapy and Oncology, 2011, 98, 162-168. | 0.6 | 68 |
| 10 | Tobacco smoking and long-lasting symptoms from the bowel and the anal-sphincter region after radiotherapy for prostate cancer. Radiotherapy and Oncology, 2011, 101, 495-501. | 0.6 | 58 |
| 11 | Inclusion of clinical risk factors into NTCP modelling of late rectal toxicity after high dose radiotherapy for prostate cancer. Radiotherapy and Oncology, 2011, 100, 124-130. | 0.6 | 65 |
| 12 | Anatomical and clinical predictors of acute bowel toxicity in whole pelvis irradiation for prostate cancer with Tomotherapy. Radiotherapy and Oncology, 2011, 101, 460-464. | 0.6 | 21 |
| 13 | The Impact of Clinical Factors on the Development of Late Radiation Toxicity: Results from the Medical Research Council RT01 Trial (ISRCTN47772397). Clinical Oncology, 2011, 23, 613-624. | 1.4 | 88 |
| 14 | Prostate Image-Guided Radiotherapy by Megavolt Cone-Beam CT. Strahlentherapie Und Onkologie, 2011, 187, 473-478. | 2.0 | 19 |
| 15 | The evolution of rectal and urinary toxicity and immune response in prostate cancer patients treated with two three-dimensional conformal radiotherapy techniques. Radiation Oncology, 2011, 6, 87. | 2.7 | 6 |
| 16 | Feasibility of safe ultra-high (EQD ₂ >100 Gy) dose escalation on dominant intra-prostatic lesions (DILs) by Helical Tomotheraphy. Acta Oncológica, 2011, 50, 25-34. | 1.8 | 42 |
| 17 | Late rectal bleeding after 3D-CRT for prostate cancer: development of a neural-network-based predictive model. Physics in Medicine and Biology, 2012, 57, 1399-1412. | 3.0 | 44 |
| 18 | Consolidating Risk Estimates for Radiation-Induced Complications in Individual Patient: Late Rectal Toxicity. International Journal of Radiation Oncology Biology Physics, 2012, 83, 53-63. | 0.8 | 15 |
| 19 | Is It Time to Tailor the Prediction of Radio-Induced Toxicity in Prostate Cancer Patients? Building the First Set of Nomograms for Late Rectal Syndrome. International Journal of Radiation Oncology Biology Physics, 2012, 82, 1957-1966. | 0.8 | 41 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 20 | The Benefits of Including Clinical Factors in Rectal Normal Tissue Complication Probability Modeling After Radiotherapy for Prostate Cancer. International Journal of Radiation Oncology Biology Physics, 2012, 82, 1233-1242. | 0.8 | 68 |
| 21 | Late Fecal Incontinence After High-Dose Radiotherapy for Prostate Cancer: Better Prediction Using Longitudinal Definitions. International Journal of Radiation Oncology Biology Physics, 2012, 83, 38-45. | 0.8 | 38 |
| 22 | Do Intermediate Radiation Doses Contribute to Late Rectal Toxicity? An Analysis of Data From Radiation Therapy Oncology Group Protocol 94-06. International Journal of Radiation Oncology Biology Physics, 2012, 84, 390-395. | 0.8 | 26 |
| 23 | Mean Absorbed Dose to the Anal-Sphincter Region and Fecal Leakage among Irradiated Prostate Cancer Survivors. International Journal of Radiation Oncology Biology Physics, 2012, 84, e181-e185. | 0.8 | 24 |
| 24 | Increasing the risk of late rectal bleeding after high-dose radiotherapy for prostate cancer: The case of previous abdominal surgery. Results from a prospective trial. Radiotherapy and Oncology, 2012, 103, 252-255. | 0.6 | 39 |
| 26 | Management of Intestinal Complications in Patients With Pelvic Radiation Disease. Clinical Gastroenterology and Hepatology, 2012, 10, 1326-1334.e4. | 4.4 | 47 |
| 27 | Functional Outcomes and Complications Following Radiation Therapy for Prostate Cancer: A Critical Analysis of the Literature. European Urology, 2012, 61, 112-127. | 1.9 | 224 |
| 28 | Clinical risk factors for late intestinal toxicity after radiotherapy: a systematic review protocol. Systematic Reviews, 2013, 2, 39. | 5.3 | 8 |
| 29 | Late rectal and bladder toxicity following radiation therapy for prostate cancer: Predictive factors and treatment results. Reports of Practical Oncology and Radiotherapy, 2013, 18, 298-303. | 0.6 | 22 |
| 30 | Multivariate normal tissue complication probability modeling of gastrointestinal toxicity after external beam radiotherapy for localized prostate cancer. Radiation Oncology, 2013, 8, 221. | 2.7 | 34 |
| 31 | Role of Early Proctoscopy in Predicting Late Symptomatic Proctitis After External Radiation Therapy for Prostate Carcinoma. International Journal of Radiation Oncology Biology Physics, 2013, 85, 1031-1037. | 0.8 | 9 |
| 32 | Quality of care indicators and their related outcomes: A population-based study in prostate cancer patients treated with radiotherapy. Radiotherapy and Oncology, 2013, 107, 358-365. | 0.6 | 10 |
| 33 | Reducing rectal injury during external beam radiotherapy for prostate cancer. Nature Reviews Urology, 2013, 10, 345-357. | 3.8 | 13 |
| 34 | Statistical simulations to estimate motion-inclusive dose-volume histograms for prediction of rectal morbidity following radiotherapy. Acta Oncológica, 2013, 52, 666-675. | 1.8 | 18 |
| 36 | Quantifying daily variation in volume and dose to the prostate, rectum and bladder using cone-beam computerised tomography. Journal of Radiotherapy in Practice, 2014, 13, 79-86. | 0.5 | 11 |
| 37 | Rectal toxicity after intensity modulated radiotherapy for prostate cancer: Which rectal dose volume constraints should we use?. Radiotherapy and Oncology, 2014, 113, 398-403. | 0.6 | 28 |
| 38 | Transitioning from conventional radiotherapy to intensity-modulated radiotherapy for localized prostate cancer: changing focus from rectal bleeding to detailed quality of life analysis. Journal of Radiation Research, 2014, 55, 1033-1047. | 1.6 | 26 |
| 40 | Long term rectal function after high-dose prostatecancer radiotherapy: Results from a prospective cohort study. Radiotherapy and Oncology, 2014, 110, 272-277. | 0.6 | 30 |

CITATION REPORT

CITATION REPORT

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 41 | MRI target delineation may reduce long-term toxicity after prostate radiotherapy. Acta Oncológica, 2014, 53, 809-814. | 1.8 | 51 |
| 42 | The REQUITE Project: Validating Predictive Models and Biomarkers of Radiotherapy Toxicity to Reduce Side-effects and Improve Quality of Life in Cancer Survivors. Clinical Oncology, 2014, 26, 739-742. | 1.4 | 73 |
| 43 | Recommendations for Prostate Cancer Survivorship Care: An Update to the 2009 Michigan Cancer Consortium Guidelines for the Primary Care Management of Prostate Cancer Post-Treatment Sequelae. Journal of Men's Health, 2014, 11, 95-107. | 0.3 | 7 |
| 44 | Pathogenesis, Diagnosis, and Management of Ulcerative Proctitis, Chronic Radiation Proctopathy, and Diversion Proctitis. Inflammatory Bowel Diseases, 2015, 21, 703-715. | 1.9 | 49 |
| 45 | Improved Beam Angle Arrangement in Intensity Modulated Proton Therapy Treatment Planning for Localized Prostate Cancer. Cancers, 2015, 7, 574-584. | 3.7 | 20 |
| 46 | Systematic Review of the Relationship between Acute and Late Gastrointestinal Toxicity after Radiotherapy for Prostate Cancer. Prostate Cancer, 2015, 2015, 1-11. | 0.6 | 37 |
| 47 | Prediction of gastrointestinal toxicity after external beam radiotherapy for localized prostate cancer. Radiation Oncology, 2015, 10, 80. | 2.7 | 24 |
| 48 | Rectal bleeding after radiation therapy for endometrial cancer. Radiotherapy and Oncology, 2015, 115, 240-245. | 0.6 | 7 |
| 49 | Contrasting analytical and data-driven frameworks for radiogenomic modeling of normal tissue toxicities in prostate cancer. Radiotherapy and Oncology, 2015, 115, 107-113. | 0.6 | 24 |
| 50 | Predictive Parameters of Symptomatic Hematochezia Following 5-Fraction Gantry-Based SABR in Prostate Cancer. International Journal of Radiation Oncology Biology Physics, 2016, 94, 1043-1051. | 0.8 | 33 |
| 51 | Modelling late stool frequency and rectal pain after radical radiotherapy in prostate cancer patients: Results from a large pooled population. Physica Medica, 2016, 32, 1690-1697. | 0.7 | 12 |
| 52 | Clinical and dosimetric predictors of late rectal bleeding of prostate cancer after TomoTherapy intensity modulated radiation therapy. Journal of Medical Radiation Sciences, 2017, 64, 172-179. | 1.5 | 6 |
| 53 | Comparison of different contouring definitions of the rectum as organ at risk (OAR) and dose–volume parameters predicting rectal inflammation in radiotherapy of prostate cancer: which definition to use?. British Journal of Radiology, 2017, 90, 20160370. | 2.2 | 7 |
| 54 | Linking CHHiP prostate cancer RCT with GP records: A study proposal to investigate the effect of co-morbidities and medications on long-term symptoms and radiotherapy-related toxicity. Technical Innovations and Patient Support in Radiation Oncology, 2017, 2, 5-12. | 1.9 | 4 |
| 55 | Daily Rectal Dose-volume Histogram Variation in Prostate Intensity-modulated Radiation Therapy: Is It Clinically Significant in the Era of Image Guidance?. Journal of Medical Imaging and Radiation Sciences, 2017, 48, 346-351. | 0.3 | 2 |
| 56 | Factors Predicting Late Rectal Disorders after Radiation Therapy for Prostate Cancer. Chinese Medical Journal, 2017, 130, 2441-2446. | 2.3 | 2 |
| 57 | Development of a Ready-to-Use Graphical Tool Based on Artificial Neural Network Classification: Application for the Prediction of Late Fecal Incontinence After Prostate Cancer Radiation Therapy. International Journal of Radiation Oncology Biology Physics, 2018, 102, 1533-1542. | 0.8 | 14 |
| 58 | Texture analysis of 3D dose distributions for predictive modelling of toxicity rates in radiotherapy. Radiotherapy and Oncology, 2018, 129, 548-553. | 0.6 | 89 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 59 | Predicting Late Fecal Incontinence Risk After Radiation Therapy for Prostate Cancer: New Insights From External Independent Validation. International Journal of Radiation Oncology Biology Physics, 2018, 102, 127-136. | 0.8 | 14 |
| 60 | Evaluating the predictive value of quantec rectum tolerance dose suggestions on acute rectal toxicity in prostate carcinoma patients treated with IMRT. Reports of Practical Oncology and Radiotherapy, 2020, 25, 50-54. | 0.6 | 4 |
| 61 | Tools for fecal incontinence assessment: lessons for inflammatory bowel disease trials based on a systematic review. United European Gastroenterology Journal, 2020, 8, 886-922. | 3.8 | 14 |
| 62 | Radiation Dose to the Rectum With Definitive Radiation Therapy and Hydrogel Spacer Versus Postprostatectomy Radiation Therapy. Advances in Radiation Oncology, 2020, 5, 1225-1231. | 1.2 | 0 |
| 63 | Late Gastrointestinal Tolerance After Prostate Radiotherapy: Is the Anal Canal the Culprit? A Narrative Critical Review. Frontiers in Oncology, 2021, 11, 666962. | 2.8 | 0 |
| 64 | Long-term Outcomes of a Dose–reduction Trial to Decrease Late Gastrointestinal Toxicity in Patients with Prostate Cancer Receiving Soft Tissue-matched Image-guided Intensity-modulated Radiotherapy. Anticancer Research, 2018, 38, 385-391. | 1.1 | 9 |
| 65 | Predictive Factors of Late-onset Rectal Mucosal Changes After Radiotherapy of Prostate Cancer. In Vivo, 2018, 31, 961-966. | 1.3 | 4 |
| 66 | Dosimetric correlation of acute and late toxicities in high-risk prostate cancer patients treated with three-dimensional conformal radiotherapy followed by intensity modulated radiotherapy boost. Indian Journal of Urology, 2016, 32, 210. | 0.6 | 2 |
| 67 | Predictive parameters of symptomatic haematochezia following 5-fraction gantry-based SABR in prostate cancer Journal of Clinical Oncology, 2016, 34, 79-79. | 1.6 | 0 |
| 68 | Interrelatedness of Urological Conditions and Anal Incontinence. Updates in Surgery Series, 2023, , 147-154. | 0.1 | 0 |
| 69 | Clinical Outcome and Risk Factors of Chronic Radiation Proctitis Following Pelvic Radiation Therapy. Anticancer Research, 2022, 42, 5951-5959. | 1.1 | 2 |
| 70 | Validation of prediction models for radiation-induced late rectal bleeding: Evidence from a large pooled population of prostate cancer patients. Radiotherapy and Oncology, 2023, 183, 109628. | 0.6 | 1 |
| 71 | A new predictive parameter for doseâ€volume metrics in intensityâ€modulated radiation therapy planning for prostate cancer: Initial phantom study. Journal of Applied Clinical Medical Physics, 2024, 25, . | 1.9 | 0 |

CITATION REPORT