

CITATION REPORT

List of articles citing

Assessment of liver function for safe hepatic resection

DOI: 10.1111/j.1872-034x.2008.00441.x
Hepatology Research, 2009, 39, 107-16.

Source: <https://exaly.com/paper-pdf/47102360/citation-report.pdf>

Version: 2024-04-28

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
161	Preoperative portal vein embolization. 176-192		
160	Use of EDTA-plasma gel-separating tubes for measurement of indocyanine green. 2009 , 47, 1159-60		
159	Das hepatozelluläre Karzinom aus der Sicht des Internisten und des Chirurgen. 2009 , 4, 301-311		
158	Radical resection of biliary tract cancers and the role of extended lymphadenectomy. 2009 , 18, 339-59, ix		2
157	Virtual liver resection and volumetric analysis of the future liver remnant using open source image processing software. 2010 , 34, 2426-33		62
156	Uses and limitations of portal vein embolization for improving perioperative outcomes in hepatocellular carcinoma. 2010 , 37, 102-9		26
155	Prediction of mortality rate after major hepatectomy in patients without cirrhosis. 2010 , 145, 1075-81		23
154	Relationship of hepatic functional parameters with changes of functional liver volume using technetium-99m galactosyl serum albumin scintigraphy in patients undergoing preoperative portal vein embolization: a follow-up report. 2010 , 164, e235-42		19
153	A fourth dimension in decision making in hepatology. <i>Hepatology Research</i> , 2010 , 40, 1143-54	5.1	10
152	Condotta generale delle epatectomie mediante laparotomia. 2011 , 17, 1-19		
151	MR imaging of the biliary tract with Gd-EOB-DTPA: effect of liver function on signal intensity. 2011 , 77, 325-9		62
150	Prediction of indocyanine green retention rate at 15 minutes by correlated liver function parameters before hepatectomy. 2011 , 169, e119-25		13
149	Conducta general en las hepatectomías por laparotomía. 2011 , 27, 1-21		
148	Toward a total and regional scintigraphic liver function test enabling assessment of future remnant liver function. <i>Nuclear Medicine Communications</i> , 2011 , 32, 669-71	1.6	6
147	Estimation of liver function using T1 mapping on Gd-EOB-DTPA-enhanced magnetic resonance imaging. 2011 , 46, 277-83		155
146	Conduite générale des hépatectomies par laparotomie. 2011 , 6, 1-19		
145	Usefulness of a modified inflammation-based prognostic system for predicting postoperative mortality of patients undergoing surgery for primary hepatocellular carcinoma. 2011 , 103, 801-6		34

144	Underweight patients show an increased rate of postoperative death after surgery for hepatocellular carcinoma. 2011 , 104, 809-13		6
143	Duration of hepatic vascular inflow clamping and survival after liver resection for hepatocellular carcinoma. 2011 , 98, 1284-90		232
142	Duration of hepatic vascular inflow clamping and survival after liver resection for hepatocellular carcinoma (Br J Surg 2011; 98: 1284-1290). 2011 , 98, 1291		
141	Quantitative evaluation of liver function with use of gadoxetate disodium-enhanced MR imaging. 2011 , 260, 727-33		143
140	Impact of an inflammation-based prognostic system on patients undergoing surgery for hepatocellular carcinoma: a retrospective study of 398 Japanese patients. 2012 , 203, 101-6		77
139	Quantitative Evaluation of Liver Function within MR Imaging. 2012 , 01,		1
138	Liver injury due to chemotherapy-induced sinusoidal obstruction syndrome is associated with sinusoidal capillarization. <i>Annals of Surgical Oncology</i> , 2012 , 19, 2230-7	3.1	45
137	Increase in left liver lobe function after preoperative right portal vein embolisation assessed with gadolinium-EOB-DTPA MRI. 2013 , 23, 2555-60		16
136	Vascular reconstruction combined with liver resection for malignant tumours. 2013 , 100, 1764-75		34
135	Lipid profile is associated with the incidence of cognitive dysfunction in viral cirrhotic patients: A data-mining analysis. <i>Hepatology Research</i> , 2013 , 43, 418-24	5.1	11
134	New paradigms in post-hepatectomy liver failure. <i>Journal of Gastrointestinal Surgery</i> , 2013 , 17, 593-605	3.3	52
133	Developments in near-infrared-guided hepatobiliary, pancreatic and other upper gastrointestinal surgery. 2013 , 8, 211-9		5
132	Survival after surgery for hepatocellular carcinoma in relation to presence or absence of viral infection. 2013 , 206, 187-93		2
131	Virtual liver resection: computer-assisted operation planning using a three-dimensional liver representation. 2013 , 20, 157-64		63
130	Safety of hepatic resection for hepatocellular carcinoma in obese patients with cirrhosis. <i>Surgery Today</i> , 2013 , 43, 1290-7	3	19
129	[Portal vein embolisation prior to extended liver resection - current and future aspects]. 2013 , 138, 442-8		1
128	Assessment of hepatic function, operative candidacy, and medical management after liver resection in the patient with underlying liver disease. 2013 , 33, 204-12		23
127	Transjugular Retrograde Obliteration prior to Liver Resection for Hepatocellular Carcinoma Associated with Hyperammonemia due to Spontaneous Portosystemic Shunt. 2013 , 2013, 809543		

126	Can the model for end-stage liver disease score replace the indocyanine green clearance test in the selection of right hemihepatectomy in Child-Pugh class A?. 2014 , 86, 122-9		8
125	Evaluation of gadolinium-EOB-DTPA uptake after portal vein embolization: value of an increased flip angle. <i>Acta Radiologica</i> , 2014 , 55, 149-54	2	5
124	Functional discrepancy between two liver lobes after hemilobe biliary drainage in patients with jaundice and bile duct cancer: an appraisal using (99m)Tc-GSA SPECT/CT fusion imaging. 2014 , 273, 444-51		21
123	Association between pretreatment retention rate of indocyanine green 15 min after administration and life prognosis in patients with HCC treated by proton beam therapy. 2014 , 113, 54-9		13
122	Consensus report from the 6th International forum for liver MRI using gadoxetic acid. <i>Journal of Magnetic Resonance Imaging</i> , 2014 , 40, 516-29	5.6	33
121	Parenteral solution of nutritional hepatotrophic factors improves regeneration in thioacetamide-induced cirrhotic livers after partial hepatectomy. 2014 , 42, 414-21		2
120	Surgical treatment of hepatocellular carcinoma associated with hepatic vein tumor thrombosis. 2014 , 61, 583-8		103
119	Evaluation of the usefulness of the indocyanine green clearance test for chemotherapy-associated liver injury in patients with colorectal cancer liver metastasis. <i>Annals of Surgical Oncology</i> , 2014 , 21, 167-72		28
118	Efficacy of liver parenchymal enhancement and liver volume to standard liver volume ratio on Gd-EOB-DTPA-enhanced MRI for estimation of liver function. 2014 , 24, 857-65		51
117	Abdomen and Thoracic Imaging. 2014 ,		3
116	Three-dimensional volumetry in 107 normal livers reveals clinically relevant inter-segment variation in size. 2014 , 16, 439-47		42
115	Minibody-indocyanine green based activatable optical imaging probes: the role of short polyethylene glycol linkers. 2014 , 5, 411-5		28
114	Rapid reversal of neuromuscular blockade by sugammadex after continuous infusion of rocuronium in patients with liver dysfunction undergoing hepatic surgery. 2014 , 52, 54-8		15
113	Effect of microbial cell preparation on renal profile and liver function among type 2 diabetics: a randomized controlled trial. 2015 , 15, 433		22
112	The natural history of cirrhosis from parenteral nutrition-associated liver disease after resolution of cholestasis with parenteral fish oil therapy. <i>Annals of Surgery</i> , 2015 , 261, 172-9	7.8	32
111	High Levels of Gamma-Glutamyl Transferase and Indocyanine Green Retention Rate at 15 min as Preoperative Predictors of Tumor Recurrence in Patients With Hepatocellular Carcinoma. 2015 , 94, e810		24
110	Preoperative estimation of future remnant liver function following portal vein embolization using relative enhancement on gadoxetic acid disodium-enhanced magnetic resonance imaging. 2015 , 16, 523-30		14
109	Gadoxetic acid: pearls and pitfalls. 2015 , 40, 2012-29		14

108	Risk factors for incisional and organ space surgical site infections after liver resection are different. 2015 , 39, 1185-92		31
107	The relationship between liver function and liver parenchymal contrast enhancement on Gd-BOPTA-enhanced MR imaging in the hepatocyte phase. 2015 , 33, 768-73		14
106	Ultrasound-guided percutaneous microwave ablation treatment of initial recurrent hepatocellular carcinoma after hepatic resection: long-term outcomes. 2015 , 41, 2391-9		5
105	T1 ρ Relaxation of the liver: A potential biomarker of liver function. <i>Journal of Magnetic Resonance Imaging</i> , 2015 , 42, 188-95	5.6	30
104	Chronic kidney disease after liver transplantation. 2015 , 62, 243-4		4
103	Reply to: "ALPPS procedure for hepatocellular carcinoma with macrovascular thrombosis: a new opportunity?". 2015 , 62, 243		5
102	Prognostic role of Child-Pugh score 5 and 6 in hepatocellular carcinoma patients who underwent curative hepatic resection. 2015 , 209, 199-205		15
101	Portal vein embolization for induction of selective hepatic hypertrophy prior to major hepatectomy: rationale, techniques, outcomes and future directions. 2016 , 13, 426-442		15
100	Indocyanine green kinetics to assess liver function: Ready for a clinical dynamic assessment in major liver surgery?. 2016 , 8, 355-67		85
99	The Evaluation of Liver Function and Surgical Influence by ICGR15 after Chemotherapy for Colorectal Liver Metastases. 2016 , 7, 595-9		8
98	Assessment of Preoperative Liver Function in Patients with Hepatocellular Carcinoma - The Albumin-Indocyanine Green Evaluation (ALICE) Grade. 2016 , 11, e0159530		36
97	Clinical Impact of Gadoteric Acid-Enhanced Magnetic Resonance Imaging on Hepatoma Management: A Prospective Study. 2016 , 61, 1197-205		12
96	Long-term impact of liver function on curative therapy for hepatocellular carcinoma: application of the ALBI grade. 2016 , 114, 744-50		119
95	Preoperative direct bilirubin to prothrombin time ratio index to prevent liver failure after minor hepatectomy. 2016 , 23, 763-770		8
94	Semi-quantitative assessment of the presence and Child-Pugh class of hepatitis B related cirrhosis by using liver lobe-based dynamic contrast-enhanced MRI. 2016 , 71, 1289-1295		2
93	MRI-based estimation of liver function by intravoxel incoherent motion diffusion-weighted imaging. 2016 , 34, 1220-5		12
92	Hyaluronic Acid Predicts Postoperative Large Amount of Ascites after Liver Resection for Patients with Hepatocellular Carcinoma. 2016 , 33, 520-8		7
91	Current strategies for preoperative conditioning of the liver to expand criteria for resectability of hepatic metastases. 2016 , 48, 180-190		

90	Clinical feasibility of Gd-EOB-DTPA-enhanced MR imaging for assessing liver function: validation with ICG tests and parenchymal cell volume. 2016 , 40, 797-800		6
89	Liver function assessment using 99mTc-GSA single-photon emission computed tomography (SPECT)/CT fusion imaging in hilar bile duct cancer: A retrospective study. 2016 , 160, 118-126		25
88	Stepwise approach to curative surgery using percutaneous transhepatic cholangiodrainage and portal vein embolization for severe bile duct injury during laparoscopic cholecystectomy: a case report. 2016 , 2, 27		
87	Platelet count is more useful for predicting posthepatectomy liver failure at surgery for hepatocellular carcinoma than indocyanine green clearance test. 2016 , 113, 565-9		37
86	Morbidity of hepatic resection for intermediate and advanced hepatocellular carcinoma. <i>Langenbecks Archives of Surgery</i> , 2016 , 401, 43-53	3-4	4
85	Use of finger-piece method for indocyanine green clearance test. <i>Hepatology Research</i> , 2017 , 47, 1235-1240	3-4	4
84	Kinetic Changes in Liver Parenchyma After Preoperative Chemotherapy for Patients with Colorectal Liver Metastases. <i>Journal of Gastrointestinal Surgery</i> , 2017 , 21, 813-821	3-3	9
83	The response of male and female rats to a high-fructose diet during adolescence following early administration of Hibiscus sabdariffa aqueous calyx extracts. 2017 , 8, 628-637		7
82	The preoperative globulin-to-albumin ratio, a novel inflammation-based prognostic system, predicts survival after potentially curative liver resection for patients with hepatocellular carcinoma. 2017 , 116, 1166-1175		8
81	Validation and Potential of Albumin-Bilirubin Grade and Prognostication in a Nationwide Survey of 46,681 Hepatocellular Carcinoma Patients in Japan: The Need for a More Detailed Evaluation of Hepatic Function. <i>Liver Cancer</i> , 2017 , 6, 325-336	9-1	102
80	3D analysis of apparent diffusion coefficient histograms in hepatocellular carcinoma: correlation with histological grade. 2017 , 17, 1		18
79	Potential use of metabolic breath tests to assess liver disease and prognosis: has the time arrived for routine use in the clinic?. 2017 , 37, 328-336		9
78	Prognostic value of the albumin-bilirubin grade in patients with hepatocellular carcinoma: Validation in a Chinese cohort. <i>Hepatology Research</i> , 2017 , 47, 731-741	5-1	45
77	Validation of Newly Proposed Time to Transarterial Chemoembolization Progression in Intermediate-Stage Hepatocellular Carcinoma Cases. 2017 , 93 Suppl 1, 120-126		17
76	Index of convexity: A novel method for assessing liver functional reserve using technetium-99m-galactosyl human serum albumin liver scintigraphy. 2017 , 11, 333-339		2
75	The Value of the C-Reactive Protein-to-Albumin Ratio is Useful for Predicting Survival of Patients with Child-Pugh Class A Undergoing Liver Resection for Hepatocellular Carcinoma. 2018 , 42, 2218-2226		13
74	Comparison of twelve liver functional reserve models for outcome prediction in patients with hepatocellular carcinoma undergoing surgical resection. 2018 , 8, 4773		21
73	Hepatocellular Carcinoma: The Gap Between Eastern and Western Clinical Practice. <i>Annals of Surgery</i> , 2018 , 267, e27-e28	7-8	2

72	A New Proposal of Criteria for the Future Remnant Liver Volume in Older Patients Undergoing Major Hepatectomy for Biliary Tract Cancer. <i>Annals of Surgery</i> , 2018 , 267, 338-345	7.8	13
71	Development of a simple indocyanine green measurement method using an automated biochemical analyser. 2018 , 55, 491-495		
70	Predictors of post-hepatectomy liver failure in patients undergoing extensive liver resections for hepatocellular carcinoma. 2018 , 22, 185-196		11
69	Hepatic venous pressure gradient after portal vein embolization: An accurate predictor of future liver remnant hypertrophy. 2018 , 164, 227-232		6
68	Prognosis of Hepatocellular Carcinoma Patients Who Achieved Long-Term Recurrence-Free Survival After Curative Therapy: Impact of the ALBI Grade. <i>Journal of Gastrointestinal Surgery</i> , 2018 , 22, 1230-1238	3.3	11
67	Liver Resection for Hepatocellular Carcinoma in Patients with Renal Dysfunction. 2018 , 42, 4054-4062		9
66	Validation of Modified ALBI Grade for More Detailed Assessment of Hepatic Function in Hepatocellular Carcinoma Patients: A Multicenter Analysis. <i>Liver Cancer</i> , 2019 , 8, 121-129	9.1	88
65	Different clinical risk scores for prediction of early mortality after liver resection for hepatocellular carcinoma: which is the best?. 2019 , 89, 1275-1280		1
64	A novel magnetic resonance contrast agent for detection of hepatic segmental function. 2019 , 254, 452-456		1
63	Value of contrast-enhanced ultrasound for preoperative assessment of liver reserve function in patients with liver tumors. 2019 , 14, e0222514		1
62	A new formula to calculate the resection limit in hepatectomy based on Gd-EOB-DTPA-enhanced magnetic resonance imaging. 2019 , 14, e0210579		8
61	An elevated neutrophil-to-lymphocyte ratio predicts a poor postoperative survival in primary hepatocellular carcinoma patients with a normal preoperative serum level of alpha-fetoprotein. <i>Surgery Today</i> , 2019 , 49, 661-669	3	5
60	Reliability of indocyanine green retention and clearance rates at 15 minutes calculated by dye-dilution cardiac output flowmetry in comparison to blood sampling in patients undergoing hepatic resection. <i>Indian Journal of Gastroenterology</i> , 2019 , 38, 441-449	1.9	2
59	Assessment of Preoperative Liver Function for Surgical Decision Making in Patients with Hepatocellular Carcinoma. <i>Liver Cancer</i> , 2019 , 8, 447-456	9.1	18
58	Diffusion kurtosis imaging in the assessment of liver function: Its potential as an effective predictor of liver function. <i>British Journal of Radiology</i> , 2019 , 92, 20170608	3.4	3
57	Perioperative and prognostic implication of albumin-bilirubin-TNM score in Child-Pugh class A hepatocellular carcinoma. <i>Annals of Gastroenterological Surgery</i> , 2019 , 3, 65-74	4.3	8
56	Predicting Patients With Insufficient Liver Enhancement in the Hepatobiliary Phase Before the Injection of Gadoteric Acid: A Practical Approach Using the Bayesian Method. <i>Journal of Magnetic Resonance Imaging</i> , 2020 , 51, 62-69	5.6	1
55	Gd-EOB-DTPA-enhanced MRI T1 relaxometry as an imaging-based liver function test compared with C-methacetin breath test. <i>Acta Radiologica</i> , 2020 , 61, 291-301	2	5

54	Zinc deficiency as an independent prognostic factor for patients with early hepatocellular carcinoma due to hepatitis virus. <i>Hepatology Research</i> , 2020 , 50, 92-100	5.1	8
53	Validation set analysis to predict postoperative outcomes by technetium-99m galactosyl serum albumin scintigraphy for hepatectomy. <i>Nuclear Medicine Communications</i> , 2020 , 41, 320-326	1.6	3
52	Is left-sided involvement of hepatocellular carcinoma an important preoperative predictive factor of poor outcome?. <i>World Journal of Surgical Oncology</i> , 2020 , 18, 317	3.4	4
51	Indocyanine Green Clearance Test for the Preoperative Assessment of Chemotherapy-Related Hepatic Injury in Patients with Colorectal Liver Metastasis. <i>Cancer Management and Research</i> , 2020 , 12, 3237-3245	3.6	1
50	Prognostic value of the albumin-bilirubin grade in patients with hepatocellular carcinoma and other liver diseases. <i>Annals of Translational Medicine</i> , 2020 , 8, 553	3.2	15
49	Assessing Liver Function in Liver Tumors Patients: The Performance of T1 Mapping and Residual Liver Volume on Gd-EOBDTPA-Enhanced MRI. <i>Frontiers in Medicine</i> , 2020 , 7, 215	4.9	2
48	Preoperative Portal Vein Embolization in Hepatic Surgery: A Review about the Embolic Materials and Their Effects on Liver Regeneration and Outcome. <i>Radiology Research and Practice</i> , 2020 , 2020, 9295852	2.3	8
47	New technical development for pure laparoscopic donor hepatectomy: indocyanine green cholangiography and three-dimensional laparoscopy. <i>Laparoscopic Surgery</i> , 2020 , 4, 17-17	0.1	
46	Effectiveness comparison of indocyanine green retention test with the cirrhotic severity scoring in evaluating the pathological severity of liver cirrhosis in patients with hepatocellular carcinoma and Child-Pugh grade A liver function. <i>World Journal of Surgical Oncology</i> , 2020 , 18, 79	3.4	6
45	Preoperative Remnant Liver Function Evaluation Using a Routine Clinical Dynamic Gd-EOB-DTPA-Enhanced MRI Protocol in Patients with Hepatocellular Carcinoma. <i>Annals of Surgical Oncology</i> , 2021 , 28, 3672-3682	3.1	4
44	Indocyanine green: An old drug with novel applications. <i>Tzu Chi Medical Journal</i> , 2021 , 33, 317-322	1.1	0
43	Impact of spontaneous splenorenal shunt on liver volume and long-term survival of liver cirrhosis. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2021 , 36, 1694-1702	4	7
42	Prediction of portal vein thrombosis after hepatectomy for hepatocellular carcinoma. <i>Langenbecks Archives of Surgery</i> , 2021 , 406, 781-789	3.4	4
41	Radiomics Analysis of Gd-EOB-DTPA Enhanced Hepatic MRI for Assessment of Functional Liver Reserve. <i>Academic Radiology</i> , 2021 ,	4.3	0
40	Prediction of survival after partial hepatectomy using a physiologically based pharmacokinetic model of indocyanine green liver function tests.		4
39	Improved indocyanine green retention after short-term lenvatinib withdrawal in three patients with hepatocellular carcinoma. <i>Clinical Journal of Gastroenterology</i> , 2021 , 14, 1484-1490	1.1	1
38	The Application of Preoperative Three-Dimensional Reconstruction Visualization Digital Technology in the Surgical Treatment of Hepatic Echinococcosis in Tibet. <i>Frontiers in Surgery</i> , 2021 , 8, 715005	2.3	0
37	Volumetric assessment and clinical predictors of cirrhosis in patients undergoing hepatectomy for hepatocellular carcinoma with presumed normal liver function. <i>Hepatology International</i> , 2021 , 15, 1258-1267	8.8	0

36	Liver Stiffness Measured by Two-Dimensional Shear Wave Elastography for Predicting Symptomatic Post-hepatectomy Liver Failure in Patients with Hepatocellular Carcinoma. <i>Annals of Surgical Oncology</i> , 2021 , 1	3.1	2
35	Functional Assessment of Liver for Radiation Oncologist. 2021 , 51-57		
34	Quantitative Evaluation of Liver Function Within MR Imaging. 2014 , 233-251		3
33	Assessment of liver function reserve by photoacoustic tomography: a feasibility study. <i>Biomedical Optics Express</i> , 2020 , 11, 3985-3995	3.5	4
32	The albumin-bilirubin score stratifies the outcomes of Child-Pugh class A patients after resection of hepatocellular carcinoma.. <i>Translational Cancer Research</i> , 2019 , 8, S233-S244	0.3	2
31	Preoperative predictors of short-term survival after hepatectomy for multinodular hepatocellular carcinoma. <i>World Journal of Gastroenterology</i> , 2012 , 18, 3272-81	5.6	25
30	CT/99mTc-GSA SPECT fusion images demonstrate functional differences between the liver lobes. <i>World Journal of Gastroenterology</i> , 2013 , 19, 3217-25	5.6	13
29	Use of magnetic resonance elastography for assessing liver functional reserve: A clinical study. <i>World Journal of Gastroenterology</i> , 2015 , 21, 7522-8	5.6	5
28	Evaluation of safety for hepatectomy in a novel mouse model with nonalcoholic-steatohepatitis. <i>World Journal of Gastroenterology</i> , 2018 , 24, 1622-1631	5.6	4
27	Preoperative imaging for hepatic resection of colorectal cancer metastasis. <i>Journal of Gastrointestinal Oncology</i> , 2012 , 3, 11-8	2.8	38
26	Small for size liver remnant following resection: prevention and management. <i>Hepatobiliary Surgery and Nutrition</i> , 2014 , 3, 303-12	2.1	16
25	Brief communication (Original). Correlation between liver signal intensity in hepatobiliary phase of Gd-EOB-DTPA enhancement and liver function reserves. <i>Asian Biomedicine</i> , 2014 , 8, 411-415	0.4	0
24	Operative Microwave Ablation for Hepatocellular Carcinoma Within 3 cm and 3 Nodules: Experience in 559 Patients. <i>Journal of Gastrointestinal Surgery</i> , 2021 , 1	3.3	1
23	Leber. 2012 , 95-105		
22	Anästhesiologische Beurteilung des Patienten: Leber. 2016 , 1-14		
21	Preoperative Preparations for Patients with Hepatocellular Carcinoma. 2019 , 7-26		
20	Other Types of Hepatic Resection for HCC. 2019 , 261-329		
19	Development of a novel rat model of heterogeneous hepatic injury by injection with colchicine the splenic vein. <i>World Journal of Gastroenterology</i> , 2018 , 24, 5005-5012	5.6	1

18	Anästhesiologische Beurteilung des Patienten: Leber. <i>Springer Reference Medizin</i> , 2019 , 117-129	0	
17	Significance of indocyanine green test in radiotherapy for hepatocellular carcinoma.. <i>Translational Cancer Research</i> , 2019 , 8, 14-16	0.3	
16	Quantitative Evaluation of Liver Function with Gd-EOB-DTPA-enhanced MR Imaging. <i>Japanese Journal of Magnetic Resonance in Medicine</i> , 2019 , 39, 137-144	0	
15	Evaluating the liver function of patients with hepatocellular carcinoma. <i>Hepatology Research</i> , 2021 , 51, 1095-1096	5.1	0
14	To determine the prognostic value of the albumin-bilirubin grade (ALBI) in patients underwent transarterial chemoembolization for unresectable hepatocellular carcinoma. <i>Gastroenterology and Hepatology From Bed To Bench</i> , 2019 , 12, 110-115	1.2	3
13	Prediction of Survival After Partial Hepatectomy Using a Physiologically Based Pharmacokinetic Model of Indocyanine Green Liver Function Tests. <i>Frontiers in Physiology</i> , 2021 , 12, 730418	4.6	3
12	Combination of portal vein embolization and neoadjuvant chemotherapy for locally advanced gallbladder cancer requiring extended hepatectomy - A novel approach.. <i>Indian Journal of Gastroenterology</i> , 2021 , 40, 580	1.9	0
11	Virtual hepatectomy and metastasectomy 3d simulation using an automated and semiautomated tool applied to preoperative CT in the treatment of colorectal cancer metastases. <i>BJR case Reports</i> , 2022 , 7, 20210117	0.7	
10	Liver Function and Posthepatectomy Liver Failure. 2022 , 23-30		
9	Comparative study of indocyanine green (ICG)-R15 and Albumin-Indocyanine Green Evaluation (ALICE) grading system in the prediction of posthepatectomy liver failure and postoperative mortality in patients with hepatic alveolar echinococcosis. <i>BMC Gastroenterology</i> , 2022 , 22, 1-10	3	
8	The Effect of a Liver Transplant Program on the Outcomes of Resectable Hepatocellular Carcinoma: A Nationwide Multicenter Analysis. <i>Annals of Surgery</i> , Publish Ahead of Print, 2022	7.8	0
7	The Liver SEntinel LYmph-node (LISELY) study: A prospective intraoperative real time evaluation of liver lymphatic drainage and sentinel lymph-node using near-infrared (NIR) imaging with Indocyanine Green (ICG). <i>European Journal of Surgical Oncology</i> , 2022 , 48, 1056-1062	3.6	
6	Impact of comorbid renal dysfunction in patients with hepatocellular carcinoma on long-term outcomes after curative resection. <i>World Journal of Gastrointestinal Surgery</i> , 2022 , 14, 670-684	2.4	0
5	Optimal tumor numbers in surgical candidates for multiple hepatocellular carcinomas. 2022 , 1-10		0
4	Quantitative evaluation of functional liver reserve using the liver-to-spleen R1 ratio based on gadoxetic acid-enhanced MRI: corrected for the effect of gadoxetic acid dose. 028418512211376		0
3	Machine learning-based Radiomics analysis of preoperative functional liver reserve with MRI and CT image.		0
2	On the relationship between metabolic capacities and in vivo viscoelastic properties of the liver. 10, 1-10		0
1	Indocyanine Green Fluorescence for Liver Assessment and Imaging-Guided Resection of Colorectal Metastases: A Case Report. 2023 , 50, 54-58		0

