Sildenafil and glyceryl trinitrate reduce tactile allodynia in streptozotocin-injected rats

DOI: 10.1016/j.ejphar.2010.01.001
European Journal of Pharmacology, 2010, 631, 17-23.

#	Paper	IF	Citations
6	Role of hydrogen sulfide in the pain processing of non-diabetic and diabetic rats. *Neuroscience*, **2013**, 250, 786-97	3.9	15
5	Angiotensin (5-8) modulates nociception at the rat periaqueductal gray via the NO-sGC pathway and an endogenous opioid. *Neuroscience*, **2013**, 231, 315-27	3.9	7
4	Drug repositioning: playing dirty to kill pain. *CNS Drugs*, **2014**, 28, 45-61	6.7	27
3	Painful neuropathy: Mechanisms. *Handbook of Clinical Neurology / Edited By P J Vinken and G W Bruyn*, **2014**, 126, 533-57	3	17
2	Fosinopril Prevents the Development of Tactile Allodynia in a Streptozotocin-Induced Diabetic Rat Model. *Drug Development Research*, **2015**, 76, 442-9	5.1	13
1	Drug Repurposing for the Development of Novel Analgesics. *Trends in Pharmacological Sciences*, **2016**, 37, 172-183	13.2	37