## CITATION REPORT List of articles citing

Nine-year incident diabetes is predicted by fatty liver indices: the French D.E.S.I.R. study

DOI: 10.1186/1471-230x-10-56 BMC Gastroenterology, 2010, 10, 56.

Source: https://exaly.com/paper-pdf/48386312/citation-report.pdf

Version: 2024-04-16

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
108	A simple index of lipid overaccumulation is a good marker of liver steatosis. <i>BMC Gastroenterology</i> , <b>2010</b> , 10, 98	3	140
107	C-reactive protein and gamma-glutamyltransferase concentrations in relation to the prevalence of type 2 diabetes diagnosed by glucose or HbA1c criteria in Chinese adults in Qingdao, China. <b>2010</b> , 2010, 761715		2
106	The Finnish Diabetes Risk Score (FINDRISC) and other non-invasive scores for screening of hepatic steatosis and associated cardiometabolic risk. <b>2011</b> , 43, 413-7		11
105	Association between markers of fatty liver disease and impaired glucose regulation in men and women from the general population: the KORA-F4-study. <b>2011</b> , 6, e22932		46
104	Systematic review: the diagnosis and staging of non-alcoholic fatty liver disease and non-alcoholic steatohepatitis. <b>2011</b> , 33, 525-40		215
103	Why does NAFLD predict type 2 diabetes?. <b>2011</b> , 11, 167-72		26
102	Elevated liver function enzymes are related to the development of prediabetes and type 2 diabetes in younger adults: the Bogalusa Heart Study. <b>2011</b> , 34, 2603-7		60
101	Combined influence of insulin resistance, overweight/obesity, and fatty liver as risk factors for type 2 diabetes. <b>2012</b> , 35, 717-22		145
100	Long-term dietary potential renal acid load during adolescence is prospectively associated with indices of nonalcoholic fatty liver disease in young women. <b>2012</b> , 142, 313-9		20
99	Associations of sex hormone-binding globulin and testosterone with diabetes among men and women (the Saku Diabetes study): a case control study. <b>2012</b> , 11, 130		25
98	Predicting total, abdominal, visceral and hepatic adiposity with circulating biomarkers in Caucasian and Japanese American women. <b>2012</b> , 7, e43502		41
97	Liver and diabetes. A vicious circle. <b>2013</b> , 43, 51-64		130
96	Hypertension and vascular dynamics in men and women with metabolic syndrome. <b>2013</b> , 61, 12-9		84
95	The Fatty Liver Index has limited utility for the detection and quantification of hepatic steatosis in obese patients. <b>2013</b> , 7, 592-9		15
94	Early growth and non-alcoholic fatty liver disease in adulthood-the NAFLD liver fat score and equation applied on the Helsinki Birth Cohort Study. <b>2013</b> , 45, 430-7		90
93	Assessment of cardiometabolic risk and prevalence of meeting treatment guidelines among patients with type 2 diabetes stratified according to their use of insulin and/or other diabetic medications: results from INSPIRE ME IAA. <b>2013</b> , 15, 629-41		4
92	Non-alcoholic fatty liver disease independently predicts prediabetes during a 7-year prospective follow-up. <b>2013</b> , 33, 1406-12		60

91	Influence of non-alcoholic fatty liver disease on the development of diabetes mellitus. 2013, 28, 142-7	66
90	External validation of the fatty liver index for identifying nonalcoholic fatty liver disease in a population-based study. <b>2013</b> , 11, 1201-4	128
89	Review article: the diagnosis of non-alcoholic fatty liver disease availability and accuracy of non-invasive methods. <b>2013</b> , 37, 392-400	126
88	Assessment of the fatty liver index as an indicator of hepatic steatosis for predicting incident diabetes independently of insulin resistance in a Korean population. <b>2013</b> , 30, 428-35	30
87	The association between nonalcoholic fatty pancreas disease and diabetes. 2013, 8, e62561	95
86	Association of serum adipocyte-specific Fatty Acid binding protein with Fatty liver index as a predictive indicator of nonalcoholic Fatty liver disease. <b>2013</b> , 28, 283-7	11
85	The role of skeletal muscle in development of nonalcoholic Fatty liver disease. <b>2013</b> , 37, 278-85	48
84	Comparison of fatty liver index with noninvasive methods for steatosis detection and quantification. <b>2013</b> , 19, 57-64	77
83	Comparison of liver fat indices for the diagnosis of hepatic steatosis and insulin resistance. <b>2014</b> , 9, e94059	88
82	The effects of vitamin D supplementation on hepatic dysfunction, vitamin D status, and glycemic control in children and adolescents with vitamin D deficiency and either type 1 or type 2 diabetes mellitus. <b>2014</b> , 9, e99646	40
81	Non-alcoholic fatty liver disease: what the clinician needs to know. <b>2014</b> , 20, 12956-80	117
80	Performance and limitations of steatosis biomarkers in patients with nonalcoholic fatty liver disease. <b>2014</b> , 40, 1209-22	196
79	Fatty liver and serum cholinesterase are independently correlated with HbA1c levels: cross-sectional analysis of 5384 people. <b>2014</b> , 42, 542-53	7
78	Elevated liver enzymes are related to progression to impaired glucose tolerance in Japanese men. <b>2014</b> , 31, 552-8	7
77	External validation of the fatty liver index and lipid accumulation product indices, using 1H-magnetic resonance spectroscopy, to identify hepatic steatosis in healthy controls and obese, insulin-resistant individuals. <b>2014</b> , 171, 561-9	80
76	[Non-invasive markers of fibrosis in the diagnosis of non-alcoholic fatty liver disease]. <b>2014</b> , 37, 503-10	5
75	Physical activity, sedentary time, and liver enzymes in adolescents: the HELENA study. <b>2014</b> , 75, 798-802	18
74	Liver in the analysis of body composition by dual-energy X-ray absorptiometry. <b>2014</b> , 87, 20140232	6

73	Evaluation of the fatty liver index as a predictor for the development of diabetes among insurance beneficiaries with prediabetes. <b>2015</b> , 6, 309-16		16
72	Sex-Specific Predictors of Metabolic Syndrome Independent of Its Components. <b>2015</b> , 63, 796-801		6
71	Serum cytokeratin-18 fragment levels predict development of type 2 diabetes mellitus in adult patients with NAFLD. <b>2015</b> , 35, 2621		5
70	Non-invasive diagnostic approach to non-alcoholic fatty liver disease: current evidence and future perspectives. <b>2015</b> , 9, 1039-53		9
69	Country of birth modifies the association of fatty liver index with insulin action in Middle Eastern immigrants to Sweden. <b>2015</b> , 110, 66-74		6
68	NAFLD: a multisystem disease. <b>2015</b> , 62, S47-64		1286
67	Identifying patients with chronic hepatitis B at high risk of type 2 diabetes mellitus: a cross-sectional study with pair-matched controls. <i>BMC Gastroenterology</i> , <b>2015</b> , 15, 32	3	16
66	A Perspective on Metabolic Syndrome and Nonalcoholic Fatty Liver Disease. <b>2015</b> , 13, 235-8		21
65	Evaluation of various biomarkers as potential mediators of the association between B desaturase, B desaturase, and stearoyl-CoA desaturase activity and incident type 2 diabetes in the European Prospective Investigation into Cancer and Nutrition-Potsdam Study. <b>2015</b> , 102, 155-64		38
64	Prediabetes and associated disorders. <b>2015</b> , 48, 371-93		86
63	Bidirectional Relationships and Disconnects between NAFLD and Features of the Metabolic Syndrome. <b>2016</b> , 17, 367		60
62	Discriminatory value of alanine aminotransferase for diabetes prediction: the Insulin Resistance Atherosclerosis Study. <b>2016</b> , 33, 348-55		7
61	Nonalcoholic fatty liver disease is associated with an almost twofold increased risk of incident type 2 diabetes and metabolic syndrome. Evidence from a systematic review and meta-analysis. <b>2016</b> , 31, 936-44		378
60	How to best diagnose NAFLD/NASH?. <b>2016</b> , 111-123		
59	Fatty liver is an independent predictor of early carotid atherosclerosis. <b>2016</b> , 65, 95-102		67
58	Adipose tissue dysfunction increases fatty liver association with pre diabetes and newly diagnosed type 2 diabetes mellitus. <i>Diabetology and Metabolic Syndrome</i> , <b>2016</b> , 8, 73	5.6	9
57	Association between liver enzymes and incident type 2 diabetes in Singapore Chinese men and women. <b>2016</b> , 4, e000296		30
56	Fatty liver index as a simple predictor of incident diabetes from the KoGES-ARIRANG study. <b>2016</b> , 95, e4447		27

## (2019-2016)

55	Fatty liver index is a risk determinant of incident type 2 diabetes in a metabolically healthy population with obesity. <b>2016</b> , 24, 1373-9	26
54	Non-alcoholic fatty liver disease and diabetes. <b>2016</b> , 65, 1096-108	266
53	Low alcohol consumption increases the risk of impaired glucose tolerance in patients with non-alcoholic fatty liver disease. <b>2016</b> , 51, 1090-1100	10
52	Analysis of the Japanese Diabetes Risk Score and fatty liver markers for incident diabetes in a Japanese cohort. <b>2016</b> , 10, 19-26	3
51	T-cadherin gene variants are associated with type 2 diabetes and the Fatty Liver Index in the French population. <b>2017</b> , 43, 33-39	13
50	Associations of the fatty liver and hepatic steatosis indices with risk of cardiovascular disease: Interrelationship with age. <b>2017</b> , 466, 54-60	17
49	Gamma-glutamyltransferase, fatty liver index and hepatic insulin resistance are associated with incident hypertension in two longitudinal studies. <b>2017</b> , 35, 493-500	40
48	The Fatty Liver Index (FLI) Relates to Diabetes-Specific Parameters and an Adverse Lipid Profile in a Cohort of Nondiabetic, Dyslipidemic Patients. <b>2017</b> , 36, 287-294	6
47	Non-alcoholic fatty liver disease and its relationship with cardiovascular disease and other extrahepatic diseases. <b>2017</b> , 66, 1138-1153	508
46	Esteatosis hep <b>l</b> ica: diagn¶tico y seguimiento. <b>2017</b> , 24, 378-389	O
45	Diable, facteur daggravation des hatopathies. <b>2017</b> , 11, 687-691	1
44	Fatty liver index predicts incident diabetes in a Japanese general population with and without impaired fasting glucose. <b>2018</b> , 48, 708-716	9
43	Associations between dietary factors and markers of NAFLD in a general Dutch adult population. <b>2018</b> , 72, 117-123	36
42	NAFLD/NASH in patients with type 2 diabetes and related treatment options. <b>2018</b> , 41, 509-521	32
41	Hepatic IRS1 and Ecatenin expression is associated with histological progression and overt diabetes emergence in NAFLD patients. <b>2018</b> , 53, 1261-1275	16
40	Fatty liver index is a predictor of incident diabetes in patients with prediabetes: The PREDAPS study. <b>2018</b> , 13, e0198327	23
39	Association of fatty liver index with risk of incident type 2 diabetes by metabolic syndrome status	7
	in an Eastern Finland male cohort: a prospective study. <b>2019</b> , 9, e026949	

37	Fatty liver index is a strong predictor of changes in glycemic status in people with prediabetes: The IT-DIAB study. <b>2019</b> , 14, e0221524	16
36	The association between plasma proneurotensin and glucose regulation is modified by country of birth. <b>2019</b> , 9, 13640	6
35	Nonalcoholic Fatty Liver Disease. <b>2019</b> , 523-546	
34	The fatty liver index, a simple and useful predictor of metabolic syndrome: analysis of the Korea National Health and Nutrition Examination Survey 2010-2011. <b>2019</b> , 12, 181-190	19
33	Liver fibrosis by FibroScan independently of established cardiovascular risk parameters associates with macrovascular and microvascular complications in patients with type 2 diabetes. <b>2020</b> , 40, 347-354	27
32	Relationships between plasma lactate, plasma alanine, genetic variations in lactate transporters and type 2 diabetes in the Japanese population. <b>2020</b> , 35, 131-138	4
31	Metabolic Footprint, towards Understanding Type 2 Diabetes beyond Glycemia. <i>Journal of Clinical Medicine</i> , <b>2020</b> , 9,	4
30	Identifying High-Risk NASH Patients: What We Know so Far. <b>2020</b> , 12, 125-138	11
29	Prevalence of elevated liver enzymes and its association with type 2 diabetes: A cross-sectional study in Bangladeshi adults. <b>2020</b> , 3, e00116	14
28	Nonalcoholic fatty liver disease and the risk of atrial fibrillation stratified by body mass index: a nationwide population-based study. <b>2021</b> , 11, 3737	7
27	SHBG as a Marker of NAFLD and Metabolic Impairments in Women Referred for Oligomenorrhea and/or Hirsutism and in Women With Sexual Dysfunction. <b>2021</b> , 12, 641446	5
26	Low handgrip strength is associated with higher liver enzyme concentrations in US adolescents. <b>2021</b> ,	
25	High level of fatty liver index predicts new onset of diabetes mellitus during a 10-year period in healthy subjects. <b>2021</b> , 11, 12830	2
24	Fatty liver index as a predictor for type 2 diabetes in subjects with normoglycemia in a nationwide cohort study. <b>2021</b> , 11, 16453	O
23	Fatty liver index as a predictive marker for the development of diabetes: A retrospective cohort study using Japanese health check-up data. <b>2021</b> , 16, e0257352	1
22	Fatty liver index predicts further metabolic deteriorations in women with previous gestational diabetes. <b>2012</b> , 7, e32710	40
21	Associations between longer habitual day napping and non-alcoholic fatty liver disease in an elderly Chinese population. <b>2014</b> , 9, e105583	15
20	Association between the Fatty Liver Index and Risk of Type 2 Diabetes in the EPIC-Potsdam Study. <b>2015</b> , 10, e0124749	38

19	Epidemiology of fatty liver: an update. <b>2014</b> , 20, 9050-4		59
18	Fatty liver disease: Disparate predictive ability for cardiometabolic risk and all-cause mortality. <b>2015</b> , 21, 13555-65		7
17	Prediction of NAFLD occurrence in prediabetes patients. <b>2020</b> , 20, 190		8
16	Nonalcoholic fatty liver disease: Synopsis of current developments. <b>2015</b> , 18, 703-12		22
15	Neck Circumference to Height Ratio is a Reliable Predictor of Liver Stiffness and Nonalcoholic Fatty Liver Disease in Prediabetes. <b>2018</b> , 22, 347-354		8
14	Vitamin D and liver enzymes' levels in Libyans with type 2 diabetes. <b>2021</b> , 5, 116		2
13	Nonalcoholic Fatty Liver Disease. <b>2022</b> , 295-323		
12	Alcohol and Nutrition as Risk Factors for Chronic Liver Disease. <b>2013</b> , 497-506		
11	Diabetes and NAFLD: Why is the Connection Important?. 62-70		
10	A STUDY OF NONALCOHOLIC FATTY LIVER DISEASE AND FATTY LIVER INDEX IN TYPE 2 DIABETES MELLITUS PATIENTS. <b>2016</b> , 3, 3001-3006		
9	Increased Serum Pentraxin 3 Is Associated with Prediabetes and Type 2 Diabetes in Obese Patients with Nonalcoholic Fatty Liver Disease. <b>2021</b> ,		0
8	Nonalcoholic Fatty Liver Disease Is a Precursor of New-Onset Metabolic Syndrome in Metabolically Healthy Young Adults <i>Journal of Clinical Medicine</i> , <b>2022</b> , 11,	5.1	
7	Validation of fatty liver index as a marker for metabolic dysfunction-associated fatty liver disease <i>Diabetology and Metabolic Syndrome</i> , <b>2022</b> , 14, 44	5.6	О
6	Insights into Non-Alcoholic Fatty Liver Disease and Non-Alcoholic Steatohepatitis. <i>Medicina Interna</i> (Bucharest, Romania: 1991), <b>2022</b> , 19, 61-77	0.1	
5	Sex and age differences in the association of fatty liver index-defined non-alcoholic fatty liver disease with cardiometabolic risk factors: a cross-sectional study. <b>2022</b> , 13,		1
4	Fatty liver index predicts the development of hypertension in a Japanese general population with and without dysglycemia.		О
3	Incidence of diabetes mellitus according to body mass index and gamma-glutamyltransferase level in Korea. <b>2022</b> , 4,		О
2	Fatty Liver Disease in Patients with Prediabetes and Overweight or Obesity. <b>2023</b> , 13, 531		О

The association between fatty liver index and onset of diabetes: secondary analysis of a population-based cohort study. **2023**, 23,

О