## Dust flux estimates for the Last Glacial Period in East Corecords of loess deposits: a review

Quaternary Science Reviews 29, 3157-3166 DOI: 10.1016/j.quascirev.2010.07.005

Citation Report

#	Article	IF	CITATIONS
1	Investigating the penultimate and last glacial cycles of the Süttő loess section (Hungary) using luminescence dating, high-resolution grain size, and magnetic susceptibility data. Quaternary International, 2011, 234, 75-85.	1.5	107
2	Tectonic versus climatic control on the evolution of a loess–paleosol sequence at Beremend, Hungary: an integrated approach based on paleoecological, clay mineralogical, and geochemical data. Quaternary International, 2011, 240, 71-86.	1.5	54
3	The Belotinac section (Southern Serbia) at the southern limit of the European loess belt: Initial results. Quaternary International, 2011, 240, 128-138.	1.5	22
4	Dust deposition and climate in the Carpathian Basin over an independently dated last glacial–interglacial cycle. Quaternary Science Reviews, 2011, 30, 662-681.	3.0	214
5	The last million years recorded at the Stari Slankamen (Northern Serbia) loess-palaeosol sequence: revised chronostratigraphy and long-term environmental trends. Quaternary Science Reviews, 2011, 30, 1142-1154.	3.0	169
6	Loess towards (geo) tourism – proposed application on loess in Vojvodina region (north Serbia). Acta Geographica Slovenica, 2011, 51, 391-406.	0.7	34
7	Radiocarbon chronology of Late Pleistocene large mammal faunas from the Pannonian basin (Hungary) Bulletin of Geosciences, 2012, , 13-19.	1.1	10
8	Establishing a luminescence chronology for a palaeosol-loess profile at Tokaj (Hungary): A comparison of quartz OSL and polymineral IRSL signals. Quaternary Geochronology, 2012, 10, 68-74.	1.4	44
9	Pleistocene environmental dynamics recorded in the loess of the middle and lower Danube basin. Quaternary Science Reviews, 2012, 41, 104-118.	3.0	157
10	Evaluating the use of clay mineralogy, Sr–Nd isotopes and zircon U–Pb ages in tracking dust provenance: An example from loess of the Carpathian Basin. Chemical Geology, 2012, 304-305, 83-96.	3.3	78
11	Upbuilding pedogenesis under active loess deposition in a super-humid, temperate climate — quantification of deposition rates, soil chemistry and pedogenic thresholds. Geoderma, 2012, 189-190, 491-501.	5.1	29
12	PALEOSOLS AND WIND-BLOWN SEDIMENTS   Biogeochemical Role of Dust in Quaternary Climate Cycles. , 2013, , 412-420.		0
13	Humid glacials, arid interglacials? Critical thoughts on pedogenesis and paleoclimate based on multi-proxy analyses of the loess–paleosol sequence Crvenka, Northern Serbia. Palaeogeography, Palaeoclimatology, Palaeoecology, 2013, 387, 165-175.	2.3	102
14	Analysis of Saharan dust intrusions into the Carpathian Basin (Central Europe) over the period of 1979–2011. Global and Planetary Change, 2013, 100, 333-342.	3.5	49
15	Sedimentary characteristics and source of loess in Baranja (Eastern Croatia). Aeolian Research, 2013, 11, 129-139.	2.7	21
16	TowardsÂidentifying the origin of metamorphic components in Austrian loess: insights from detrital rutile chemistry, thermometry and U–Pb geochronology. Quaternary Science Reviews, 2013, 75, 132-142.	3.0	29
17	The chronology of the Åarengrad II loess-palaeosol section (Eastern Croatia). Geologia Croatica, 2013, 66, 191-203.	0.8	25
18	Late Pleistocene variations of the background aeolian dust concentration in the Carpathian Basin: an estimate using decomposition of grain-size distribution curves of loess deposits. Geologie En Mijnbouw/Netherlands Journal of Geosciences, 2013, 91, 159-171	0.9	15

CITATION REPORT

#	Article	IF	CITATIONS
19	Loess in the Vojvodina region (Northern Serbia): an essential link between European and Asian Pleistocene environments. Geologie En Mijnbouw/Netherlands Journal of Geosciences, 2013, 91, 173-188.	0.9	57
20	Eolian contribution to geochemical and mineralogical characteristics of some soil types in Medvednica Mountain, Croatia. Catena, 2014, 117, 145-156.	5.0	8
21	Loess accumulation during the last glacial maximum: Evidence from Urluia, southeastern Romania. Quaternary International, 2014, 334-335, 74-85.	1.5	39
22	Vegetation and environmental responses to climate forcing during the Last Glacial Maximum and deglaciation in the East Carpathians: attenuated response to maximum cooling and increased biomass burning. Quaternary Science Reviews, 2014, 106, 278-298.	3.0	65
23	Introducing an improved multi-proxy approach for paleoenvironmental reconstruction of loess–paleosol archives applied on the Late Pleistocene Nussloch sequence (SW Germany). Palaeogeography, Palaeoclimatology, Palaeoecology, 2014, 410, 300-315.	2.3	53
24	The Late Pleistocene Belotinac section (southern Serbia) at the southern limit of the European loess belt: Environmental and climate reconstruction using grain size and stable C and N isotopes. Quaternary International, 2014, 334-335, 10-19.	1.5	50
25	The conceptual sedimentary model of the Lower Danube loess basin: Sedimentogenetic implications. Quaternary International, 2014, 351, 14-24.	1.5	28
26	Tracing the origin of loess in Hungary with the help of heavy mineral composition data. Quaternary International, 2014, 319, 11-21.	1.5	25
27	Two possible source regions for central Greenland last glacial dust. Geophysical Research Letters, 2015, 42, 10,399.	4.0	39
28	Aeolian dynamics at the Orlovat loess–paleosol sequence, northern Serbia, based on detailed textural and geochemical evidence. Aeolian Research, 2015, 18, 69-81.	2.7	56
29	Core-Shell Processing of Natural Pigment: Upper Palaeolithic Red Ochre from Lovas, Hungary. PLoS ONE, 2015, 10, e0131762.	2.5	19
30	Results of paleoecological studies in the loess region of Szeged-Öthalom (SE Hungary). Quaternary International, 2015, 372, 66-78.	1.5	30
31	Danube loess stratigraphy — Towards a pan-European loess stratigraphic model. Earth-Science Reviews, 2015, 148, 228-258.	9.1	241
32	Climate vs. parent material — Which is the key of Stagnosol diversity in Croatia?. Geoderma, 2015, 241-242, 250-261.	5.1	14
33	Compositional trends through the Holocene mud succession of the southwestern Black Sea shelf: Implications for sedimentary provenance and water-level history. Sedimentary Geology, 2015, 316, 13-25.	2.1	15
34	Fine and coarse-quartz SAR-OSL dating of Last Glacial loess in Southern Romania. Quaternary International, 2015, 357, 33-43.	1.5	27
35	Mineralogical discrimination of the pleistocene loess/paleosol sections in Srijem and Baranja, Croatia. Aeolian Research, 2016, 21, 151-162.	2.7	6
36	High-resolution trace element and stable/radiogenic isotope profiles of late Pleistocene to Holocene speleothems from Dim Cave, SW Turkey. Palaeogeography, Palaeoclimatology, Palaeoecology, 2016, 452, 68-79.	2.3	19

ARTICLE IF CITATIONS # Saharan dust deposition in the Carpathian Basin and its possible effects on interglacial soil 37 2.7 46 formation. Aeolian Research, 2016, 22, 1-12. Sedimentological and mineralogical characteristics of the Pleistocene loess/paleosol sections in the 2.7 Eastern Croatia. Aeolian Research, 2016, 20, 7-23. 39 The history of Danube loess research. Quaternary International, 2016, 399, 86-99. 1.5 27 Coupled European and Greenland last glacial dust activity driven by North Atlantic climate. Proceedings of the National Academy of Sciences of the United States of America, 2017, 114, E10632-E10638. Reconstructing palaeoenvironments on desert margins: New perspectives from Eurasian loess and 41 3.0 26 Australian dry lake shorelines. Quaternary Science Reviews, 2017, 171, 1-19. Loess correlations  $\hat{a} \in \mathcal{E}$  Between myth and reality. Palaeogeography, Palaeoclimatology, Palaeoecology, 2.3 2018, 509, 4-23. The Crvenka loess-paleosol sequence: A record of continuous grassland domination in the southern Carpathian Basin during the Late Pleistocene. Palaeogeography, Palaeoclimatology, Palaeoecology, 43 2.3 38 2018, 509, 33-46. Early Upper Paleolithic surface collections from loess-like sediments in the northern Carpathian 44 1.5 9 Basín. Quaternary International, 2018, 485, 167-182. Discriminating luminescence age uncertainty composition for a robust Bayesian modelling. 45 1.4 26 Quaternary Geochronology, 2018, 43, 30-39. The Last Glacial aeolian record of the Island of Susak (Croatia) as seen from a high-resolution 1.5 grain–size and rock magnetic analysis. Quaternary International, 2018, 494, 211-224. New chronology of the best developed loess/paleosol sequence of Hungary capturing the past 1.1 ma: Implications for correlation and proposed pan-Eurasian stratigraphic schemes. Quaternary Science 47 3.030 Reviews, 2018, 191, 144-166. Approaches and challenges to the study of loess—Introduction to the LoessFest Special Issue. 48 Quaternary Research, 2018, 89, 563-618. Editorial: Eurasian loess records. Palaeogeography, Palaeoclimatology, Palaeoecology, 2018, 509, 1-3. 49 2.31 Quartz OSL dating of late quaternary Chinese and Serbian loess: A cross Eurasian comparison of dust 1.5 44 mass accumulation rates. Quaternary International, 2019, 502, 30-44. Radiocarbon dated malacological records of two Late Pleistocene loess-paleosol sequences from 52 7 1.5 SW-Hungary: Paleoecological inferences. Quaternary International, 2019, 504, 108-117. High-resolution proxy record of the environmental response to climatic variations during transition MIS3/MIS2 and MIS2 in Central Europe: The loess-paleosol sequence of KatymÃ;r brickyard (Hungary). Quaternary International, 2019, 504, 40-55. High-resolution paleoclimatic proxy data from the MIS3/2 transition recorded in northeastern 54 1.521 Hungarian loess. Quaternary International, 2019, 502, 95-107. Holocene sedimentation in the southwestern Black Sea: Interplay between riverine supply, coastal eddies of the Rim Current, surface and internal waves, and saline underflow through the Strait of 2.1 Bosphorus. Marine Geology, 2020, 420, 106092.

CITATION REPORT

CITATION REPORT

#	Article	IF	CITATIONS
56	Initial quartz OSL and dust mass accumulation rate investigation of the Kisiljevo loess sequence in north-eastern Serbia. Quaternary International, 2020, , .	1.5	5
57	A 180 ka record of environmental change at Erdut (Croatia): a new chronology for the loess–palaeosol sequence and its implications for environmental interpretation. Journal of Quaternary Science, 2020, 35, 582-593.	2.1	16
58	A postâ€ <scp>IR IRSL</scp> chronology and dust mass accumulation rates of the Nosak loessâ€palaeosol sequence in northeastern Serbia. Boreas, 2020, 49, 841-857.	2.4	16
59	Abrupt last glacial dust fall over southeast England associated with dynamics of the British-Irish ice sheet. Quaternary Science Reviews, 2020, 250, 106641.	3.0	26
60	Reâ€analysis of late Quaternary dust mass accumulation rates in Serbia using new luminescence chronology for loess–palaeosol sequence at Surduk. Boreas, 2020, 49, 634-652.	2.4	14
61	Plio-Pleistocene Dust Traps on Paleokarst Surfaces: A Case Study From the Carpathian Basin. Frontiers in Earth Science, 2020, 8, .	1.8	3
62	Linkage between dust cycle and loess of the Last Glacial Maximum in Europe. Atmospheric Chemistry and Physics, 2020, 20, 4969-4986.	4.9	35
63	<sup>14</sup> C Dated Chronology of the Thickest and Best Resolved Loess/Paleosol Record of the LGM from SE Hungary Based on Comparing Precision and Accuracy of Age-Depth Models. Radiocarbon, 2020, 62, 403-417.	1.8	11
64	Simulated regional dust cycle in the Carpathian Basin and the Adriatic Sea region during the Last Glacial Maximum. Quaternary International, 2021, 581-582, 114-127.	1.5	17
65	Chronological Assessment of the Balta Alba Kurgan Loess-Paleosol Section (Romania) – A Comparative Study on Different Dating Methods for a Robust and Precise Age Model. Frontiers in Earth Science, 2021, 8, .	1.8	16
66	Loess landscapes of Europe – Mapping, geomorphology, and zonal differentiation. Earth-Science Reviews, 2021, 215, 103496.	9.1	104
67	Palaeoecological background of the Upper Palaeolithic site of Ságvár, Hungary: radiocarbonâ€dated malacological and sedimentological studies on the Late Pleistocene environment. Journal of Quaternary Science, 0, , .	2.1	3
68	Selected Grain-Size and Geochemical Analyses of the Loess-Paleosol Sequence of Pécel (Northern) Tj ETQq0 0 0 Quaternary, 2021, 4, 17.	rgBT /C 2.0	Overlock 10 Ti 4
69	Spatial differences in the luminescence sensitivity of quartz extracted from Carpathian Basin fluvial sediments. Quaternary Geochronology, 2021, 64, 101166.	1.4	10
70	Chronology and dust mass accumulation history of the Wenchuan loess on eastern Tibetan Plateau since the last glacial. Aeolian Research, 2021, 53, 100748.	2.7	8
71	Dunaszekcső Castle Hill: Landslides Along the Danubian Loess Bluff. World Geomorphological Landscapes, 2015, , 113-120.	0.3	7
72	New chronology and extended palaeoenvironmental data to the 1975 loess profile of Madaras brickyard, South Hungary. Journal of Quaternary Science, 2021, 36, 1364-1381.	2.1	3
73	A tale of two signals: Global and local influences on the Late Pleistocene loess sequences in Bulgarian Lower Danube. Quaternary Science Reviews, 2021, 274, 107264.	3.0	16

**CITATION REPORT** 

#	Article	IF	CITATIONS
74	Detailed Luminescence Dating of Dust Mass Accumulation Rates Over the Last Two Glacial-Interglacial Cycles from the Irig Loess-Palaeosol Sequence, Carpathian Basin. SSRN Electronic Journal, 0, , .	0.4	0
75	A remarkable last glacial loess sedimentation at Roxolany in the Dniester Liman (Southern Ukraine). Quaternary Science Reviews, 2022, 285, 107521.	3.0	2
76	Modeling dust emission in alpine regions with low air temperature and low air pressure – A case study on the Qinghai-Tibetan Plateau (QTP). Geoderma, 2022, 422, 115930.	5.1	6
77	Detailed luminescence dating of dust mass accumulation rates over the last two glacial-interglacial cycles from the Irig loess-palaeosol sequence, Carpathian Basin. Global and Planetary Change, 2022, 215, 103895.	3.5	5
78	Greenland Ice Core Record of Last Glacial Dust Sources and Atmospheric Circulation. Journal of Geophysical Research D: Atmospheres, 2022, 127, .	3.3	17
79	Southeast Asian dust flux reconstructed using accurately dated stalagmite thorium concentrations. Results in Geochemistry, 2022, 7-8, 100017.	0.8	0
80	Last glacial loess dynamics in the Southern Caucasus (NE-Armenia) and the phenomenon of missing loess deposition during MIS-2. Scientific Reports, 2022, 12, .	3.3	4
81	Dataset of dust mass accumulation rates for the loess-palaeosol sequences from the Carpathian Basin. Data in Brief, 2022, 44, 108555.	1.0	1
82	The patchwork loess of Central Asia: Implications for interpreting aeolian dynamics and past climate circulation in piedmont regions. Journal of Quaternary Science, 2023, 38, 526-543.	2.1	3
83	High-resolution OSL dating of loess in Adventdalen, Svalbard: Late Holocene dust activity and permafrost development. Quaternary Science Reviews, 2023, 310, 108137.	3.0	3
84	Disentangling variations of dust concentration in Greenland ice cores over the last glaciation: An overview of current knowledge and new initiative. Earth-Science Reviews, 2023, 242, 104451.	9.1	3
85	Last Glacial loess in Europe: luminescence database and chronology of deposition. Earth System Science Data, 2023, 15, 4689-4711.	9.9	0
86	Mercury records covering the past 90 000Âyears from lakes Prespa and Ohrid, SEÂEurope. Biogeosciences, 2024, 21, 531-556.	3.3	1
87	Paleo±Dust: quantifying uncertainty in paleo-dust deposition across archive types. Earth System Science Data, 2024, 16, 941-959.	9.9	0
88	Ground-penetrating radar investigation of regolith thickness on a periglacial alpine summit flat, Uinta Mountains, Utah, USA. Arctic, Antarctic, and Alpine Research, 2024, 56, .	1.1	0