

# Dust flux estimates for the Last Glacial Period in East C records of loess deposits: a review

Quaternary Science Reviews

29, 3157-3166

DOI: [10.1016/j.quascirev.2010.07.005](https://doi.org/10.1016/j.quascirev.2010.07.005)

Citation Report

#	ARTICLE	IF	CITATIONS
1	Investigating the penultimate and last glacial cycles of the S <sub>1</sub> loess section (Hungary) using luminescence dating, high-resolution grain size, and magnetic susceptibility data. <i>Quaternary International</i> , 2011, 234, 75-85.	0.7	107
2	Tectonic versus climatic control on the evolution of a loess-paleosol sequence at Beremend, Hungary: an integrated approach based on paleoecological, clay mineralogical, and geochemical data. <i>Quaternary International</i> , 2011, 240, 71-86.	0.7	54
3	The Belotinac section (Southern Serbia) at the southern limit of the European loess belt: Initial results. <i>Quaternary International</i> , 2011, 240, 128-138.	0.7	22
4	Dust deposition and climate in the Carpathian Basin over an independently dated last glacial-interglacial cycle. <i>Quaternary Science Reviews</i> , 2011, 30, 662-681.	1.4	214
5	The last million years recorded at the Stari Slankamen (Northern Serbia) loess-palaeosol sequence: revised chronostratigraphy and long-term environmental trends. <i>Quaternary Science Reviews</i> , 2011, 30, 1142-1154.	1.4	169
6	Loess towards (geo) tourism – proposed application on loess in Vojvodina region (north Serbia). <i>Acta Geographica Slovenica</i> , 2011, 51, 391-406.	0.3	34
7	Radiocarbon chronology of Late Pleistocene large mammal faunas from the Pannonian basin (Hungary). <i>Bulletin of Geosciences</i> , 2012, , 13-19.	0.5	10
8	Establishing a luminescence chronology for a palaeosol-loess profile at Tokaj (Hungary): A comparison of quartz OSL and polymineral IRSL signals. <i>Quaternary Geochronology</i> , 2012, 10, 68-74.	0.6	44
9	Pleistocene environmental dynamics recorded in the loess of the middle and lower Danube basin. <i>Quaternary Science Reviews</i> , 2012, 41, 104-118.	1.4	157
10	Evaluating the use of clay mineralogy, Sr-Nd isotopes and zircon U-Pb ages in tracking dust provenance: An example from loess of the Carpathian Basin. <i>Chemical Geology</i> , 2012, 304-305, 83-96.	1.4	78
11	Upbuilding pedogenesis under active loess deposition in a super-humid, temperate climate – quantification of deposition rates, soil chemistry and pedogenic thresholds. <i>Geoderma</i> , 2012, 189-190, 491-501.	2.3	29
12	PALEOSOLS AND WIND-BLOWN SEDIMENTS   Biogeochemical Role of Dust in Quaternary Climate Cycles. , 2013, , 412-420.		0
13	Humid glacials, arid interglacials? Critical thoughts on pedogenesis and paleoclimate based on multi-proxy analyses of the loess-paleosol sequence Crvenka, Northern Serbia. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2013, 387, 165-175.	1.0	102
14	Analysis of Saharan dust intrusions into the Carpathian Basin (Central Europe) over the period of 1979-2011. <i>Global and Planetary Change</i> , 2013, 100, 333-342.	1.6	49
15	Sedimentary characteristics and source of loess in Baranja (Eastern Croatia). <i>Aeolian Research</i> , 2013, 11, 129-139.	1.1	21
16	Towards identifying the origin of metamorphic components in Austrian loess: insights from detrital rutile chemistry, thermometry and U-Pb geochronology. <i>Quaternary Science Reviews</i> , 2013, 75, 132-142.	1.4	29
17	The chronology of the Aarengard II loess-palaeosol section (Eastern Croatia). <i>Geologia Croatica</i> , 2013, 66, 191-203.	0.3	25
18	Late Pleistocene variations of the background aeolian dust concentration in the Carpathian Basin: an estimate using decomposition of grain-size distribution curves of loess deposits. <i>Geologie En Mijnbouw/Netherlands Journal of Geosciences</i> , 2013, 91, 159-171.	0.6	15

#	ARTICLE	IF	CITATIONS
19	Loess in the Vojvodina region (Northern Serbia): an essential link between European and Asian Pleistocene environments. <i>Geologie En Mijnbouw/Netherlands Journal of Geosciences</i> , 2013, 91, 173-188.	0.6	57
20	Eolian contribution to geochemical and mineralogical characteristics of some soil types in Medvednica Mountain, Croatia. <i>Catena</i> , 2014, 117, 145-156.	2.2	8
21	Loess accumulation during the last glacial maximum: Evidence from Urluia, southeastern Romania. <i>Quaternary International</i> , 2014, 334-335, 74-85.	0.7	39
22	Vegetation and environmental responses to climate forcing during the Last Glacial Maximum and deglaciation in the East Carpathians: attenuated response to maximum cooling and increased biomass burning. <i>Quaternary Science Reviews</i> , 2014, 106, 278-298.	1.4	65
23	Introducing an improved multi-proxy approach for paleoenvironmental reconstruction of loessâ€“paleosol archives applied on the Late Pleistocene Nussloch sequence (SW Germany). <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2014, 410, 300-315.	1.0	53
24	The Late Pleistocene Belotinac section (southern Serbia) at the southern limit of the European loess belt: Environmental and climate reconstruction using grain size and stable C and N isotopes. <i>Quaternary International</i> , 2014, 334-335, 10-19.	0.7	50
25	The conceptual sedimentary model of the Lower Danube loess basin: Sedimentogenetic implications. <i>Quaternary International</i> , 2014, 351, 14-24.	0.7	28
26	Tracing the origin of loess in Hungary with the help of heavy mineral composition data. <i>Quaternary International</i> , 2014, 319, 11-21.	0.7	25
27	Two possible source regions for central Greenland last glacial dust. <i>Geophysical Research Letters</i> , 2015, 42, 10,399.	1.5	39
28	Aeolian dynamics at the Orlovat loessâ€“paleosol sequence, northern Serbia, based on detailed textural and geochemical evidence. <i>Aeolian Research</i> , 2015, 18, 69-81.	1.1	56
29	Core-Shell Processing of Natural Pigment: Upper Palaeolithic Red Ochre from Lovas, Hungary. <i>PLoS ONE</i> , 2015, 10, e0131762.	1.1	19
30	Results of paleoecological studies in the loess region of Szeged-Ã–thalom (SE Hungary). <i>Quaternary International</i> , 2015, 372, 66-78.	0.7	30
31	Danube loess stratigraphy â€” Towards a pan-European loess stratigraphic model. <i>Earth-Science Reviews</i> , 2015, 148, 228-258.	4.0	241
32	Climate vs. parent material â€” Which is the key of Stagnosol diversity in Croatia?. <i>Geoderma</i> , 2015, 241-242, 250-261.	2.3	14
33	Compositional trends through the Holocene mud succession of the southwestern Black Sea shelf: Implications for sedimentary provenance and water-level history. <i>Sedimentary Geology</i> , 2015, 316, 13-25.	1.0	15
34	Fine and coarse-quartz SAR-OSL dating of Last Glacial loess in Southern Romania. <i>Quaternary International</i> , 2015, 357, 33-43.	0.7	27
35	Mineralogical discrimination of the pleistocene loess/paleosol sections in Srijem and Baranja, Croatia. <i>Aeolian Research</i> , 2016, 21, 151-162.	1.1	6
36	High-resolution trace element and stable/radiogenic isotope profiles of late Pleistocene to Holocene speleothems from Dim Cave, SW Turkey. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2016, 452, 68-79.	1.0	19

#	ARTICLE	IF	CITATIONS
37	Saharan dust deposition in the Carpathian Basin and its possible effects on interglacial soil formation. <i>Aeolian Research</i> , 2016, 22, 1-12.	1.1	46
38	Sedimentological and mineralogical characteristics of the Pleistocene loess/paleosol sections in the Eastern Croatia. <i>Aeolian Research</i> , 2016, 20, 7-23.	1.1	18
39	The history of Danube loess research. <i>Quaternary International</i> , 2016, 399, 86-99.	0.7	27
40	Coupled European and Greenland last glacial dust activity driven by North Atlantic climate. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2017, 114, E10632-E10638.	3.3	77
41	Reconstructing palaeoenvironments on desert margins: New perspectives from Eurasian loess and Australian dry lake shorelines. <i>Quaternary Science Reviews</i> , 2017, 171, 1-19.	1.4	26
42	Loess correlations – Between myth and reality. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2018, 509, 4-23.	1.0	31
43	The Crvenka loess-paleosol sequence: A record of continuous grassland domination in the southern Carpathian Basin during the Late Pleistocene. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2018, 509, 33-46.	1.0	38
44	Early Upper Paleolithic surface collections from loess-like sediments in the northern Carpathian Basin. <i>Quaternary International</i> , 2018, 485, 167-182.	0.7	9
45	Discriminating luminescence age uncertainty composition for a robust Bayesian modelling. <i>Quaternary Geochronology</i> , 2018, 43, 30-39.	0.6	26
46	The Last Glacial aeolian record of the Island of Susak (Croatia) as seen from a high-resolution grain size and rock magnetic analysis. <i>Quaternary International</i> , 2018, 494, 211-224.	0.7	26
47	New chronology of the best developed loess/paleosol sequence of Hungary capturing the past 1.1 ma: Implications for correlation and proposed pan-Eurasian stratigraphic schemes. <i>Quaternary Science Reviews</i> , 2018, 191, 144-166.	1.4	30
48	Approaches and challenges to the study of loess – Introduction to the LoessFest Special Issue. <i>Quaternary Research</i> , 2018, 89, 563-618.	1.0	92
49	Editorial: Eurasian loess records. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2018, 509, 1-3.	1.0	1
51	Quartz OSL dating of late quaternary Chinese and Serbian loess: A cross Eurasian comparison of dust mass accumulation rates. <i>Quaternary International</i> , 2019, 502, 30-44.	0.7	44
52	Radiocarbon dated malacological records of two Late Pleistocene loess-paleosol sequences from SW-Hungary: Paleoeological inferences. <i>Quaternary International</i> , 2019, 504, 108-117.	0.7	7
53	High-resolution proxy record of the environmental response to climatic variations during transition MIS3/MIS2 and MIS2 in Central Europe: The loess-paleosol sequence of Katymár brickyard (Hungary). <i>Quaternary International</i> , 2019, 504, 40-55.	0.7	18
54	High-resolution paleoclimatic proxy data from the MIS3/2 transition recorded in northeastern Hungarian loess. <i>Quaternary International</i> , 2019, 502, 95-107.	0.7	21
55	Holocene sedimentation in the southwestern Black Sea: Interplay between riverine supply, coastal eddies of the Rim Current, surface and internal waves, and saline underflow through the Strait of Bosphorus. <i>Marine Geology</i> , 2020, 420, 106092.	0.9	4

#	ARTICLE	IF	CITATIONS
56	Initial quartz OSL and dust mass accumulation rate investigation of the Kisljevo loess sequence in north-eastern Serbia. <i>Quaternary International</i> , 2020, , .	0.7	5
57	A 180 ka record of environmental change at Erdut (Croatia): a new chronology for the loessâ€‘palaeosol sequence and its implications for environmental interpretation. <i>Journal of Quaternary Science</i> , 2020, 35, 582-593.	1.1	16
58	A postâ€‘IR IRSL</sc> chronology and dust mass accumulation rates of the Nosak loessâ€‘palaeosol sequence in northeastern Serbia. <i>Boreas</i> , 2020, 49, 841-857.	1.2	16
59	Abrupt last glacial dust fall over southeast England associated with dynamics of the British-Irish ice sheet. <i>Quaternary Science Reviews</i> , 2020, 250, 106641.	1.4	26
60	Reâ€‘analysis of late Quaternary dust mass accumulation rates in Serbia using new luminescence chronology for loessâ€‘palaeosol sequence at Surduk. <i>Boreas</i> , 2020, 49, 634-652.	1.2	14
61	Plio-Pleistocene Dust Traps on Paleokarst Surfaces: A Case Study From the Carpathian Basin. <i>Frontiers in Earth Science</i> , 2020, 8, .	0.8	3
62	Linkage between dust cycle and loess of the Last Glacial Maximum in Europe. <i>Atmospheric Chemistry and Physics</i> , 2020, 20, 4969-4986.	1.9	35
63	<sup>14</sup> C Dated Chronology of the Thickest and Best Resolved Loess/Paleosol Record of the LGM from SE Hungary Based on Comparing Precision and Accuracy of Age-Depth Models. <i>Radiocarbon</i> , 2020, 62, 403-417.	0.8	11
64	Simulated regional dust cycle in the Carpathian Basin and the Adriatic Sea region during the Last Glacial Maximum. <i>Quaternary International</i> , 2021, 581-582, 114-127.	0.7	17
65	Chronological Assessment of the Balta Alba Kurgan Loess-Paleosol Section (Romania) â€‘ A Comparative Study on Different Dating Methods for a Robust and Precise Age Model. <i>Frontiers in Earth Science</i> , 2021, 8, .	0.8	16
66	Loess landscapes of Europe â€‘ Mapping, geomorphology, and zonal differentiation. <i>Earth-Science Reviews</i> , 2021, 215, 103496.	4.0	104
67	Palaeoecological background of the Upper Palaeolithic site of SÃ¡gvÃ¡r, Hungary: radiocarbonâ€‘dated malacological and sedimentological studies on the Late Pleistocene environment. <i>Journal of Quaternary Science</i> , 0, , .	1.1	3
68	Selected Grain-Size and Geochemical Analyses of the Loess-Paleosol Sequence of PÃ¡cel (Northern) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 5 <i>Quaternary</i> , 2021, 4, 17.	1.0	4
69	Spatial differences in the luminescence sensitivity of quartz extracted from Carpathian Basin fluvial sediments. <i>Quaternary Geochronology</i> , 2021, 64, 101166.	0.6	10
70	Chronology and dust mass accumulation history of the Wenchuan loess on eastern Tibetan Plateau since the last glacial. <i>Aeolian Research</i> , 2021, 53, 100748.	1.1	8
71	DunaszekcsÃ‘ Castle Hill: Landslides Along the Danubian Loess Bluff. <i>World Geomorphological Landscapes</i> , 2015, , 113-120.	0.1	7
72	New chronology and extended palaeoenvironmental data to the 1975 loess profile of Madaras brickyard, South Hungary. <i>Journal of Quaternary Science</i> , 2021, 36, 1364-1381.	1.1	3
73	A tale of two signals: Global and local influences on the Late Pleistocene loess sequences in Bulgarian Lower Danube. <i>Quaternary Science Reviews</i> , 2021, 274, 107264.	1.4	16

#	ARTICLE	IF	CITATIONS
74	Detailed Luminescence Dating of Dust Mass Accumulation Rates Over the Last Two Glacial-Interglacial Cycles from the Irig Loess-Palaeosol Sequence, Carpathian Basin. SSRN Electronic Journal, 0, , .	0.4	0
75	A remarkable last glacial loess sedimentation at Roxolany in the Dniester Liman (Southern Ukraine). Quaternary Science Reviews, 2022, 285, 107521.	1.4	2
76	Modeling dust emission in alpine regions with low air temperature and low air pressure – A case study on the Qinghai-Tibetan Plateau (QTP). Geoderma, 2022, 422, 115930.	2.3	6
77	Detailed luminescence dating of dust mass accumulation rates over the last two glacial-interglacial cycles from the Irig loess-palaeosol sequence, Carpathian Basin. Global and Planetary Change, 2022, 215, 103895.	1.6	5
78	Greenland Ice Core Record of Last Glacial Dust Sources and Atmospheric Circulation. Journal of Geophysical Research D: Atmospheres, 2022, 127, .	1.2	17
79	Southeast Asian dust flux reconstructed using accurately dated stalagmite thorium concentrations. Results in Geochemistry, 2022, 7-8, 100017.	0.3	0
80	Last glacial loess dynamics in the Southern Caucasus (NE-Armenia) and the phenomenon of missing loess deposition during MIS-2. Scientific Reports, 2022, 12, .	1.6	4
81	Dataset of dust mass accumulation rates for the loess-palaeosol sequences from the Carpathian Basin. Data in Brief, 2022, 44, 108555.	0.5	1
82	The patchwork loess of Central Asia: Implications for interpreting aeolian dynamics and past climate circulation in piedmont regions. Journal of Quaternary Science, 2023, 38, 526-543.	1.1	3