

CITATION REPORT

List of articles citing

Incurred sample reanalysis (ISR): a decisive tool in bioanalytical research

DOI: 10.4155/bio.11.76
Bioanalysis, 2011, 3, 1007-24.

Source: <https://exaly.com/paper-pdf/49966464/citation-report.pdf>

Version: 2024-04-10

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
117	Determination of (S)-(+)- and (R)-(-)-ibuprofen enantiomers in human plasma after chiral precolumn derivatization by reversed-phase LC-ESI-MS/MS. <i>Bioanalysis</i> , 2012 , 4, 2909-27	2.1	11
116	Regulated drug bioanalysis for human pharmacokinetic studies and therapeutic drug management. <i>Bioanalysis</i> , 2012 , 4, 1919-31	2.1	7
115	Development of an SPE-LC-MS/MS method for simultaneous quantification of bosentan and its active metabolite hydroxybosentan in human plasma to support a bioequivalence study. 2012 , 70, 462-70		9
114	Challenges in the simultaneous quantitation of sumatriptan and naproxen in human plasma: application to a bioequivalence study. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2012 , 902, 122-31	3.2	21
113	Highly sensitive and rapid ultra-performance liquid chromatography-tandem mass spectrometry method for the determination of nifedipine in human plasma and its application to a bioequivalence study. <i>Biomedical Chromatography</i> , 2012 , 26, 1509-18	1.7	6
112	Comparison of fasting bioavailability among 100-mg commercial, 100-mg generic, and 50-mg chewable generic sildenafil tablets in healthy male Mexican volunteers: a single-dose, 3-period, crossover study. 2012 , 34, 689-98		4
111	Application of a rapid and sensitive liquid chromatography-tandem mass spectrometry method for determination of bumetanide in human plasma for a bioequivalence study. 2012 , 66, 365-70		6
110	Analysis of 21-hydroxy deflazacort in human plasma by UPLC-MS/MS: application to a bioequivalence study in healthy volunteers. 2013 , 85, 213-7		3
109	Simultaneous quantification of prodrug oseltamivir and its metabolite oseltamivir carboxylate in human plasma by LC-MS/MS to support a bioequivalence study. 2013 , 3, 149-160		10
108	Current Understanding of Bioanalytical Assay Reproducibility: Incurred Sample Reanalysis, Incurred Sample Stability, and Incurred Sample Accuracy. 2013 , 47-64		2
107	Investigation of ex vivo stability of fesoterodine in human plasma and its simultaneous determination together with its active metabolite 5-HMT by LC-ESI-MS/MS: Application to a bioequivalence study. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2013 , 913-914, 1-11	3.2	2
106	Simultaneous analysis of oxybutynin and its active metabolite N-desethyl oxybutynin in human plasma by stable isotope dilution LC-MS/MS to support a bioequivalence study. 2013 , 84, 244-55		11
105	Practical approaches to incurred sample LC-MS/MS reanalysis: confirming unexpected results. 2013 , 85, 2405-13		2
104	SPE-UPLC-MS/MS method for sensitive and rapid determination of aripiprazole in human plasma to support a bioequivalence study. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2013 , 925, 20-5	3.2	21
103	UPLC-MS/MS assay for the simultaneous quantification of carvedilol and its active metabolite 4-hydroxyphenyl carvedilol in human plasma to support a bioequivalence study in healthy volunteers. <i>Biomedical Chromatography</i> , 2013 , 27, 974-86	1.7	16
102	Sensitive and selective determination of diflunisal in human plasma by LC-MS. 2013 , 51, 872-82		8
101	Phase I and beyond - the importance of switching to patient plasma: considerations and perspectives on bioanalytical procedures. <i>Bioanalysis</i> , 2013 , 5, 2215-8	2.1	

100	Ex vivo conversion of prodrug prothionamide to its metabolite prothionamide sulfoxide with different extraction techniques and their estimation in human plasma by LC-MS/MS. <i>Bioanalysis</i> , 2013 , 5, 185-200	2.1	2
99	Quantitation of nitrofurantoin in human plasma by liquid chromatography tandem mass spectrometry. 2013 , 63, 141-58		8
98	Reliable and sensitive determination of dutasteride in human plasma by liquid chromatography-tandem mass spectrometry. <i>Biomedical Chromatography</i> , 2013 , 27, 1168-76	1.7	5
97	Stable-isotope dilution LC-MS/MS assay for determination of iloperidone and its two major metabolites, P 88 and P 95, in human plasma: application to a bioequivalence study. <i>Bioanalysis</i> , 2013 , 5, 669-86	2.1	6
96	Incurring sample reanalysis evaluation and impact on drug preclinical and clinical development. 2014 , 134-147		
95	ISR: background, evolution and implementation, with specific consideration for ligand-binding assays. <i>Bioanalysis</i> , 2014 , 6, 393-402	2.1	7
94	Systematic evaluation of plasma phospholipids for reliable and precise determination of dronedarone and desbutyldronedarone by LC-MS/MS. <i>Bioanalysis</i> , 2014 , 6, 2635-50	2.1	1
93	Impact of electrospray ion source platforms on matrix effect due to plasma phospholipids in the determination of rivastigmine by LC-MS/MS. <i>Bioanalysis</i> , 2014 , 6, 2301-16	2.1	3
92	Simultaneous determination of ornidazole and its main metabolites in human plasma by LC-MS/MS: application to a pharmacokinetic study. <i>Bioanalysis</i> , 2014 , 6, 2343-56	2.1	18
91	Development and validation of a rapid and sensitive LC-MS/MS method for the determination of aripiprazole in human plasma: Application to a bioequivalence study. 2014 , 26, 203-227		4
90	Simultaneous determination of metoprolol and its metabolites, hydroxymetoprolol and O-desmethylnmetoprolol, in human plasma by liquid chromatography with tandem mass spectrometry: Application to the pharmacokinetics of metoprolol associated with CYP2D6 genotypes. 2014 , 37, 1256-64		23
89	Incurring sample reanalysis in bioequivalence studies for abbreviated new drug applications. <i>Bioanalysis</i> , 2014 , 6, 3349-54	2.1	2
88	Determination of mycophenolic acid in human plasma by ultra performance liquid chromatography tandem mass spectrometry. 2014 , 4, 205-216		12
87	Method of variability optimization in pharmacokinetic data analysis. 2014 , 39, 111-9		5
86	Development of a sensitive and rapid method for quantitation of (-)- and (+)-metoprolol in human plasma by chiral LC-ESI-MS/MS. 2014 , 4, 63-79		9
85	WITHDRAWN: Simultaneous determination of glimepiride and fluoxetine in human plasma by ultra performance liquid chromatography tandem mass spectrometry. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2014 ,	3.2	
84	Parallel achiral-chiral determination of oxybutynin, N-desethyl oxybutynin and their enantiomers in human plasma by LC-MS/MS to support a bioequivalence trial. 2014 , 88, 81-91		6
83	Rapid and sensitive UPLC-MS-MS determination of tacrolimus in Wistar rats and human blood. 2014 , 52, 59-67		9

82	FDA Bioequivalence Standards. <i>AAPS Advances in the Pharmaceutical Sciences Series</i> , 2014 ,	0.5	9
81	Analysis and pharmacokinetics studies of gastrodin and p-hydroxybenzyl alcohol in dogs using ultra fast liquid chromatography-tandem mass spectrometry method. 2014 , 99, 83-8		16
80	Sensitive, selective and rapid determination of lafutidine in human plasma by solid phase extraction-liquid chromatography-tandem mass spectrometry. 2014 , 69, 448-460		
79	Good Clinical Practice. 2014 , 756-787		
78	Application of a UPLC-MS/MS method for the analysis of alosetron in human plasma to support a bioequivalence study in healthy males and females. <i>Biomedical Chromatography</i> , 2015 , 29, 1527-34	1.7	1
77	Simultaneous analysis of losartan, its active metabolite, and hydrochlorothiazide in human plasma by a UPLC-MS/MS method. 2015 , 39, 714-733		14
76	Determination of cilostazol and its active metabolite 3,4-dehydro cilostazol from small plasma volume by UPLC-MS/MS. 2015 , 5, 1-11		5
75	Bioavailability and pharmacokinetics profile of helcid in beagle dogs using gradient elution high performance liquid chromatography electrospray ionization mass spectrometry. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2015 , 988, 8-12	3.2	6
74	Validated UPLC-MS/MS method for determination of hordenine in rat plasma and its application to pharmacokinetic study. 2015 , 111, 131-7		5 ¹
73	Selective and rapid determination of raltegravir in human plasma by liquid chromatography-tandem mass spectrometry in the negative ionization mode. 2015 , 5, 101-109		2
72	High-sensitivity simultaneous liquid chromatography-tandem mass spectrometry assay of ethinyl estradiol and levonorgestrel in human plasma. 2015 , 5, 316-326		11
71	Quantification of lamotrigine in patient plasma using a fast liquid chromatography-tandem mass spectrometry method with backflush technology. 2015 , 37, 188-97		7
70	Reliable LC-MS/MS assay for the estimation of rilpivirine in human plasma: application to a bioequivalence study and incurred sample reanalysis. 2015 , 7, 290-9		3
69	Rat dried blood spot analysis of (R,S)-(-)- and (S,R)-(+)- enantiomers of emtricitabin on immobilized tris-(3,5-dimethylphenyl carbamate) amylose silica as a chiral stationary phase. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2015 , 1002, 160-8	3.2	9
68	Analysis of imprecision in incurred sample reanalysis for small molecules. <i>AAPS Journal</i> , 2015 , 17, 206-153.7		5
67	Simultaneous quantification of atenolol and chlorthalidone in human plasma by ultra-performance liquid chromatography-tandem mass spectrometry. <i>Biomedical Chromatography</i> , 2016 , 30, 208-16	1.7	15
66	A simple LC-ESI-MS/MS method for quantification of bacopaside I in rat plasma and its application to a pharmacokinetic study. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2016 , 1027, 103-9	3.2	
65	An improved LC-MS/MS method for the simultaneous determination of pyrazinamide, pyrazinoic acid and 5-hydroxy pyrazinoic acid in human plasma for a pharmacokinetic study. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2016 , 1017-1018, 52-61	3.2	8

64	Simultaneous bioanalysis of rasagiline and its major metabolites in human plasma by LC-MS/MS: Application to a clinical pharmacokinetic study. 2016 , 125, 280-5		6
63	A novel, sensitive and selective method of UPLC/MS-MS for rapid simultaneous determination of midodrine and its active metabolite desglymidodrine in human plasma: Application to support bioequivalence study in healthy human volunteers. 2016 , 131, 355-363		7
62	SPE-UPLC-MS/MS assay for determination of letrozole in human plasma and its application to bioequivalence study in healthy postmenopausal Indian women. 2016 , 6, 276-281		11
61	LC-tandem mass spectrometry method for the simultaneous determination of metformin and sitagliptin in human plasma after ion-pair solid phase extraction. 2016 , 131, 64-70		32
60	Determination of torasemide in human plasma and its bioequivalence study by high-performance liquid chromatography with electrospray ionization tandem mass spectrometry. 2016 , 6, 95-102		3
59	Platform switching from ELISA to Gyrolab® a novel generic reagent omits the need to change critical reagents. <i>Bioanalysis</i> , 2016 , 8, 807-14	2.1	1
58	Application of an LC-MS/MS method for reliable determination of amodiaquine, N-desethylamodiaquine, artesunate and dihydroartemisinin in human plasma for a bioequivalence study in healthy Indian subjects. 2016 , 124, 67-78		7
57	Liquid chromatography--tandem mass spectrometry method for simultaneous determination of albendazole and albendazole sulfoxide in human plasma for bioequivalence studies. 2016 , 6, 226-234		10
56	Determination of lercanidipine in human plasma by an improved UPLC-MS/MS method for a bioequivalence study. 2016 , 6, 87-94		6
55	Simultaneous determination of imperatorin and its metabolite xanthotoxol in rat plasma and urine by LC-MS/MS and its application to pharmacokinetic studies. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2017 , 1044-1045, 30-38	3.2	7
54	Simultaneous quantification of amiloride and hydrochlorothiazide in human plasma by liquid chromatography-tandem mass spectrometry. 2017 , 7, 288-296		8
53	Determination of ergocalciferol in human plasma after Diels-Alder derivatization by LC-MS/MS and its application to a bioequivalence study. 2017 , 7, 417-422		7
52	Fast and sensitive LC-MS/MS method for the simultaneous determination of lisinopril and hydrochlorothiazide in human plasma. 2017 , 7, 163-169		9
51	An improved LC-MS/MS method for the quantification of alverine and para hydroxy alverine in human plasma for a bioequivalence study. 2017 , 7, 95-102		1
50	Simultaneous analysis of glucocorticosteroid fluticasone propionate and its metabolite fluticasone propionate 17β-carboxylic acid in human plasma by UPLC-MS/MS at sub pg/mL level. 2017 , 135, 1-7		8
49	An HPLC-MS/MS method for quantitation of Gly-MCA in mouse plasma: Application to a pharmacokinetic study. 2017 , 146, 53-58		1
48	Comprehensive graphical presentation of data from incurred sample reanalysis. <i>Bioanalysis</i> , 2017 , 9, 947-956	2.1	10
47	Assessment of critical extraction and chromatographic parameters for the determination of bupropion and its three primary metabolites in human plasma by LC-MS/MS. 2017 , 135, 81-90		7

46	Application of an LC-MS/MS method for the analysis of amlodipine, valsartan and hydrochlorothiazide in polypill for a bioequivalence study. 2017 , 7, 309-316		16
45	LC-MS/MS analysis of metformin, saxagliptin and 5-hydroxy saxagliptin in human plasma and its pharmacokinetic study with a fixed-dose formulation in healthy Indian subjects. <i>Biomedical Chromatography</i> , 2017 , 31, e3809	1.7	18
44	Simultaneous analysis of allopurinol and oxypurinol using a validated liquid chromatography-tandem mass spectrometry method in human plasma. 2017 , 7, 56-62		12
43	Review of the bioanalytical methods for the determination of methotrexate and its metabolites in in vitro, preclinical and clinical studies: Case studies and perspectives. <i>Biomedical Chromatography</i> , 2017 , 31, e3849	1.7	5
42	Sensitive and rapid determination of amantadine without derivatization in human plasma by LC-MS/MS for a bioequivalence study. 2018 , 8, 202-207		5
41	Quantification of metronidazole in human plasma using a highly sensitive and rugged LC-MS/MS method for a bioequivalence study. <i>Biomedical Chromatography</i> , 2018 , 32, e4242	1.7	8
40	Quantitation of tadalafil in human plasma using a sensitive and rapid LC-MS/MS method for a bioequivalence study. 2018 , 8, 271-276		11
39	Accurate determination of a novel vasodilatory Eblocker TJ0711 using LC-MS/MS: Resolution of an isobaric metabolite interference in dog plasma. <i>Biomedical Chromatography</i> , 2018 , 32, e4196	1.7	0
38	Highly sensitive LC-MS/MS method to estimate doxepin and its metabolite nordoxepin in human plasma for a bioequivalence study. 2018 , 8, 378-385		6
37	Determination of silodosin and its active glucuronide metabolite KMD-3213G in human plasma by LC-MS/MS for a bioequivalence study. <i>Biomedical Chromatography</i> , 2018 , 32, e4041	1.7	6
36	Quantification of root phosphite concentrations for evaluating the potential of foliar phosphonate sprays for the management of avocado root rot. 2018 , 103, 87-97		7
35	Ultra-performance liquid chromatography-tandem mass spectrometry assay for determination of plasma norgestrel acetate and estradiol in healthy postmenopausal women. <i>Biomedical Chromatography</i> , 2018 , 32, e4086	1.7	3
34	Overcoming interference of plasma phospholipids using HybridSPE for the determination of trimetazidine by UPLC-MS/MS. <i>Biomedical Chromatography</i> , 2018 , 32, e4095	1.7	3
33	Simultaneous determination of etonogestrel and ethinyl estradiol in human plasma by UPLC-MS/MS and its pharmacokinetic study. <i>Biomedical Chromatography</i> , 2018 , 32, e4165	1.7	2
32	Therapeutic Drug Monitoring of Busulfan for the Management of Pediatric Patients: Cross-Validation of Methods and Long-Term Performance. 2018 , 40, 84-92		12
31	ESTIMATION OF CELECOXIB IN HUMAN PLASMA BY RAPID AND SELECTIVE LC-MS/MS METHOD FOR A BIOEQUIVALENCE STUDY. 2018 , 10, 16		4
30	A decade of incurred sample reanalysis: failures, investigations and impact. <i>Bioanalysis</i> , 2018 , 10, 1767-1772		4
29	Incurred sample reanalysis at GSK: what have we learned?. <i>Bioanalysis</i> , 2018 , 10, 1755-1766	2.1	2

28	Simultaneous quantitation of rosuvastatin and ezetimibe in human plasma by LC-MS/MS: Pharmacokinetic study of fixed-dose formulation and separate tablets. <i>Biomedical Chromatography</i> , 2018 , 32, e4291	1.7	2
27	Determination of asenapine in presence of its inactive metabolites in human plasma by LC-MS/MS. 2018 , 8, 341-347		7
26	Ion-pair solid phase extraction for the simultaneous separation and quantitation of metformin and canagliflozin in human plasma by LC-MS/MS. 2018 , 143, 181-189		13
25	Determination of a PDE4 inhibitor Hemay005 in human plasma and urine by UPLC-MS/MS and its application to a PK study. <i>Bioanalysis</i> , 2018 , 10, 863-875	2.1	4
24	Challenges in Simultaneous Determination of Hydrochlorothiazide and Ramipril in Human Plasma: Application to a Bioequivalence Study. 2018 , 56, 867-878		2
23	An ultra HPLC-MS/MS method for quantification of hypidone hydrochloride (YL-0919) and application to a pharmacokinetic study. <i>Bioanalysis</i> , 2019 , 11, 1243-1254	2.1	0
22	CX1003 quantification by ultra-performance LC-MS/MS in human plasma and its application to a pharmacokinetic study in solid tumor patients. <i>Bioanalysis</i> , 2019 , 11, 1483-1493	2.1	
21	Determination of BPI15086 and its metabolite in human plasma by ultra-high performance liquid chromatography-MS/MS and its application to a pharmacokinetic study. <i>Bioanalysis</i> , 2019 , 11, 773-784	2.1	
20	Determination of terbinafine in human plasma using UPLC-MS/MS: Application to a bioequivalence study in healthy subjects. <i>Biomedical Chromatography</i> , 2019 , 33, e4543	1.7	3
19	Determination of apatinib and its three active metabolites by UPLC-MS/MS in a Phase IV clinical trial in NSCLC patients. <i>Bioanalysis</i> , 2019 , 11, 2049-2060	2.1	1
18	Mixed-mode solid phase extraction combined with LC-MS/MS for determination of empagliflozin and linagliptin in human plasma. 2019 , 145, 523-531		21
17	Simultaneous quantitation of metformin and dapagliflozin in human plasma by LC-MS/MS: Application to a pharmacokinetic study. <i>Biomedical Chromatography</i> , 2019 , 33, e4453	1.7	12
16	Development and validation of a rapid and sensitive UPLC-MS/MS assay for the quantification of tofacitinib in human plasma. <i>Biomedical Chromatography</i> , 2019 , 33, e4458	1.7	5
15	Quantification of fenoprofen in human plasma using UHPLC-tandem mass spectrometry for pharmacokinetic study in healthy subjects. <i>Biomedical Chromatography</i> , 2020 , 34, e4708	1.7	2
14	Optimization of chromatography to overcome matrix effect for reliable estimation of four small molecular drugs from biological fluids using LC-MS/MS. <i>Biomedical Chromatography</i> , 2020 , 34, e4777	1.7	2
13	Bioanalysis of biological matrix samples using liquid chromatography-tandem mass spectrometry detection. 2020 , 641-659		0
12	Selective quantification of lacosamide in human plasma using UPLC-MS/MS: Application to pharmacokinetic study in healthy subjects with different doses. <i>Biomedical Chromatography</i> , 2020 , 34, e4928	1.7	1
11	Development and Validation of Liquid Chromatography-Tandem Mass Spectrometry Method for Simple Analysis of Sumatriptan and its Application in Bioequivalence Study. <i>Pharmaceuticals</i> , 2020 , 13,	5.2	4

10	Factors affecting the stability of drugs and their metabolites in biological matrices. 2021 , 517-537		1
9	Development of Fast and Simple LC-ESI-MS/MS Technique for the Quantification of Regorafenib; Application to Pharmacokinetics in Healthy Rabbits. <i>Current Pharmaceutical Analysis</i> , 2021 , 17, 554-563	0.6	1
8	A Highly Sensitive LC-MS/MS Method Development and Validation of Fedratinib in Human Plasma and Pharmacokinetic Evaluation in Healthy Rabbits. <i>Current Pharmaceutical Analysis</i> , 2021 , 17, 782-791	0.6	1
7	Using a stable isotope-labeled internal standard for liquid chromatography-tandem mass spectrometry quantitation of meloxicam in human plasma. <i>Biomedical Chromatography</i> , 2021 , 35, e5217	1.7	1
6	Bioanalysis. <i>AAPS Advances in the Pharmaceutical Sciences Series</i> , 2014 , 419-458	0.5	1
5	Application of a Reliable LC-MS/MS Method for Determination of Rizatriptan in Healthy Subject Samples: Demonstration of Assay Reproducibility by Incurred Sample Reanalysis. <i>ISRN Chromatography</i> , 2012 , 2012, 1-10		1
4	Handling unstable analytes: literature review and expert panel survey by Japan Bioanalysis Forum Discussion Group.. <i>Bioanalysis</i> , 2021 ,	2.1	
3	Is Incurred Sample Reanalysis (ISR) Applicable in Biomarker Assays?. <i>AAPS Journal</i> , 2022 , 24, 65	3.7	
2	Liquid chromatography-tandem mass spectrometry methods for quantification of roxadustat (FG-4592) in human plasma and urine and the applications in two clinical pharmacokinetic studies. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2022 , 123274	3.2	0
1	Recent molecularly imprinted polymers applications in bioanalysis.		0