CITATION REPORT List of articles citing

Impact of switching from an open to a closed infusion system on rates of central line-associated bloodstream infection: a meta-analysis of time-sequence cohort studies in 4 countries

DOI: 10.1086/657632 Infection Control and Hospital Epidemiology, 2011, 32, 50-8.

Source: https://exaly.com/paper-pdf/50057536/citation-report.pdf

Version: 2024-04-17

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
565	Open versus closed IV infusion systems: a state based model to predict risk of catheter associated blood stream infections. 2011 , 1, e000188		5
564	Influence of parenteral nutrition delivery system on the development of bloodstream infections in critically ill patients: an international, multicenter, prospective, open-label, controlled studyEPICOS study. 2012 , 36, 574-86		44
563	A window into diaphragm kinetics: feasibility, precision, and physiological meaning of ultrasound measurements of diaphragm thickness. <i>Critical Care</i> , 2013 , 17,	10.8	1
562	Impact of an international nosocomial infection control consortium multidimensional approach on central line-associated bloodstream infection rates in adult intensive care units in eight cities in India. <i>International Journal of Infectious Diseases</i> , 2013 , 17, e1218-24	10.5	50
561	Abstracts of the 33rd International Symposium on Intensive Care and Emergency Medicine. Brussels, Belgium. March 19-22, 2013. <i>Critical Care</i> , 2013 , 17 Suppl 2, P1-545	10.8	5
560	Reversible depressive effect of TNFIbn a model of isolated perfused rat heart. <i>Critical Care</i> , 2013 , 17,	10.8	78
559	Effect of cdp-choline on microcirculatory alterations during endotoxemia. <i>Critical Care</i> , 2013 , 17,	10.8	78
558	Immune response after stimulation with wall components of Gram-positive bacteria. <i>Critical Care</i> , 2013 , 17,	10.8	78
557	Correlation of the oxygen radical activity and antioxidants and severity in critically ill surgical patients: preliminary report. <i>Critical Care</i> , 2013 , 17,	10.8	78
556	Simultaneous analysis of the expression of CD64 and HLA-DR in the peripheral blood and bronchoalveolar lavage fluid in sepsis. <i>Critical Care</i> , 2013 , 17,	10.8	78
555	Anti-inflammatory effects of Kupffer cells through 🛭 -nicotinic acetylcholine receptors. <i>Critical Care</i> , 2013 , 17,	10.8	1
554	Prevention of sepsis by correcting apoptosis. <i>Critical Care</i> , 2013 , 17,	10.8	78
553	Immune paralysis in trauma patients; implications for prehospital intervention. <i>Critical Care</i> , 2013 , 17,	10.8	2
552	Is hemoglobin concentration affected by sepsis in the acute phase?. Critical Care, 2013, 17,	10.8	3
551	Do changes in red blood cell deformability in patients with septic shock correlate with changes in SOFA scores?. <i>Critical Care</i> , 2013 , 17,	10.8	78
550	Do erythrocytes subjected to cardiopulmonary bypass exhibit changes in their membrane mechanical properties?. <i>Critical Care</i> , 2013 , 17,	10.8	78
549	Platelet-associated oxidative stress and ADAMTS-13 levels are inversely associated with a poor prognosis in septic shock. <i>Critical Care</i> , 2013 , 17,	10.8	78

548	Neutrophil gelatinase-associated lipocalin/lipocalin2, derived from gut crypt cells, exerts intestinal antimicrobial effect via bacterial stimulation of Toll-like receptor 4 and 9. <i>Critical Care</i> , 2013 , 17,	10.8	78
547	Lethal influenza virus A H1N1 infection in two relatives with autosomal dominant GATA-2 deficiency. <i>Critical Care</i> , 2013 , 17,	10.8	2
546	Bacterial translocation primes proinflammatory responses and is connected to early death in an experimental model of lethal injury. <i>Critical Care</i> , 2013 , 17,	10.8	78
545	Effects of the common 34C>T variant of the AMPD1 gene on immune function, multiorgan dysfunction and mortality in patients with sepsis. <i>Critical Care</i> , 2013 , 17,	10.8	78
544	Exploring the translational disconnect between the murine and human inflammatory response: in vitro analysis of the dose-response relationship of LPS and NFB activation in murine and human immune cells. <i>Critical Care</i> , 2013 , 17,	10.8	78
543	Hypogammaglobulinemia in sepsis is not correlated to high circulating angiopoietin-2 levels. <i>Critical Care</i> , 2013 , 17,	10.8	1
542	TREM-1 levels are elevated in patients with liver cirrhosis. Critical Care, 2013, 17,	10.8	78
541	Effects of different doses and serotypes of LPS on blood-brain barrier permeability in Sprague-Dawley rats. <i>Critical Care</i> , 2013 , 17,	10.8	1
540	Bioenergetic imbalance and oxidative stress in the pathophysiology of septic encephalopathy. <i>Critical Care</i> , 2013 , 17,	10.8	78
539	Pathophysiology of sepsis-associated brain dysfunction: an experimental study of cerebral microdialysis and mitochondrial function. <i>Critical Care</i> , 2013 , 17,	10.8	78
538	Cholinergic modulation of hippocampal activity during septic encephalopathy. <i>Critical Care</i> , 2013 , 17,	10.8	1
537	Neutrophil-lymphocyte count ratio as a biomarker of severe sepsis in Escherichia coli infections in adults. <i>Critical Care</i> , 2013 , 17,	10.8	3
536	Obesity and inflammatory markers in severe sepsis. Critical Care, 2013, 17,	10.8	78
535	Cytokine gene expression can predict infectious complications following severe trauma. <i>Critical Care</i> , 2013 , 17,	10.8	3
534	A cohort study of routinely used sepsis biomarkers and 28-day mortality. Critical Care, 2013, 17,	10.8	78
533	Procalcitonin as prognostic marker of mortality. <i>Critical Care</i> , 2013 , 17,	10.8	78
532	Changes in circulating procalcitonin versus C-reactive protein in predicting evolution of infectious disease in febrile, critically ill patients. <i>Critical Care</i> , 2013 , 17,	10.8	78
531	Procalcitonin-guided antibiotic therapy in patients with congestive heart failure and suspicion of lower respiratory tract infection: results from a randomized trial. <i>Critical Care</i> , 2013 , 17,	10.8	2

530	Pro-adrenomedullin as a clinical predictor after cardiac surgery. Critical Care, 2013, 17,	10.8	78
529	Pro-adrenomedullin as prognostic biomarker in the sepsis. <i>Critical Care</i> , 2013 , 17,	10.8	78
528	Early IL-6 response in sepsis is correlated with mortality and severity score. <i>Critical Care</i> , 2013 , 17,	10.8	6
527	Compared values of presepsin (sCD14-ST) and procalcitonin as early markers of outcome in severe sepsis and septic shock: a preliminary report from the Albumin Italian Outcome Sepsis (ALBIOS) study. <i>Critical Care</i> , 2013 , 17,	10.8	78
526	Usefulness of presepsin in the diagnosis of sepsis in acute kidney injury patients. <i>Critical Care</i> , 2013 , 17,	10.8	78
525	Serum levels of presepsin reffects the APACHE II and SOFA scores in patients with sepsis. <i>Critical Care</i> , 2013 , 17,	10.8	1
524	Relationship between serum nitrite/nitrate levels in the early phase of septic acute lung injury and prognosis. <i>Critical Care</i> , 2013 , 17,	10.8	78
523	Simple and fast prediction of Legionella sp. in community-acquired pneumonia: validation of a prediction rule. <i>Critical Care</i> , 2013 , 17,	10.8	78
522	Host genetic variants associated with community-acquired pneumonia. Critical Care, 2013, 17,	10.8	1
521	Outcome of severe community-acquired pneumonia: the impact of comorbidities. <i>Critical Care</i> , 2013 , 17,	10.8	1
520	Prognostic scores and infective endocarditis. <i>Critical Care</i> , 2013 , 17,	10.8	78
519	Effectiveness of early ureteric stenting for urosepsis associated with urinary tract calculi. <i>Critical Care</i> , 2013 , 17,	10.8	78
518	Outcomes in urinary sepsis. <i>Critical Care</i> , 2013 , 17,	10.8	78
517	Nosocomial pneumonia in the postoperative period after heart transplantation. <i>Critical Care</i> , 2013 , 17,	10.8	2
516	Abdominal infection plays a role in the incidence of ventilator-associated pneumonia. <i>Critical Care</i> , 2013 , 17,	10.8	78
515	Incidence and risk factors for ventilator-associated pneumonia in Siriraj Hospital. <i>Critical Care</i> , 2013 , 17,	10.8	78
514	Hospital-acquired bloodstream infection: Indian perspective. Critical Care, 2013, 17,	10.8	78
513	Risk factors for catheter-related bloodstream infections in a surgical ICU. Critical Care, 2013, 17,	10.8	78

512	Analysis of risk factors for catheter-related bloodstream infection in a parenteral nutrition population. <i>Critical Care</i> , 2013 , 17,	10.8	78
511	Can we predict the postoperative infections?. <i>Critical Care</i> , 2013 , 17,	10.8	78
510	Results for introduction of a new hand hygiene program in the ICU. Critical Care, 2013, 17,	10.8	78
509	Improving hand hygiene performance in a surgical ICU: a novel method for data collection to change physician behavior. <i>Critical Care</i> , 2013 , 17,	10.8	78
508	Compliance with the implementation of an ICU cluster-randomized trial assessing the benefits and potential harms of universal glove and gowning. <i>Critical Care</i> , 2013 , 17,	10.8	1
507	Fighting hospital sepsis. <i>Critical Care</i> , 2013 , 17,	10.8	78
506	Sinks as a correctable source of ESBL contamination for patients in the ICU. Critical Care, 2013, 17,	10.8	78
505	Hemodynamic and mitochondrial effects of enalapril in experimental sepsis. Critical Care, 2013, 17,	10.8	78
504	Effects of intraperitoneal immunoglobulin therapy on behavior and cognitive functions in CLP-induced sepsis model in rats. <i>Critical Care</i> , 2013 , 17,	10.8	78
503	Clinical efficacy of recombinant human soluble thrombomodulin in patients with septic disseminated intravascular coagulation. <i>Critical Care</i> , 2013 , 17,	10.8	78
502	Antithrombin III concentrate therapy may have an efficacy in sepsis. Critical Care, 2013, 17,	10.8	78
501	A clinical randomized trial on the use of atorvastatin in patients with sepsis or septic shock: effects on endothelial function. <i>Critical Care</i> , 2013 , 17,	10.8	3
500	A multicenter randomized controlled study of an extracorporeal cytokine hemoadsorption device in septic patients. <i>Critical Care</i> , 2013 , 17,	10.8	26
499	Continuous renal replacement therapy with the adsorbent membrane oXiris in septic patients: a clinical experience. <i>Critical Care</i> , 2013 , 17,	10.8	5
498	Combined use of pumpless extracorporeal lung assist system and continuous renal replacement therapy with citrate anticoagulation in polytrauma patients. <i>Critical Care</i> , 2013 , 17,	10.8	78
497	Abstract withdrawn. <i>Critical Care</i> , 2013 , 17,	10.8	78
496	Use of extracorporeal endotoxin elimination therapy for septic shock. Critical Care, 2013, 17,	10.8	1
495	Selective adsorption of lipopolysaccharide in the complex treatment of patients with severe sepsis. <i>Critical Care</i> , 2013 , 17,	10.8	78

494	Alteco endotoxin hemoadsorption in Gram-negative septic shock patients. Critical Care, 2013, 17,	10.8	78
493	Steroid therapy in septic shock: survey of practice amongst UK critical care physicians. <i>Critical Care</i> , 2013 , 17,	10.8	78
492	Need for restoration of cortisol serum levels for successful antimicrobial therapy in experimental sepsis. <i>Critical Care</i> , 2013 , 17,	10.8	78
491	Impact of time to initiation of appropriate antibiotic therapy on early mortality of patients with septic shock. <i>Critical Care</i> , 2013 , 17,	10.8	1
490	Comparison of survivors and nonsurvivors in 27 confirmed injectional anthrax cases from the 2009 outbreak in Scotland. <i>Critical Care</i> , 2013 , 17,	10.8	3
489	Efficacy and utility of preemptive anti-Gr(+) coverage in selected critically ill patients. <i>Critical Care</i> , 2013 , 17,	10.8	78
488	Gram-negative resistance and need for ICU among urinary tract infections in the United States. <i>Critical Care</i> , 2013 , 17,	10.8	1
487	Multidrug resistance among P. aeruginosa and Enterobacteriaceae in US hospitals, 2000 to 2009. <i>Critical Care</i> , 2013 , 17,	10.8	78
486	Prevalence of multidrug resistance, extensive drug resistance and pandrug resistance among multiple Gram-negative isolates: experience in a tertiary-care hospital ICU in North India. <i>Critical Care</i> , 2013 , 17,	10.8	78
485	Risk factors associated with Acinetobacter baumannii septicemia and its mortality rates in critically ill patients. <i>Critical Care</i> , 2013 , 17,	10.8	O
484	Prevalence of colonization by multiresistant bacteria on admission to the ICU in the French military hospital in Kabul, Afghanistan. <i>Critical Care</i> , 2013 , 17,	10.8	78
483	Extended-spectrum Elactamase Providencia stuartii in a general ICU. Critical Care, 2013, 17,	10.8	78
482	High-dose tigecycline use in severe infections. <i>Critical Care</i> , 2013 , 17,	10.8	1
481	In vitro efficiency of Amikacin Inhale, a novel drug-device delivery system. <i>Critical Care</i> , 2013 , 17,	10.8	4
480	Analysis of parameters in critically ill patients influencing amikacin peak levels. <i>Critical Care</i> , 2013 , 17,	10.8	78
479	Cost implication of antibiotic-associated diarrhoea and the financial impact of probiotic use in its prevention. <i>Critical Care</i> , 2013 , 17,	10.8	1
478	Empirical versus preemptive antimycotic therapy in terms of outcome benefit. <i>Critical Care</i> , 2013 , 17,	10.8	1
477	Impact of echinocandins as first-choice therapy in cardiothoracic high-risk patients with candidemia. <i>Critical Care</i> , 2013 , 17,	10.8	78

476	Fungal ventricular assist device infections occur in colonized patients and are associated with high mortality rate. <i>Critical Care</i> , 2013 , 17,	10.8	78
475	Cost-effectiveness analysis of anidulafungin in the treatment of candidaemia. <i>Critical Care</i> , 2013 , 17,	10.8	78
474	Prevalence and impact of invasive fungal infections in intensive care. Critical Care, 2013, 17,	10.8	78
473	Beliefs and actual practice of oxygen therapy in the ICU. Critical Care, 2013, 17,	10.8	1
472	FiO2/PEEP index: a simple tool for opitimizing ventilator settings. <i>Critical Care</i> , 2013 , 17,	10.8	78
471	Use of high-flow nasal canulae: effect on alveolar pressure and its limitation. Critical Care, 2013, 17,	10.8	3
470	Relation between PaO2/FiO2 ratio, SpO2/FiO2 ratio, oxygenation index and ventilation ratio in critically ill patients. <i>Critical Care</i> , 2013 , 17,	10.8	78
469	Decreased respiratory complexity during apnea in acute respiratory distress syndrome. <i>Critical Care</i> , 2013 , 17,	10.8	78
468	Acute respiratory distress syndrome - the Berlin definition: impact on an ICU of a university hospital. <i>Critical Care</i> , 2013 , 17,	10.8	78
467	Extravascular lung water, B-type natriuretic peptide and blood volume contraction for diagnosing weaning-induced pulmonary edema. <i>Critical Care</i> , 2013 , 17,	10.8	1
466	Pressure support ventilation with minimal sedation as the main ventilatory mode in critically ill patients with lung injury: effect on mortality and incidence of complications. <i>Critical Care</i> , 2013 , 17,	10.8	78
465	Comparison of a fully automatic ARDSNet protocol and a feedback-controlled open lung management concept. <i>Critical Care</i> , 2013 , 17,	10.8	78
464	Comparison of an entirely automated ventilation mode, Intellivent-ASV, with conventional ventilation in ARDS patients: a 48-hour study. <i>Critical Care</i> , 2013 , 17,	10.8	1
463	Difference in pulmonary permeability between indirect and direct acute respiratory distress syndrome assessed by the transpulmonary thermodilution technique. <i>Critical Care</i> , 2013 , 17,	10.8	78
462	Biomarkers for early stage of acute respiratory distress syndrome in septic patients: surfactant protein D and Clara cell protein. <i>Critical Care</i> , 2013 , 17,	10.8	1
461	Intravenous perfluorocarbons for prevention of ventilator-associated ARDS. Critical Care, 2013, 17,	10.8	78
460	Effect of angiotensin-converting enzyme gene I/D polymorphism and its expression on clinical outcome in acute respiratory distress syndrome. <i>Critical Care</i> , 2013 , 17,	10.8	78
459	End-expiratory esophageal pressure versus lower inflection point in acute lung injury. <i>Critical Care</i> , 2013 , 17,	10.8	78

458	Lung recruitment induced by Sigh in hypoxemic intubated critically ill patients. <i>Critical Care</i> , 2013 , 17,	10.8	78
457	Quantification of stress raisers in ARDS. <i>Critical Care</i> , 2013 , 17,	10.8	4
456	Recruited lung tissue does not resume normal mechanical properties. Critical Care, 2013, 17,	10.8	2
455	Intraoperative management of hypoxemia with recruitment maneuvers: are the benefits worth the costs?. <i>Critical Care</i> , 2013 , 17,	10.8	78
454	Haemodynamic changes during alveolar recruitment manoeuvre in patients with diastolic dysfunction. <i>Critical Care</i> , 2013 , 17,	10.8	78
453	Alveolar Recruitment for ARDS Trial: preliminary results. <i>Critical Care</i> , 2013 , 17,	10.8	1
452	Reliability of pressure measurements via balloon catheters is high: an evaluation of six esophageal catheters. <i>Critical Care</i> , 2013 , 17,	10.8	78
451	Effect of protective ventilation on organ-specific cytokine production in an experimental postoperative sepsis model. <i>Critical Care</i> , 2013 , 17,	10.8	78
450	Time required for gas exchange equilibration after a change of positive end-expiratory pressure in acute respiratory distress syndrome. <i>Critical Care</i> , 2013 , 17,	10.8	78
449	Comparison between the standard and low-dose chest CT scans on the lung quantitative analysis in critically ill patients. <i>Critical Care</i> , 2013 , 17,	10.8	78
448	Preliminary experience with airway pressure release ventilation on hemodynamics in patients with septic shock in a medical/surgical ICU. <i>Critical Care</i> , 2013 , 17,	10.8	78
447	Feasibility and effectiveness of prone position in morbidly obese ARDS patients: a case-control clinical study. <i>Critical Care</i> , 2013 , 17,	10.8	78
446	Provision of mechanical ventilation to pregnant/postpartum women with H1N1 influenza: a case-control study. <i>Critical Care</i> , 2013 , 17,	10.8	78
445	Inhalation injury in critical burn patients. <i>Critical Care</i> , 2013 , 17,	10.8	78
444	Beneficial effects of intrapulmonary percussive ventilation in patients with respiratory insufficiency in the ICU. <i>Critical Care</i> , 2013 , 17,	10.8	1
443	High-frequency ventilation for acute traumatic and nontraumatic lung injury. Critical Care, 2013, 17,	10.8	78
442	Improvement in oxygenation with high-frequency oscillatory ventilation combined with tracheal gas insufflation is correlated to the extravascular lung water index. <i>Critical Care</i> , 2013 , 17,	10.8	78
441	Physiologic effects and regional ventilation of high-frequency positive-pressure ventilation using a conventional ventilator in a severe ARDS animal model associated with an inspiratory pause or recruitment maneuvers. <i>Critical Care</i> , 2013 , 17,	10.8	78

440	High-frequency percussive ventilation at altitude: study in a hypobaric chamber with a mechanical test lung. <i>Critical Care</i> , 2013 , 17,	10.8	78
439	Recruitable volume is comparable in acute respiratory distress syndrome and in healthy lungs. <i>Critical Care</i> , 2013 , 17,	10.8	78
438	Nosocomial bloodstream infection and venovenous extracorporeal membrane oxygenation: a retrospective cohort study. <i>Critical Care</i> , 2013 , 17,	10.8	2
437	Monitoring anticoagulation during extracorporeal membrane oxygenation in patients with acute respiratory failure. <i>Critical Care</i> , 2013 , 17,	10.8	3
436	Extracorporeal membrane oxygenation: the MOTOR of cytokine production?. <i>Critical Care</i> , 2013 , 17,	10.8	78
435	Effect of extracorporeal CO2 removal on respiratory rate in spontaneously breathing patients with chronic obstructive pulmonary disease exacerbation. <i>Critical Care</i> , 2013 , 17,	10.8	8
434	ECCO2 removal with a phosphorylcholine-coated membrane oxygenator in difficult respiratory weaning patients. <i>Critical Care</i> , 2013 , 17,	10.8	2
433	Lung elastance can be determined without esophageal pressure measurements. <i>Critical Care</i> , 2013 , 17,	10.8	1
432	Feasibility of minimally invasive stimulation of the phrenic nerves for supporting ventilation in fully anesthetized swine. <i>Critical Care</i> , 2013 , 17,	10.8	78
431	Prehospital management of COPD patients in respiratory failure and short-term outcome. <i>Critical Care</i> , 2013 , 17,	10.8	78
430	N-terminal pro-brain natriuretic peptide as a prognostic marker in patients with exacerbation of chronic obstructive pulmonary disease. <i>Critical Care</i> , 2013 , 17,	10.8	78
429	Novel biomarkers for prediction of mortality after acute exacerbation of chronic obstructive pulmonary disease. <i>Critical Care</i> , 2013 , 17,	10.8	78
428	High incidence of respiratory viruses in critically ill adult patients with respiratory failure. <i>Critical Care</i> , 2013 , 17,	10.8	78
427	Role of bronchoscopy during non-invasive ventilation in hypercapnic respiratory failure. <i>Critical Care</i> , 2013 , 17,	10.8	78
426	SMART-COP score for patients admitted with community-acquired pneumonia (CAP) to an ICU in a district general hospital: a smarter way of identifying patients with severe CAP?. <i>Critical Care</i> , 2013 , 17,	10.8	78
425	Effects of steroid therapies to severe streptococcal pneumoniae that required mechanical ventilation. <i>Critical Care</i> , 2013 , 17,	10.8	78
424	Risk factors for ventilator-associated pneumonia in a surgical ICU. Critical Care, 2013, 17,	10.8	78
423	Ventilation distribution measured with electrical impedance tomography at varying levels of assist during pressure support versus neurally adjusted ventilatory assist in ICU patients. <i>Critical Care</i> , 2013 , 17,	10.8	78

422	Analysis of bilevel positive airway pressure therapy in children based on weight. <i>Critical Care</i> , 2013 , 17,	10.8	78
421	Determinants of non-invasive ventilation success or failure in morbidly obese patients in acute respiratory failure. <i>Critical Care</i> , 2013 , 17,	10.8	78
420	Non-invasive ventilation in patients with a clinical diagnosis of pneumonia: a clinical audit. <i>Critical Care</i> , 2013 , 17,	10.8	78
419	Analysis of management of non-invasive ventilation support in prehospital care for COPD patients and short-term outcome. <i>Critical Care</i> , 2013 , 17,	10.8	1
418	Retrospective review of utilisation and outcomes of diaphragmatic EMG monitoring and neurally adjusted ventilatory assist in a central London teaching hospital over a 3-year period. <i>Critical Care</i> , 2013 , 17,	10.8	78
417	Validation of a lung ultrasound protocol in acute respiratory failure: preliminary results. <i>Critical Care</i> , 2013 , 17,	10.8	2
416	Risk of weaning failure with PAOP 🛭 5 versus PAOP 🗗 8. Critical Care, 2013, 17,	10.8	O
415	Impact of continuous positive airway pressure on the treatment of acute exacerbation of asthma: a randomised controlled trial. <i>Critical Care</i> , 2013 , 17,	10.8	1
414	Novel method to confirm tracheal intubation based on airway pressures: validation in patients with lung disease. <i>Critical Care</i> , 2013 , 17,	10.8	78
413	Main indications, diagnostic and therapeutic yield of bronchoscopy in the ICU. <i>Critical Care</i> , 2013 , 17,	10.8	78
412	Benchtop study comparing leakages across cuffs of three endotracheal tubes. <i>Critical Care</i> , 2013 , 17,	10.8	1
411	Endotracheal tube with tapered-type cuff for preventing ventilator-associated pneumonia: a randomized clinical trial. <i>Critical Care</i> , 2013 , 17,	10.8	1
410	Effect of body position changes on endotracheal tube cuff pressure in ventilated patients. <i>Critical Care</i> , 2013 , 17,	10.8	78
409	Towards safer airway management in the critically ill: lessons from National Audit Project 4. <i>Critical Care</i> , 2013 , 17,	10.8	78
408	Quantitative assessment of endotracheal tube (ETT) biofilm by micro-CT scan: evaluation of the effectiveness of ETT cleaning devices. <i>Critical Care</i> , 2013 , 17,	10.8	78
407	Early identification of patients at risk of difficult intubation in the ICU: development and validation of the MACOCHA score in a multicenter cohort study. <i>Critical Care</i> , 2013 , 17,	10.8	O
406	Incidence of major complications related to endotracheal intubation in ICUs of nonacademic Brazilian public hospitals. <i>Critical Care</i> , 2013 , 17,	10.8	78
405	Tracheal intubation for a difficult airway using Airway scope[] , KingVision[] and McGRATH[] : a comparative manikin study of inexperienced personnel. <i>Critical Care</i> , 2013 , 17,	10.8	4

404	Resistance of neonatal endotracheal tubes: a comparison of four commercially available types. <i>Critical Care</i> , 2013 , 17,	10.8	2
403	NT-proBNP and cardiac cycle efficiency changes during extubation process in critically ill patients. <i>Critical Care</i> , 2013 , 17,	10.8	78
402	Comparison of outcomes between early and late tracheostomy for critically ill patients. <i>Critical Care</i> , 2013 , 17,	10.8	3
401	Ultrasound scanning for percutaneous dilatational tracheostomy: a systematic review. <i>Critical Care</i> , 2013 , 17,	10.8	78
400	Evaluation of a new device for emergency transcricoid ventilation in a manikin model. <i>Critical Care</i> , 2013 , 17,	10.8	2
399	Device for measuring bronchodilator delivery and response in resource-limited settings. <i>Critical Care</i> , 2013 , 17,	10.8	78
398	Evidence-based guidelines and protocols for the management of adult patients with a tracheostomy: a systematic review. <i>Critical Care</i> , 2013 , 17,	10.8	78
397	Survey of emergency tracheostomy management in the East of England region. <i>Critical Care</i> , 2013 , 17,	10.8	78
396	Can we keep neck breathers safe? A survey on training adequacy of medical staff caring for tracheostomy/laryngectomy patients. <i>Critical Care</i> , 2013 , 17,	10.8	78
395	Laryngeal mask as a safe and effective ventilatory device during Blue Dolphin tracheostomy in the ICU. <i>Critical Care</i> , 2013 , 17,	10.8	78
394	Real-time ultrasound-guided balloon dilatational percutaneous tracheostomy is a safe procedure in critically ill patients: an evaluation study. <i>Critical Care</i> , 2013 , 17,	10.8	16
393	Right ventricular restriction in interventional lung assist for acute respiratory distress syndrome. <i>Critical Care</i> , 2013 , 17,	10.8	78
392	Left ventricular electromechanical dyssynchrony and mortality in cardiothoracic intensive care. <i>Critical Care</i> , 2013 , 17,	10.8	78
391	Safety of ultrasound-guided central venous access in critically ill patients with uncorrected coagulopathy. <i>Critical Care</i> , 2013 , 17,	10.8	78
390	Cyanoacrylate glue prevents early bleeding of the exit site after CVC or PICC placement. <i>Critical Care</i> , 2013 , 17,	10.8	7
389	Intracardiac ECG for confirmation of correct positioning of central venous catheters is safe and cost-effective. <i>Critical Care</i> , 2013 , 17,	10.8	78
388	Is automated brachial cuff measurement of arterial pressure less accurate in cases of arrhythmia?. <i>Critical Care</i> , 2013 , 17,	10.8	78
387	Cardiac output monitoring in brain-stem-dead potential organ donors: an audit of current UK practice. <i>Critical Care</i> , 2013 , 17,	10.8	78

386	Indocyanine green plasma disappearance rate for assessment of liver function: re-evaluation of normal ranges and impact of biometric data. <i>Critical Care</i> , 2013 , 17,	10.8	2
385	Prediction of 28-day mortality by indocyanine green disappearance rate, other markers of hepatic function and transpulmonary thermodilution parameters: a prospective study in 154 patients. <i>Critical Care</i> , 2013 , 17,	10.8	78
384	Mixed and central venous oxygen saturation are not interchangeable in patients with cardiogenic shock after cardiac surgery. <i>Critical Care</i> , 2013 , 17,	10.8	78
383	Pulmonary hemodynamic disorder in pediatric sepsis and their correction with L-arginine infusion. <i>Critical Care</i> , 2013 , 17,	10.8	78
382	Utility of transesophageal echocardiography in the ICU: a preliminary US perspective. <i>Critical Care</i> , 2013 , 17,	10.8	78
381	Cardiopulmonary monitoring in Thai ICUs: results of ICU-RESOURCE I surveys. <i>Critical Care</i> , 2013 , 17,	10.8	O
380	Establishing a critical care echocardiography laboratory. Critical Care, 2013, 17,	10.8	78
379	Contrast-enhanced ultrasonography in the ICU: promising tool or exciting toy?. <i>Critical Care</i> , 2013 , 17,	10.8	1
378	Hemodynamic disclosure of septic shock patients by intensive care ultrasound. <i>Critical Care</i> , 2013 , 17,	10.8	78
377	Influence of positive end-expiratory pressure on four-chamber longitudinal strain analysis by speckle tracking echocardiography. <i>Critical Care</i> , 2013 , 17,	10.8	78
376	Evaluation of a new calibration index suggesting recalibration of the pulse contour cardiac index by transpulmonary thermodilution: a prospective study. <i>Critical Care</i> , 2013 , 17,	10.8	78
375	Validation of less-invasive hemodynamic monitoring with Pulsioflex in critically ill patients: interim results of a multicentre study. <i>Critical Care</i> , 2013 , 17,	10.8	1
374	Interim results of an ongoing study on the use of non-invasive hemodynamic monitoring with Nexfin in critically ill patients. <i>Critical Care</i> , 2013 , 17,	10.8	78
373	Cost-effectiveness analysis of stroke volume variation guided perioperative hemodynamic optimization. <i>Critical Care</i> , 2013 , 17,	10.8	78
372	EV1000/VolumeView: a new device for a more reliable measurement of extravascular lung water index in patients with lung resections. <i>Critical Care</i> , 2013 , 17,	10.8	78
371	Goal-directed resuscitation therapy in high-risk patients undergoing cardiac surgery (GRICS study): a randomized controlled trial - preliminary results. <i>Critical Care</i> , 2013 , 17,	10.8	1
370	Perioperative volume management for esophageal cancer surgery. Critical Care, 2013, 17,	10.8	78
369	Treatment-related cardiac complications associated with goal-directed therapy in high-risk surgical patients: a meta-analysis. <i>Critical Care</i> , 2013 , 17,	10.8	2

368	Estimation of potential cost-savings related to the implementation of perioperative hemodynamic goal-directed therapy. <i>Critical Care</i> , 2013 , 17,	10.8	78
367	Cardiac output estimation: Vigileo and Mostcare versus echocardiography. Critical Care, 2013, 17,	10.8	78
366	Brachial pulse waveform characteristics predict development of organ failure in septic patients. <i>Critical Care</i> , 2013 , 17,	10.8	78
365	Computerized decision support system improves consistency of haemodynamic assessment amongst ICU team members. <i>Critical Care</i> , 2013 , 17,	10.8	78
364	Volume quantification by contrast-enhanced ultrasound and thermodilution: an in vitro comparison. <i>Critical Care</i> , 2013 , 17,	10.8	78
363	Successful introduction of cardiac index, fluid responsiveness and oxygen delivery data into the primary survey at a central London major trauma centre and impact on time to CT, fluid resuscitation and disposal. <i>Critical Care</i> , 2013 , 17,	10.8	78
362	Fluid responsiveness in pressure support ventilation: role of asynchrony. Critical Care, 2013, 17,	10.8	1
361	Comparison of pulse pressure variation with radial arterial systolic, diastolic and pulse transit time interval variation in pediatric patients undergoing liver transplantation. <i>Critical Care</i> , 2013 , 17,	10.8	78
360	Mini-fluid challenge: how much fluid and what parameter to use?. Critical Care, 2013, 17,	10.8	1
359	Looking for the inflexion point of the Frank-Starling curve. Critical Care, 2013, 17,	10.8	0
358	Prediction of stroke volume response to fluid bolus in 100 children. <i>Critical Care</i> , 2013 , 17,	10.8	2
357	Microcirculatory changes during goal-directed or mean arterial pressure-guided fluid therapy in abdominal surgery. <i>Critical Care</i> , 2013 , 17,	10.8	78
356	Postoperative imaging of the intestinal microcirculation. Critical Care, 2013, 17,	10.8	1
355	Monitoring microcirculatory blood flow with a new sublingual tonometer in a porcine model of haemorrhagic shock. <i>Critical Care</i> , 2013 , 17,	10.8	78
354	Sublingual microcirculatory changes during transient intra-abdominal hypertension: a study in laparoscopic surgery patients. <i>Critical Care</i> , 2013 , 17,	10.8	78
353	Changes in microcirculatory perfusion during cardiac surgery are paralleled by alterations in glycocalyx integrity. <i>Critical Care</i> , 2013 , 17,	10.8	5
352	Microvascular response to cold stress in healthy humans. Critical Care, 2013, 17,	10.8	78
351	Microcirculatory response to experimentally-induced whole body heat stress in healthy humans. <i>Critical Care</i> , 2013 , 17,	10.8	78

350	Observational study of the effects of age, diabetes mellitus, cirrhosis and chronic kidney disease on sublingual microvascular flow. <i>Critical Care</i> , 2013 , 17,	10.8	78
349	Lactate clearance as a determinant of mortality in surgical patients. <i>Critical Care</i> , 2013 , 17,	10.8	78
348	Assessment of the mottling score as a mortality predictor in critically ill patients. <i>Critical Care</i> , 2013 , 17,	10.8	3
347	Impact of different haemodynamic resuscitation strategies on brain perfusion and tissue oedema markers in a model of severe haemorrhagic shock. <i>Critical Care</i> , 2013 , 17,	10.8	78
346	Accuracy of the shock index and various modified shock indexes to predict early mortality in patients suffering from gastrointestinal haemorrhage. <i>Critical Care</i> , 2013 , 17,	10.8	3
345	Inhaled carbon monoxide or nebulized sodium nitrite protect against hemorrhagic shock-induced mitochondrial dysfunction. <i>Critical Care</i> , 2013 , 17,	10.8	78
344	Is norepinephrine more effective than other vasopressors for septic shock? A systematic review and meta-analysis. <i>Critical Care</i> , 2013 , 17,	10.8	78
343	Vasopressin Versus Norepinephrine for the Management of Shock After Cardiac Surgery (VaNCS study): a randomized controlled trial. <i>Critical Care</i> , 2013 , 17,	10.8	1
342	Intraaortic counterpulsation in a second-level institution: indications, management and outcome. <i>Critical Care</i> , 2013 , 17,	10.8	78
341	Impact of left ventricular ejection fraction and elevated filling pressures on mortality in mechanically ventilated patients in severe sepsis or septic shock. <i>Critical Care</i> , 2013 , 17,	10.8	78
340	Effect of Eblockers and calcium channel blockers on shock index predictability in patients suffering from urosepsis. <i>Critical Care</i> , 2013 , 17,	10.8	78
339	Levosimendan in critically ill adults: a utilisation review. Critical Care, 2013, 17,	10.8	78
338	Levosimendan versus dobutamine in Tako-tsubo cardiomyopathy. Critical Care, 2013, 17,	10.8	2
337	Combined effects of ivabradine with dobutamine or levosimendan in isolated perfused hearts. <i>Critical Care</i> , 2013 , 17,	10.8	78
336	Influence of temperature on the cardiac action of dobutamine and levosimendan in isolated perfused hearts. <i>Critical Care</i> , 2013 , 17,	10.8	78
335	Conflicting results between V/Q SPECT and angioscan in pulmonary embolism: what to do?. <i>Critical Care</i> , 2013 , 17,	10.8	78
334	Intraabdominal hypertension and abdominal compartment syndrome in patients admitted to a medical and surgical ICU at a third referral university hospital. <i>Critical Care</i> , 2013 , 17,	10.8	1
333	Red cell distribution width predicts cardiovascular complications after high-risk surgery. <i>Critical Care</i> , 2013 , 17,	10.8	78

332	Consequences of obesity in outcomes after cardiac surgery: analysis of the ARIAM registry. <i>Critical Care</i> , 2013 , 17,	10.8	78
331	Myocardial injury in critically ill patients admitted with noncardiac diagnoses. Critical Care, 2013, 17,	10.8	78
330	Validation of the GRACE score risk in our population. <i>Critical Care</i> , 2013 , 17,	10.8	78
329	Outcomes of incomplete and complete revascularization in ACS. Critical Care, 2013, 17,	10.8	78
328	Non-invasive valve implantation with TAVI versus conventional aortic valve replacement. <i>Critical Care</i> , 2013 , 17,	10.8	78
327	Time to recover from shock is determinant of a positive fluid balance in septic shock. <i>Critical Care</i> , 2013 , 17,	10.8	78
326	Lactate clearance is a predictor of sustained bleeding in emergency room patients with moderate upper gastrointestinal bleeding. <i>Critical Care</i> , 2013 , 17,	10.8	78
325	Muscular glucose assessed by microdialysis and blood glucose can predict mortality in septic shock. <i>Critical Care</i> , 2013 , 17,	10.8	78
324	Constipation in critically ill patients and its relationship to feeding and weaning from respiratory support. <i>Critical Care</i> , 2013 , 17,	10.8	2
323	Nasogastric tube dislodgement: a problem on our ICU. Critical Care, 2013, 17,	10.8	78
322	Handling of dietary protein in critically ill patients. Critical Care, 2013, 17,	10.8	1
321	Confirmation of nasogastric tube placement in critical care. <i>Critical Care</i> , 2013 , 17,	10.8	Ο
320	pH testing to confirm nasogastric tube position on the ICU: are we wasting our time?. <i>Critical Care</i> , 2013 , 17,	10.8	78
319	Effect of probiotic containing lactobacillus, bifidobacterium and streptococcus thermophilus in critically ill patients. <i>Critical Care</i> , 2013 , 17,	10.8	78
318	Selenium levels in patients with different sources of sepsis. Critical Care, 2013, 17,	10.8	1
317	Is a combined i.v. and enteral glutamine regimen superior to a single i.v. glutamine regimen in severe thoracic trauma?. <i>Critical Care</i> , 2013 , 17,	10.8	78
316	Clinical effect of altered lipid emulsion composition containing fish oil in postoperative patients following abdominal surgery: a prospective, randomized, open-label, comparative, multicenter phase 3 clinical study. <i>Critical Care</i> , 2013 , 17,	10.8	1
315	Influence of fat-based versus glucose-based enteral nutrition formulas on glucose homeostasis. <i>Critical Care</i> , 2013 , 17,	10.8	3

314	Effect of hypocaloric versus normocaloric parenteral nutrition on whole body protein kinetics in critically ill neurosurgical patients. <i>Critical Care</i> , 2013 , 17,	10.8	78
313	Withholding parenteral nutrition for 1 week reduces ICU-acquired weakness. Critical Care, 2013, 17,	10.8	2
312	Impact of early parenteral nutrition on muscle and adipose tissue compartments during critical illness. <i>Critical Care</i> , 2013 , 17,	10.8	1
311	Withholding parenteral nutrition during the first week of critical illness increases plasma bilirubin but lowers the incidence of cholestasis and gallbladder sludge. <i>Critical Care</i> , 2013 , 17,	10.8	O
310	Impact of early parenteral nutrition on catabolism. Critical Care, 2013, 17,	10.8	78
309	Impact of early versus late parenteral nutrition on morphological and molecular markers of atrophy and autophagy in skeletal muscle of critically ill patients. <i>Critical Care</i> , 2013 , 17,	10.8	1
308	Impact of closing an emergency department on a neighbouring teaching hospital: the concentrate effect. <i>Critical Care</i> , 2013 , 17,	10.8	78
307	Overcrowding estimation in the emergency department: is the simplest score the best?. <i>Critical Care</i> , 2013 , 17,	10.8	2
306	GP-led walk-in centre in the UK: another way for urgent healthcare provision. <i>Critical Care</i> , 2013 , 17,	10.8	2
305	Establishing a new emergency department: effects on patient flow. Critical Care, 2013, 17,	10.8	78
304	Prehospital EKG evaluation in Rio de Janeiro ambulances. <i>Critical Care</i> , 2013 , 17,	10.8	78
303	Technique of ultrasound-guided peripheral venous access in the emergency room. <i>Critical Care</i> , 2013 , 17,	10.8	78
302	FIB the fractured femur. <i>Critical Care</i> , 2013 , 17,	10.8	78
301	Abdominal pain in adolescent females: a single-centre audit and review of management. <i>Critical Care</i> , 2013 , 17,	10.8	78
300	Multidisciplinary approach to improving documentation of visual acuity in patients presenting with ocular trauma. <i>Critical Care</i> , 2013 , 17,	10.8	78
299	BiPAP for treating moderate and severe asthma exacerbations in a PED. <i>Critical Care</i> , 2013 , 17,	10.8	78
298	Report of an outbreak of toxicity from a novel drug of abuse in the UK: Eric-3. <i>Critical Care</i> , 2013 , 17,	10.8	78
297	Abstract withdrawn. <i>Critical Care</i> , 2013 , 17,	10.8	78

296	Thallium group poisoning incident in Japan 2011. <i>Critical Care</i> , 2013 , 17,	10.8	1
295	Heart rate variability in children with tricyclic antidepressant intoxication. <i>Critical Care</i> , 2013 , 17,	10.8	78
294	Diagnosing anisakiasis in the emergency department. Critical Care, 2013, 17,	10.8	1
293	Implementation of a sepsis protocol in a community hospital using a Telemedicine Program. <i>Critical Care</i> , 2013 , 17,	10.8	1
292	Initial Brazilian experience of Telestroke for thrombolysis in a community hospital. <i>Critical Care</i> , 2013 , 17,	10.8	1
291	Analysis of emergency calls achieved in a French emergency dispatching centre: what resources for which patients. <i>Critical Care</i> , 2013 , 17,	10.8	78
290	Reform of emergency services: immediate effects on cardiac care unit and ICU patient intake. <i>Critical Care</i> , 2013 , 17,	10.8	78
289	Overtriage and undertriage in a prehospital system over 7 years. <i>Critical Care</i> , 2013 , 17,	10.8	1
288	Lorca@hospital evacuation and medical attention by the emergency department. <i>Critical Care</i> , 2013 , 17,	10.8	78
287	War injuries and civilian accidents in Afghan children. <i>Critical Care</i> , 2013 , 17,	10.8	Ο
286	Intensive care and disaster medicine: the role of a compendium. Critical Care, 2013, 17,	10.8	78
285	Factors involved in ICU mortality following medical retrieval: identifying differences between ICU survivors and nonsurvivors. <i>Critical Care</i> , 2013 , 17,	10.8	78
284	Algorithm for the resuscitation of traumatic cardiac arrest patients in a physician-staffed helicopter emergency medical service. <i>Critical Care</i> , 2013 , 17,	10.8	1
283	Are physicians required during HEMS winch rescue missions?. <i>Critical Care</i> , 2013 , 17,	10.8	78
282	Improving maintenance of critical care land and aeromedical transfer equipment. <i>Critical Care</i> , 2013 , 17,	10.8	78
281	Safety standards for intrahospital transfer of critical care patients. <i>Critical Care</i> , 2013 , 17,	10.8	78
280	Prehospital hyperoxemia does not influence the functional neurological outcome in polytraumatized patients with traumatic head injury. <i>Critical Care</i> , 2013 , 17,	10.8	78
279	Innate immune response-mediated late increase in SuPAR in multi-trauma patients. <i>Critical Care</i> , 2013 , 17,	10.8	78

278	Single-dose estrogen infusion can amplify brain levels of Sonic hedgehog, a signal protein for neuro stem cells and repair following the indirect brain injury resulting after severe torso burns. <i>Critical Care</i> , 2013 , 17,	10.8	2
277	Feasibility and experience of using exception from informed consent in a pilot study of immediate estrogen infusion for hypotensive trauma patients. <i>Critical Care</i> , 2013 , 17,	10.8	78
276	Early administration of a single dose of parenteral estrogen decreases inflammation in the heart for 45 days after severe burns. <i>Critical Care</i> , 2013 , 17,	10.8	78
275	Accuracy of FAST scan in blunt abdominal trauma in a major London trauma centre. <i>Critical Care</i> , 2013 , 17,	10.8	5
274	Level of lipopolysaccharide-binding protein and microbial landscape with account of severity of sepsis syndromes in polytrauma. <i>Critical Care</i> , 2013 , 17,	10.8	78
273	Risk factors for the development of complications following blunt chest trauma: a new risk stratification tool. <i>Critical Care</i> , 2013 , 17,	10.8	O
272	Guidelines for chest drain insertion do not protect relevant anatomical structures. <i>Critical Care</i> , 2013 , 17,	10.8	78
271	Prediction score of indications for whole body computed tomography in blunt trauma patients. <i>Critical Care</i> , 2013 , 17,	10.8	78
270	Prehospital blood transfusion: 5-year experience of an Australian helicopter emergency medical service. <i>Critical Care</i> , 2013 , 17,	10.8	15
269	Severe hyperkalemia is prevalent and can be detected by quantitative electrocardiographic findings and medication history of calcium channel blocker among the patients with symptomatic or extreme bradycardia. <i>Critical Care</i> , 2013 , 17,	10.8	1
268	Prehospital risk stratification following out-of-hospital cardiac arrest. Critical Care, 2013, 17,	10.8	78
267	Multidisciplinary Medical Emergency Team System increases survival in cancer patients undergoing cardiopulmonary resuscitation. <i>Critical Care</i> , 2013 , 17,	10.8	78
266	Focused cardiac ultrasound in resuscitation. <i>Critical Care</i> , 2013 , 17,	10.8	78
265	Acute intrathoracic gastric herniation as a rare cause of cardiac arrest. Critical Care, 2013, 17,	10.8	O
264	Implementation of dispatcher-assisted cardiopulmonary instructions using the ALERT protocol: preliminary results in Belgium. <i>Critical Care</i> , 2013 , 17,	10.8	78
263	Prehospital epinephrine administration and survival among patients with unshockable initial rhythm after out-of-hospital cardiac arrest. <i>Critical Care</i> , 2013 , 17,	10.8	78
262	Impact of termination of resuscitation for out-of-hospital cardiopulmonary arrest in Japan. <i>Critical Care</i> , 2013 , 17,	10.8	78
261	Survival from cardiopulmonary arrest after regular working hours in a tertiary-care hospital: retrospective study. <i>Critical Care</i> , 2013 , 17,	10.8	1

260	Cognitive performance and capacity to return home following out-of-hospital cardiac arrest. <i>Critical Care</i> , 2013 , 17,	10.8	78
259	Start value of cerebral saturation in prehospital cardiac arrest patients: does it mean something?. <i>Critical Care</i> , 2013 , 17,	10.8	78
258	Intensive care admission after cardiac arrest: cardiac versus noncardiac causes and consequences for treatment. <i>Critical Care</i> , 2013 , 17,	10.8	78
257	Comparison of 1-year neurological outcome between intra-hospital and extra-hospital cardiac arrest survivors submitted to mild therapeutic hypothermia in a community-based setting in Brazil. <i>Critical Care</i> , 2013 , 17,	10.8	78
256	Primary percutaneous coronary angioplasty and therapeutic hypothermia in out-of-hospital cardiac arrest. <i>Critical Care</i> , 2013 , 17,	10.8	78
255	Infrared pupillometry for outcome prediction after cardiac arrest and therapeutic hypothermia. <i>Critical Care</i> , 2013 , 17,	10.8	78
254	Effect of moderate hyperventilation and induced hypertension on cerebral tissue oxygen saturation in comatose post-cardiac arrest patients treated with hypothermia. <i>Critical Care</i> , 2013 , 17,	10.8	1
253	Effects of viscosity on cerebral blood flow after cardiac arrest. Critical Care, 2013, 17,	10.8	1
252	Therapeutic mild hypothermia after cardiac arrest in shockable and nonshockable rhythms: does it improve both survival and neurological outcome?. <i>Critical Care</i> , 2013 , 17,	10.8	78
251	Agreement between temperature measurements during cooling and warming through the esophagus. <i>Critical Care</i> , 2013 , 17,	10.8	78
250	Association between oxygenation and 6-month mortality during post-cardiac arrest care. <i>Critical Care</i> , 2013 , 17,	10.8	78
249	Neuron-specific enolase and bispectral index/suppression ratio for prognostication after cardiac arrest. <i>Critical Care</i> , 2013 , 17,	10.8	78
248	Motor scores, therapeutic hypothermia and neurological outcome after cardiac arrest. <i>Critical Care</i> , 2013 , 17,	10.8	78
247	Differences of calcium binding protein immunoreactivities in the young hippocampal CA1 region from the adult following transient ischemic damage. <i>Critical Care</i> , 2013 , 17,	10.8	78
246	Early hypothermia improves survival and reduces the rise of serum biomarkers after traumatic brain injury in swine. <i>Critical Care</i> , 2013 , 17,	10.8	78
245	Epidemiology of moderate and severe traumatic brain injury in Cairo University Hospital in 2010. <i>Critical Care</i> , 2013 , 17,	10.8	1
244	Alcohol intoxication impedes the recognition of traumatic brain injury in the prehospital setting and may worsen 6-month outcome. <i>Critical Care</i> , 2013 , 17,	10.8	78
243	Predictive ability of the IMPACT-TBI prognostic model is improved by adding markers of coagulation. <i>Critical Care</i> , 2013 , 17,	10.8	78

242	Nonischemic endogenous lactate production in humans with severe traumatic brain injury. <i>Critical Care</i> , 2013 , 17,	10.8	78
241	Japan Coma Scale used in the prehospital setting can predict clinical outcome in severe pediatric trauma. <i>Critical Care</i> , 2013 , 17,	10.8	10
240	Effect of positive end-expiratory pressure on ultrasound measurement of optical sheath nerve diameter: preliminary study. <i>Critical Care</i> , 2013 , 17,	10.8	78
239	National survey of current protocols and management of the traumatic brain injury patients in UK ICUs. <i>Critical Care</i> , 2013 , 17,	10.8	78
238	Low-frequency autoregulation index for calculation of optimal cerebral perfusion pressure in severe traumatic brain injury. <i>Critical Care</i> , 2013 , 17,	10.8	1
237	Cerebral rSO2 monitoring in pediatric altered mental status patients. <i>Critical Care</i> , 2013 , 17,	10.8	78
236	Intravenous paracetamol for fever control in acute brain-injured patients: cerebral and hemodynamic effects. <i>Critical Care</i> , 2013 , 17,	10.8	2
235	Retrospective analysis of the hemodynamic effects of induction of barbiturate coma in patients with refractory elevated intracranial pressure. <i>Critical Care</i> , 2013 , 17,	10.8	78
234	Hospital readmissions following traumatic brain injury. Critical Care, 2013, 17,	10.8	4
233	Risk factors of pituitary disorders following traumatic brain injury. <i>Critical Care</i> , 2013 , 17,	10.8	78
232	Vasopressin reduces the cumulative epinephrine dose in hypothermic swine with traumatic brain injury. <i>Critical Care</i> , 2013 , 17,	10.8	78
231	Long-term functional outcome in adults with severe TBI: a meta-analysis. Critical Care, 2013, 17,	10.8	Ο
230	Monitoring of severe traumatic brain injury patients in UK ICUs: a national survey. <i>Critical Care</i> , 2013 , 17,	10.8	1
229	Survival analysis after head injury: is a normal INR reassuring?. Critical Care, 2013, 17,	10.8	1
228	One-year extended Glasgow Outcome Scale and hospital mortality predictors in patients with severe traumatic brain injury in Brazil. <i>Critical Care</i> , 2013 , 17,	10.8	78
227	Early respiratory deterioration predicts poor outcome after aneurysmal subarachnoid hemorrhage. <i>Critical Care</i> , 2013 , 17,	10.8	78
226	Evaluation of proximal-distal velocity gradient in spastic middle cerebral artery after aneurismal subarachnoid hemorrhage. <i>Critical Care</i> , 2013 , 17,	10.8	78
225	Microparticles in aneurismal subarachnoid hemorrhage: role in acute and delayed cerebral ischemia. <i>Critical Care</i> , 2013 , 17,	10.8	78

224	Intracranial pressure after subarachnoid hemorrhage. Critical Care, 2013, 17,	10.8	78
223	Prevalence of pituitary disorders associated with aneurysmal subarachnoid hemorrhage. <i>Critical Care</i> , 2013 , 17,	10.8	78
222	Correlation between the timing of endovascular coiling or microsurgical clipping and long-term outcomes of patients after aneurysmal subarachnoid haemorrhage. <i>Critical Care</i> , 2013 , 17,	10.8	78
221	Time course of neuropsychological functioning after aneurysmal subarachnoid hemorrhage. <i>Critical Care</i> , 2013 , 17,	10.8	Ο
220	Neuroprotective effect of a new synthetic aspirin-decursinol adduct in a rat model of ischemic stroke. <i>Critical Care</i> , 2013 , 17,	10.8	78
219	Prevalence of acute critical neurological disease in children: a global epidemiological assessment (PANGEA). <i>Critical Care</i> , 2013 , 17,	10.8	4
218	Survival analysis of patients taking dabigatran who consulted an emergency department for acute bleeding: a first alarm. <i>Critical Care</i> , 2013 , 17,	10.8	78
217	Bedside anti-Xa measurement for therapeutic assessment of a prophylactic anticoagulation regimen. <i>Critical Care</i> , 2013 , 17,	10.8	78
216	Effective dosage of enoxaparin for intensive care patients. Critical Care, 2013, 17,	10.8	78
215	Anti-Xa activities following 5,000 IU and 7,500 IU of s.c. dalteparin in critically ill patients and 5,000 IU in medical patients: a prospective randomized study. <i>Critical Care</i> , 2013 , 17,	10.8	78
214	Randomized double-blind placebo-controlled PK/PD study on the effects of a single intravenous dose of the anti-hepcidin Spiegelmer NOX-H94 on serum iron during experimental human endotoxemia. <i>Critical Care</i> , 2013 , 17,	10.8	6
213	Haemoglobin concentration on admission to intensive care influences hospital mortality rates and length of stay: a retrospective study. <i>Critical Care</i> , 2013 , 17,	10.8	78
212	Influence of epidural anesthesia on the hemocoagulation disorders and quantity of septic complications in patients with acute necrotizing pancreatitis. <i>Critical Care</i> , 2013 , 17,	10.8	О
211	Hemoviscoelastography: a new method of hemostasis monitoring. Critical Care, 2013, 17,	10.8	78
210	Thromboelastographic monitoring of fondaparinux in surgical patients. Critical Care, 2013, 17,	10.8	2
209	Urokinase-induced fibrinolysis in thromboelastography (UKIF-TEG) to assess fibrinolysis in critically ill patients. <i>Critical Care</i> , 2013 , 17,	10.8	78
208	Bleeding complications after central line insertions and the relevance of pre-procedure coagulation tests and blood component therapy. <i>Critical Care</i> , 2013 , 17,	10.8	0
207	Efficacy of tranexamic acid in decreasing blood loss during cesarean section. <i>Critical Care</i> , 2013 , 17,	10.8	78

206	Dose-response of fibrinogen and factor XIII concentrate for correcting albumin-induced coagulopathy. <i>Critical Care</i> , 2013 , 17,	10.8	78
205	Intraoperative cell savage for life-threatening haemorrhagic caesarean section. <i>Critical Care</i> , 2013 , 17,	10.8	78
204	Does protamine infusion reduce postoperative blood loss and heparin rebound in cardiac surgery patients? A retrospective analysis of 240 patients on the cardiac ICU. <i>Critical Care</i> , 2013 , 17,	10.8	78
203	Transfusion Requirements in Surgical Oncology Patients (TRISOP): a randomized, controlled trial. <i>Critical Care</i> , 2013 , 17,	10.8	2
202	Red blood cell transfusion is a predictor of mortality and morbidity in children undergoing cardiac surgery. <i>Critical Care</i> , 2013 , 17,	10.8	78
201	Red blood cell transfusion results in worse outcomes in patients with left ventricular dysfunction undergoing on-pump cardiac surgery. <i>Critical Care</i> , 2013 , 17,	10.8	78
200	Defining the need for a massive transfusion in severe blunt traumatic patients. <i>Critical Care</i> , 2013 , 17,	10.8	1
199	Association between red blood cell transfusions and clinical outcome in critically ill children. <i>Critical Care</i> , 2013 , 17,	10.8	78
198	Microcirculation and blood transfusions during sepsis: leukoreduced (LR) versus non-LR red blood cells. <i>Critical Care</i> , 2013 , 17,	10.8	78
197	Massive blood transfusion for obstetric haemorrhage. <i>Critical Care</i> , 2013 , 17,	10.8	78
196	Blood transfusion practices in the ICU of a level 1 trauma centre and tertiary cardiac unit. <i>Critical Care</i> , 2013 , 17,	10.8	1
195	C1 inhibitor attenuates pulmonary inflammation in an in vivo model of transfusion-related acute lung injury. <i>Critical Care</i> , 2013 , 17,	10.8	78
194	Incidence, risk factors and outcome of TRALI after cardiac surgery. Critical Care, 2013, 17,	10.8	2
193	Albumin-induced coagulopathy is less severe and more effectively reversed with fibrinogen concentrate than is synthetic colloid-induced coagulopathy. <i>Critical Care</i> , 2013 , 17,	10.8	78
192	Network meta-analysis of clinical trials of fluid treatments in sepsis demonstrates improved survival with albumin compared with crystalloid and hydroxyethyl starch. <i>Critical Care</i> , 2013 , 17,	10.8	78
191	Assessing the economic impact of catheter-related bloodstream infections when switching from open to closed system delivery of human albumin solutions in ICUs. <i>Critical Care</i> , 2013 , 17,	10.8	78
190	Comparing the hemodynamic effects of six different colloids. Critical Care, 2013, 17,	10.8	78
189	Strong ion difference and arterial bicarbonate concentration as cornerstones of the impact of fluid therapy on acid-base balance. <i>Critical Care</i> , 2013 , 17,	10.8	78

188	Microcirculatory response to fluid challenge: should we prefer balanced colloids to rebalance tissue perfusion?. <i>Critical Care</i> , 2013 , 17,	10.8	78
187	Procedural sedation in the emergency department. Critical Care, 2013, 17,	10.8	78
186	Impact of an age, kidney and liver function adjusted sedation protocol in critically ill patients. <i>Critical Care</i> , 2013 , 17,	10.8	78
185	Risk and benefit of protocol sedation and daily interruption of sedation for mechanically ventilated patients: a systematic review. <i>Critical Care</i> , 2013 , 17,	10.8	78
184	Chemical and physical compatibility of continuous intravenous drug infusion combinations used in paediatric intensive care. <i>Critical Care</i> , 2013 , 17,	10.8	78
183	Abstract withdrawn. <i>Critical Care</i> , 2013 , 17,	10.8	78
182	Benzodiazepines induce hyperglycemia in rats by affecting peripheral disposal of glucose. <i>Critical Care</i> , 2013 , 17,	10.8	78
181	Prolonged sedation in ARDS patients with inhaled anesthetics: our experience. <i>Critical Care</i> , 2013 , 17,	10.8	2
180	Involvement of mitochondrial ATP-sensitive K+ channels in fentanyl-induced mitochondrial dysfunction of cultured human hepatocytes. <i>Critical Care</i> , 2013 , 17,	10.8	78
179	Effects of clonidine on microcirculatory alterations during endotoxemia. Critical Care, 2013, 17,	10.8	1
178	Conjunctival administration of S+ketamine in rabbits. <i>Critical Care</i> , 2013 , 17,	10.8	78
177	Use of propofol for procedural sedation reduces length of stay in the emergency department. <i>Critical Care</i> , 2013 , 17,	10.8	78
176	Pre-PRIS? Prospective monitoring for early markers of propofol infusion syndrome. <i>Critical Care</i> , 2013 , 17,	10.8	0
175	Safety profile of 1,008 cases of propofol sedation in the emergency department. <i>Critical Care</i> , 2013 , 17,	10.8	78
174	Use of physical restraint in Dutch ICUs: prevalence and motives. Critical Care, 2013, 17,	10.8	78
173	Factors influencing the use of physical restraints in Canadian ICUs. Critical Care, 2013, 17,	10.8	1
172	Current practices in ICU delirium management: a prospective multicenter study in the Netherlands. <i>Critical Care</i> , 2013 , 17,	10.8	78
171	Intraoperative dexamethasone and delirium after cardiac surgery: a randomized clinical trial. <i>Critical Care</i> , 2013 , 17,	10.8	1

170	Current perspectives, beliefs and practices concerning delirium in critically ill patients: a multicenter survey among Dutch healthcare professionals. <i>Critical Care</i> , 2013 , 17,	10.8	1
169	International validation of the delirium prediction model for ICU patients (PREDELIRIC): a multicenter observational study. <i>Critical Care</i> , 2013 , 17,	10.8	1
168	Delirium in ICU patients is associated with increased temperature variability. <i>Critical Care</i> , 2013 , 17,	10.8	1
167	Long-term outcome of delirium in critically ill patients. <i>Critical Care</i> , 2013 , 17,	10.8	1
166	Gastrointestinal symptoms in the ICU: a comparison of primary and secondary acute gastrointestinal injury. <i>Critical Care</i> , 2013 , 17,	10.8	78
165	Effects of proton pump inhibitor versus H2-receptor antagonist stress ulcer prophylaxis on ventilator-associated pneumonia: a pilot study. <i>Critical Care</i> , 2013 , 17,	10.8	4
164	Early diagnosis of liver failure in septic patients using the maximal liver function capacity test (LiMAx test): comparison with conventional methods. <i>Critical Care</i> , 2013 , 17,	10.8	78
163	Impact of liver resection on portal venous pressure and renal function. Critical Care, 2013, 17,	10.8	78
162	Value of the preoperative Model for End-stage Liver Disease score in predicting postoperative outcome in living donor liver transplantation recipients. <i>Critical Care</i> , 2013 , 17,	10.8	78
161	Does just fungal evidence increase the mortality of critically ill patients with liver cirrhosis? A retrospective analysis. <i>Critical Care</i> , 2013 , 17,	10.8	78
160	Predictors of persistent organ failure in ICU patients with severe acute pancreatitis. <i>Critical Care</i> , 2013 , 17,	10.8	78
159	Increased plasma levels of thrombopoietin in patients with severe acute pancreatitis. <i>Critical Care</i> , 2013 , 17,	10.8	1
158	Severe renal dysfunction measured by pRIFLE in children is an independent factor associated with death and not predicted by PIM2. <i>Critical Care</i> , 2013 , 17,	10.8	78
157	Thorough evaluation for the new acute kidney injury criteria by Kidney Disease Improving Global Outcomes. <i>Critical Care</i> , 2013 , 17,	10.8	2
156	Urinary neutrophil gelatinase-associated lipocalin as an early predictor of acute kidney injury in cardiac surgery patients. <i>Critical Care</i> , 2013 , 17,	10.8	0
155	Atrial natriuretic peptide attenuates metabolic acidosis and inflammation of the kidney, lung and heart in a rat model of renal ischemia-reperfusion injury. <i>Critical Care</i> , 2013 , 17,	10.8	78
154	Perioperative sodium bicarbonate to prevent acute kidney injury after cardiac surgery: a multicenter double-blind randomized controlled trial. <i>Critical Care</i> , 2013 , 17,	10.8	78
153	Tubular damage biomarkers linked to inflammation or iron metabolism predict acute kidney injury. <i>Critical Care</i> , 2013 , 17,	10.8	1

152	Renal response and acid-base balance alterations during furosemide administration. <i>Critical Care</i> , 2013 , 17,	10.8	2
151	Red cell distribution width is not a predictor of mortality in acute kidney injury. <i>Critical Care</i> , 2013 , 17,	10.8	78
150	Value of Tc-99m DMSA renal scan in assessing the prognosis and outcome of acute renal failure due to sepsis. <i>Critical Care</i> , 2013 , 17,	10.8	78
149	Estimated glomerular filtration rate based on hospital discharge creatinine may significantly overestimate renal function and underestimate chronic kidney disease in survivors of critical illness. <i>Critical Care</i> , 2013 , 17,	10.8	1
148	Varying models of intra-abdominal hypertension and their effect on renal function in a porcine model. <i>Critical Care</i> , 2013 , 17,	10.8	78
147	Renal resistive index can predict response to furosemide in septic patients with oliguria. <i>Critical Care</i> , 2013 , 17,	10.8	78
146	Urinary chitinase 3-like 1, a novel biomarker for acute kidney injury in adult cardiac intensive care patients: a pilot study. <i>Critical Care</i> , 2013 , 17,	10.8	1
145	Use, type and duration of cardiopulmonary bypass are related to acute kidney injury occurrence and urinary NGAL concentrations. <i>Critical Care</i> , 2013 , 17,	10.8	1
144	Symmetrical dimethylarginine is a more sensitive biomarker of renal dysfunction than creatinine. <i>Critical Care</i> , 2013 , 17,	10.8	8
143	Prognostic value of neutrophil gelatinase-associated lipocalin and transpulmonary thermodilution-derived parameters within 48 hours after admission. <i>Critical Care</i> , 2013 , 17,	10.8	1
142	Assessment of neutrophil gelatinase-associated lipocalin as a biomarker of acute kidney injury in acute heart failure patients. <i>Critical Care</i> , 2013 , 17,	10.8	О
141	Neutrophil gelatinase-associated lipocalin predicts postoperative fluid overload after cardiac surgery. <i>Critical Care</i> , 2013 , 17,	10.8	78
140	Survivors of acute kidney injury requiring renal replacement therapy rarely receive follow-up: identification of an unmet need. <i>Critical Care</i> , 2013 , 17,	10.8	78
139	Aetiology and outcomes for dialysis-dependent acute kidney injury patients on the ICU. <i>Critical Care</i> , 2013 , 17,	10.8	78
138	Efficiency of substitutive renal therapy in the complex intensive care of multiple organ failure in patients with polytrauma. <i>Critical Care</i> , 2013 , 17,	10.8	78
137	Can treatment with the molecular adsorbent recirculation system be the solution for type-1 hepatorenal syndrome?. <i>Critical Care</i> , 2013 , 17,	10.8	78
136	Effect of continuous high-volume hemofiltration on patients with acute respiratory distress syndrome. <i>Critical Care</i> , 2013 , 17,	10.8	2
135	Dialyzer clearance of myoglobin with middle molecule filter and low blood flow CVVHD in patients with rhabdomyolysis-associated acute kidney injury. <i>Critical Care</i> , 2013 , 17,	10.8	78

134	Daily fluid balance predicts hospital mortality in critically ill patients receiving continuous renal replacement therapy. <i>Critical Care</i> , 2013 , 17,	10.8	78
133	Fluid balance and renal outcomes in patients requiring renal replacement therapy in the ICU. <i>Critical Care</i> , 2013 , 17,	10.8	78
132	Pharmacokinetics and pharmacodynamics of esomeprazole and ranitidine during continuous venovenous hemodiafiltration. <i>Critical Care</i> , 2013 , 17,	10.8	78
131	Pharmacokinetics of gentamicin and vancomycin during continuous venovenous hemofiltration in critically ill septic patients with acute kidney injury. <i>Critical Care</i> , 2013 , 17,	10.8	1
130	Effects of two different regional citrate anticoagulation CVVH protocols on acid-base status and phosphate supplementation. <i>Critical Care</i> , 2013 , 17,	10.8	78
129	Abstract withdrawn. <i>Critical Care</i> , 2013 , 17,	10.8	78
128	Muscle wasting in the ICU can be reliably monitored using ultrasound. Critical Care, 2013, 17,	10.8	78
127	Critical illness induces nutrient-independent adipogenesis and accumulation of alternatively activated tissue macrophages. <i>Critical Care</i> , 2013 , 17,	10.8	78
126	Abstract withdrawn. <i>Critical Care</i> , 2013 , 17,	10.8	78
125	Hypocalcemia during magnesium therapy for obstetric ICU admissions. <i>Critical Care</i> , 2013 , 17,	10.8	O
124	Hyponatremia and intensive care outcome. <i>Critical Care</i> , 2013 , 17,	10.8	78
123	Strong ion gap can be accurately estimated with a simple bedside equation. <i>Critical Care</i> , 2013 , 17,	10.8	78
122	Relative adrenal insufficiency in burns. <i>Critical Care</i> , 2013 , 17,	10.8	78
121	Albumin-adjusted calcium concentration should not be used to identify hypocalcaemia in critical illness. <i>Critical Care</i> , 2013 , 17,	10.8	2
120	Vitamin D status and overall and sepsis mortality in critically ill patients. Critical Care, 2013, 17,	10.8	2
119	Etiology of low testosterone levels in male patients with severe sepsis requiring mechanical ventilation. <i>Critical Care</i> , 2013 , 17,	10.8	4
118	Efficacy of calcium replacement in hypocalcaemia. <i>Critical Care</i> , 2013 , 17,	10.8	78
117	Endogenous melatonin in critically ill patients. <i>Critical Care</i> , 2013 , 17,	10.8	78

116	Succinate ameliorates mitochondrial oxygen consumption of metformin-intoxicated human platelets. <i>Critical Care</i> , 2013 , 17,	10.8	78
115	Does chronic metformin therapy offer cardiomyoprotection in coronary artery bypass grafting surgery patients? A case-control study. <i>Critical Care</i> , 2013 , 17,	10.8	78
114	Metformin-associated lactic acidosis: risk factors and prognostic factors. <i>Critical Care</i> , 2013 , 17,	10.8	1
113	Multiscale modeling of acute insulin resistance in critical care. <i>Critical Care</i> , 2013 , 17,	10.8	78
112	Impact of a real-time electronic persistent hyperglycemia alert on glucose control in the ICU. <i>Critical Care</i> , 2013 , 17,	10.8	1
111	Individualized targeted glucose control to avoid hypoglycemia. Critical Care, 2013, 17,	10.8	78
110	Hyperglycemia in intensive care identifies patients at high risk of developing diabetes: preliminary data of a multicentre study. <i>Critical Care</i> , 2013 , 17,	10.8	78
109	Association between average glucose levels and hospital mortality among critically ill patients. <i>Critical Care</i> , 2013 , 17,	10.8	1
108	Continuous glucose monitoring in critically ill adults: comparison of two different calibration protocols. <i>Critical Care</i> , 2013 , 17,	10.8	1
107	Root cause analysis of hypoglycemic events in critically ill patients. <i>Critical Care</i> , 2013 , 17,	10.8	78
106	Continuous arterial and venous glucose monitoring by quenched chemical fluorescence in ICU patients after cardiac surgery. <i>Critical Care</i> , 2013 , 17,	10.8	2
105	Performance of the Medtronic Sentrino continuous glucose management system in the cardiac ICU. <i>Critical Care</i> , 2013 , 17,	10.8	2
104	Intensive insulin therapy in critically ill children: impact on blood glucose dynamics and its relation with mortality. <i>Critical Care</i> , 2013 , 17,	10.8	78
103	Implementation of evidence-based protocols improves survival: a 15-year surgical ICU experience in 10,172 patients. <i>Critical Care</i> , 2013 , 17,	10.8	78
102	Decubitus ulcers are associated with prolonged length of stay in critically ill patients. <i>Critical Care</i> , 2013 , 17,	10.8	78
101	HIV patients admitted to an ICU of a university hospital - experience of 15 years: 1995 to 2009. <i>Critical Care</i> , 2013 , 17,	10.8	78
100	Prevalence and prognosis of organ dysfunctions of AIDS critically ill patients. <i>Critical Care</i> , 2013 , 17,	10.8	78
99	Mortality predictors in septic HIV patients in the ICU. <i>Critical Care</i> , 2013 , 17,	10.8	78

98	Outcome of intensive care treatment for lung cancer patients. Critical Care, 2013, 17,	10.8	78
97	Outcome of older patients with cancer requiring ICU admission. <i>Critical Care</i> , 2013 , 17,	10.8	78
96	Identifying risk factors for readmission to the ICU: a qualitative approach. Critical Care, 2013, 17,	10.8	78
95	Clinical and laboratory postoperative impact on outcome of living donor liver transplantation recipients. <i>Critical Care</i> , 2013 , 17,	10.8	78
94	Are APACHE II scores better predictors of mortality than routine laboratory values?. <i>Critical Care</i> , 2013 , 17,	10.8	78
93	Complementarity of prognosis models SAPS 3 and EuroSCORE in cardiac surgery. <i>Critical Care</i> , 2013 , 17,	10.8	3
92	Comparing intensive care treated primary, secondary and tertiary sepsis with respect to length of stay, organ failure and mortality: a pilot study. <i>Critical Care</i> , 2013 , 17,	10.8	78
91	Survival of older patients after critical care. <i>Critical Care</i> , 2013 , 17,	10.8	78
90	Eldicus scores for admission to our ICU: a retrospective study. Critical Care, 2013, 17,	10.8	78
89	Developing a screening instrument for predicting psychological morbidity after critical illness. <i>Critical Care</i> , 2013 , 17,	10.8	78
88	Prediction of sequences of organ dysfunction in critical patients studied with Bayesian analysis. <i>Critical Care</i> , 2013 , 17,	10.8	78
87	Comparison of IGS 2 score and a new neural network biomarker score performance in predicting 1-year mortality. <i>Critical Care</i> , 2013 , 17,	10.8	78
86	Significance of preadmission quality of life for mortality in the ICU: a prospective cohort study. <i>Critical Care</i> , 2013 , 17,	10.8	78
85	Predicting physical function and mental health in trauma intensive care patients 2 years after hospitalisation. <i>Critical Care</i> , 2013 , 17,	10.8	78
84	Do ICU patients Odischarge characteristics change with Swiss Diagnosis Related Groups?. <i>Critical Care</i> , 2013 , 17,	10.8	78
83	Do Diagnosis Related Groups change admission practice to a large Swiss ICU?. <i>Critical Care</i> , 2013 , 17,	10.8	1
82	Variable cost of a Greek ICU. <i>Critical Care</i> , 2013 , 17,	10.8	78
81	Audit of the medical staffing of and referrals to a mixed adult ICU in a teaching hospital. <i>Critical Care</i> , 2013 , 17,	10.8	78

80	Impact of the time elapsed between ICU request and actual admission on mortality and length of stay. <i>Critical Care</i> , 2013 , 17,	10.8	78
79	Medical problems: review from the major nuclear incidents. <i>Critical Care</i> , 2013 , 17,	10.8	78
78	Variation in acute care burden and supply across diverse urban settings. Critical Care, 2013, 17,	10.8	3
77	Differences in severity and resource utilization for medical and surgical ICU patients. <i>Critical Care</i> , 2013 , 17,	10.8	О
76	Educational effectiveness of introductory training for the Rapid Response System in Japan. <i>Critical Care</i> , 2013 , 17,	10.8	78
75	Standards of referrals and admissions of critically ill patients to the ICU. Critical Care, 2013, 17,	10.8	78
74	Knowledge of confidentiality, consent and information disclosure is not affected by position or experience in adult critical care. <i>Critical Care</i> , 2013 , 17,	10.8	78
73	Survey of alcohol-related admissions to critical care and the resource and financial implications. <i>Critical Care</i> , 2013 , 17,	10.8	78
72	Association of sleep disordered breathing with clinical trajectories in patients undergoing cardiac surgery. <i>Critical Care</i> , 2013 , 17,	10.8	78
71	Can massage therapy help cardiac patients? A literature review plus a case study. <i>Critical Care</i> , 2013 , 17,	10.8	78
70	Ischemic stroke in different ICU types. <i>Critical Care</i> , 2013 , 17,	10.8	78
69	Development of a severe influenza critical care curriculum and training materials for resource-limited settings. <i>Critical Care</i> , 2013 , 17,	10.8	3
68	Quality assurance in severe sepsis: an individualised audit/feedback system results in substantial improvements at a UK teaching hospital. <i>Critical Care</i> , 2013 , 17,	10.8	О
67	Safety climate perception for staff nurses in an ICU. Critical Care, 2013, 17,	10.8	78
66	Physico-technical medicine: teaching technical skills to the intensivist. Critical Care, 2013, 17,	10.8	78
65	First-hour protocol clarifies resource management and nursing staff education in the ICU. <i>Critical Care</i> , 2013 , 17,	10.8	78
64	Influence of critical care nurse education and work environment on outcomes of mechanically ventilated older adults. <i>Critical Care</i> , 2013 , 17,	10.8	78
63	Activity and case-mix changes in a medical ICU after the geographical transfer of a third-level university hospital. <i>Critical Care</i> , 2013 , 17,	10.8	78

62	Are general wards sufficiently staffed to care for level 1 patients?. Critical Care, 2013, 17,	10.8	78
61	Sleep quality among ICU workers. <i>Critical Care</i> , 2013 , 17,	10.8	78
60	Work-related stress amongst doctors in intensive care, anaesthetics, accident and emergency and general medicine. <i>Critical Care</i> , 2013 , 17,	10.8	78
59	Audit of obstetric critical care admissions in a high-risk population. <i>Critical Care</i> , 2013 , 17,	10.8	78
58	Influence of adverse drug events on the mortality and the length of hospital stay in ICUs in Japan: the JADE Study. <i>Critical Care</i> , 2013 , 17,	10.8	78
57	Attitude and knowledge of intensive care staff concerning donation in Hungary: it is the first step to change. <i>Critical Care</i> , 2013 , 17,	10.8	78
56	Specialist nurse for organ donation in an emergency department will increase organ donation. <i>Critical Care</i> , 2013 , 17,	10.8	78
55	Sepsis in HIV patients admitted to the ICU. <i>Critical Care</i> , 2013 , 17,	10.8	78
54	Mortality related to respiratory failure among pediatric hematology patients requiring intensive care. <i>Critical Care</i> , 2013 , 17,	10.8	78
53	Survival of critically ill patients with haematological malignancies compared with patients without haematological malignancy. <i>Critical Care</i> , 2013 , 17,	10.8	78
52	Outcome of critically ill patients with haematological malignancy admitted to the ICU as an emergency. <i>Critical Care</i> , 2013 , 17,	10.8	78
51	Effect of organizational structure of the ICU on the prognosis: open format versus semi-closed format. <i>Critical Care</i> , 2013 , 17,	10.8	2
50	Use of defaults on an electronic prescribing tool influences the type of fluid received by patients. <i>Critical Care</i> , 2013 , 17,	10.8	78
49	Automated, electronic monitoring for early detection of sepsis. Critical Care, 2013, 17,	10.8	2
48	Admission to the ICU overnight: is it really a bad thing?. Critical Care, 2013, 17,	10.8	78
47	Assessing the quality of interdisciplinary rounds. <i>Critical Care</i> , 2013 , 17,	10.8	1
46	Leadership training and quality improvement of interdisciplinary rounds in the ICU. <i>Critical Care</i> , 2013 , 17,	10.8	1
45	Let@go round again! Quality improvement through intentional rounding. <i>Critical Care</i> , 2013 , 17,	10.8	78

44	Prospective observational study of handover in a medical ICU. Critical Care, 2013, 17,	10.8	78
43	Impact of knowledge transfer through the implementation of a telemedicine program in a community hospital in Brazil. <i>Critical Care</i> , 2013 , 17,	10.8	78
42	Utilization of telemedicine to improve burn care in a developing country. Critical Care, 2013, 17,	10.8	78
41	Different aspects of therapy limitation: a comparative study of the nurse@view. <i>Critical Care</i> , 2013 , 17,	10.8	78
40	Alternative to improve palliative care for all patients and families in critical care units: development and preliminary evaluation following MRC guidance of the King@Psychosocial, Assessment and Care tool. <i>Critical Care</i> , 2013 , 17,	10.8	78
39	End-of life thoughts in the ICU: results of a survey. Critical Care, 2013, 17,	10.8	78
38	End-of-life decisions in Slovenian ICUs: a cross-sectional survey. Critical Care, 2013, 17,	10.8	78
37	Attitudes of intensivists in the UK to withdrawal of futile therapy. Critical Care, 2013, 17,	10.8	78
36	Parent@ perception on end-of-life care in Brazil. Critical Care, 2013, 17,	10.8	78
35	Getting critical care patients home for end-of-life care. <i>Critical Care</i> , 2013 , 17,	10.8	78
34	Project PIX (Post Intensive care eXercise): impact on physical fitness and focus group analysis of quality of life following exercise rehabilitation. <i>Critical Care</i> , 2013 , 17,	10.8	78
33	Feasibility of neuromuscular electrical stimulation in critically ill patients. <i>Critical Care</i> , 2013 , 17,	10.8	1
32	Follow-up and counseling service in trauma patients: needs and goals - a preliminary study. <i>Critical Care</i> , 2013 , 17,	10.8	
31	Post-traumatic stress disorder - are we doing all we can? A UK national-based survey. <i>Critical Care</i> , 2013 , 17,	10.8	78
30	Implementation of an ICU follow-up clinic: outcomes and patient satisfaction after 1 year. <i>Critical Care</i> , 2013 , 17,	10.8	2
29	Acute exercise response in the critically ill. <i>Critical Care</i> , 2013 , 17,	10.8	78
28	Randomized, controlled pilot study of early rehabilitation strategies in acute respiratory failure. <i>Critical Care</i> , 2013 , 17,	10.8	5
27	Early results of a 6-week exercise programme in post-ICU patients. <i>Critical Care</i> , 2013 , 17,	10.8	4

26	Recognition of the primary stressors affecting intensive care patients: a systematic review. <i>Critical Care</i> , 2013 , 17,	10.8	78
25	Quality of care in the ICU from the perspective of relatives. <i>Critical Care</i> , 2013 , 17,	10.8	1
24	Family satisfaction in the ICU: enhancing patient experience. Critical Care, 2013, 17,	10.8	0
23	Family satisfaction with patient care in ICUs of nonacademic Brazilian public hospitals. <i>Critical Care</i> , 2013 , 17,	10.8	78
22	Device-associated infection rates, device use, length of stay, and mortality in intensive care units of 4 Chinese hospitals: International Nosocomial Control Consortium findings. <i>American Journal of Infection Control</i> , 2013 , 41, 301-6	3.8	40
21	Impact of a multidimensional infection control approach on central line-associated bloodstream infections rates in adult intensive care units of 8 cities of Turkey: findings of the International Nosocomial Infection Control Consortium (INICC). Annals of Clinical Microbiology and Antimicrobials,	6.2	42
20	Abstract withdrawn. <i>Critical Care</i> , 2013 , 17,	10.8	78
19	Comparative risk of bloodstream infection in hospitalized patients receiving intravenous medication by open, point-of-care, or closed delivery systems. <i>American Journal of Health-System Pharmacy</i> , 2013 , 70, 957-65	2.2	4
18	In vitro correction of hypothermic and dilutive crystalloid andcolloid rotational thromboelastography-monitored coagulopathywith fi brinogen and factor XIII. <i>Critical Care</i> , 2013 , 17,	10.8	78
17	Erratum to: Urinary neutrophil gelatinase-associated lipocalin as an early predictor of acute kidney injury in cardiac surgery patients. <i>Critical Care</i> , 2013 , 17,	10.8	78
16	Correction: The use, type and duration of cardiopulmonary bypass are related to acute kidney injury occurrence and urinary NGAL concentrations. <i>Critical Care</i> , 2013 , 17, P547	10.8	78
15	Correction: Current perspectives, beliefs and practices concerning delirium in critically ill patients: a multicenter survey among Dutch healthcare professionals. <i>Critical Care</i> , 2013 , 17, P549	10.8	78
14	Closed catheter access system implementation in reducing the bloodstream infection rate in low birth weight preterm infants. <i>Frontiers in Pediatrics</i> , 2015 , 3, 20	3.4	6
13	Evidence-based measures to prevent central line-associated bloodstream infections: a systematic review. <i>Revista Latino-Americana De Enfermagem</i> , 2016 , 24, e2787	1.5	17
12	Eliminating the use of intravenous glass bottles using a FOCUS-PDCA model and providing a practical stability reference guide. <i>International Journal of Pharmacy Practice</i> , 2016 , 24, 271-82	1.7	6
11	Device-Associated Infection Rates in 20 Cities of India, Data Summary for 2004-2013: Findings of the International Nosocomial Infection Control Consortium. <i>Infection Control and Hospital Epidemiology</i> , 2016 , 37, 172-81	2	37
10	Multicenter study in Colombia: Impact of a multidimensional International Nosocomial Infection Control Consortium (INICC) approach on central line-associated bloodstream infection rates. American Journal of Infection Control, 2016, 44, e235-e241	3.8	16
9	International Nosocomial Infection Control Consortium (INICC) resources: INICC multidimensional approach and INICC surveillance online system. <i>American Journal of Infection Control</i> , 2016 , 44, e81-90	3.8	49

CITATION REPORT

8	Prlention von Infektionen, die von Geflathetern ausgehen. <i>Bundesgesundheitsblatt - Gesundheitsforschung - Gesundheitsschutz</i> , 2017 , 60, 171-206	7.5	38
7	Prevention of central venous line associated bloodstream infections in adult intensive care units: A systematic review. <i>Intensive and Critical Care Nursing</i> , 2017 , 43, 12-22	3.1	29
6	Central-line-associated bloodstream infections in a resource-limited South African neonatal intensive care unit. <i>South African Medical Journal</i> , 2017 , 107, 758-762	1.5	9
5	Conflicts of interest in infection prevention and control research: no smoke without fire. A narrative review. <i>Intensive Care Medicine</i> , 2018 , 44, 1679-1690	14.5	3
4	Managing and preventing vascular catheter infections: A position paper of the international society for infectious diseases. <i>International Journal of Infectious Diseases</i> , 2019 , 84, 22-29	10.5	18
3	The 100 top-cited systematic reviews/meta-analyses in central venous catheter research: A PRISMA-compliant systematic literature review and bibliometric analysis. <i>Intensive and Critical Care Nursing</i> , 2020 , 57, 102803	3.1	8
2	Prevention of Central-Line Associated Bloodstream Infections: 2021 [Update. <i>Infectious Disease Clinics of North America</i> , 2021 , 35, 841-856	6.5	2
1	SHEA neonatal intensive care unit (NICU) white paper series: Practical approaches for the prevention of central line-associated bloodstream infections <i>Infection Control and Hospital Epidemiology</i> , 2022 , 1-46	2	Ο