

CITATION REPORT

List of articles citing

Pretreatment diffusion-weighted imaging lesion volume predicts favorable outcome after intravenous thrombolysis with tissue-type plasminogen activator in acute ischemic stroke

DOI: 10.1161/strokeaha.110.600148
Stroke, 2011, 42, 1251-4.

Source: <https://exaly.com/paper-pdf/51639945/citation-report.pdf>

Version: 2024-04-28

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
47	Inhaled Particles and Vapours. <i>Occupational and Environmental Medicine</i> , 1962 , 19, 145-145	2.1	
46	Final infarct volume is a stronger predictor of outcome than recanalization in patients with proximal middle cerebral artery occlusion treated with endovascular therapy. <i>Stroke</i> , 2012 , 43, 3238-44	6.7	140
45	[Industry-funded therapy studies: what is in the pipeline?]. <i>Der Nervenarzt</i> , 2012 , 83, 1260-9	0.5	
44	Outcome of MRI-based intravenous thrombolysis in carotid-T occlusion. <i>Journal of Neurology</i> , 2012 , 259, 2141-6	5.5	8
43	Noninvasive limb remote ischemic preconditioning contributes neuroprotective effects via activation of adenosine A1 receptor and redox status after transient focal cerebral ischemia in rats. <i>Brain Research</i> , 2012 , 1459, 81-90	3.7	99
42	Predictive factors of outcome and hemorrhage after acute ischemic stroke treated by mechanical thrombectomy with a stent-retriever. <i>Neuroradiology</i> , 2013 , 55, 977-987	3.2	60
41	Prediction of recanalization trumps prediction of tissue fate: the penumbra: a dual-edged sword. <i>Stroke</i> , 2013 , 44, 1014-9	6.7	39
40	Stroke size correlates with functional outcome on the simplified modified Rankin Scale questionnaire. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2013 , 22, 781-3	2.8	15
39	Total mismatch in anterior circulation stroke patients before thrombolysis. <i>Journal of Neuroradiology</i> , 2013 , 40, 158-63	3.1	13
38	Use of DWI-only MR protocol for screening stroke mimics. <i>Journal of the Neurological Sciences</i> , 2013 , 328, 37-40	3.2	19
37	Clinical outcomes of fast MRI-based trombolysis in wake-up strokes compared to superacute ischemic strokes within 12 hours. <i>Neurological Research</i> , 2013 , 35, 492-7	2.7	23
36	DWI-ASPECTS as a predictor of dramatic recovery after intravenous recombinant tissue plasminogen activator administration in patients with middle cerebral artery occlusion. <i>Stroke</i> , 2013 , 44, 534-7	6.7	30
35	Leukoaraiosis and sex predict the hyperacute ischemic core volume. <i>Stroke</i> , 2013 , 44, 61-7	6.7	50
34	Microembolism and catheter ablation II: effects of cerebral microemboli injection in a canine model. <i>Circulation: Arrhythmia and Electrophysiology</i> , 2013 , 6, 23-30	6.4	52
33	Early changes in tissue perfusion after tissue plasminogen activator administration in hyperacute ischemic stroke: initial experiences with arterial spin labeling perfusion magnetic resonance imaging. <i>Neurologia Medico-Chirurgica</i> , 2013 , 53, 213-6	2.6	2
32	To do or not to do; dilemma of intra-arterial revascularization in acute ischemic stroke. <i>PLoS ONE</i> , 2014 , 9, e99261	3.7	1
31	High-sensitivity C-reactive protein, lipoprotein-related phospholipase A2, and acute ischemic stroke. <i>Neuropsychiatric Disease and Treatment</i> , 2014 , 10, 1451-7	3.1	22

30	Immediate changes in stroke lesion volumes post thrombolysis predict clinical outcome. <i>Stroke</i> , 2014 , 45, 3275-9	6.7	24
29	The ischemic penumbra: the location rather than the volume of recovery determines outcome. <i>Current Opinion in Neurology</i> , 2014 , 27, 35-41	7.1	21
28	Imaging predictors of outcome following intravenous thrombolysis in acute stroke. <i>Acta Neurologica Belgica</i> , 2014 , 114, 81-6	1.5	2
27	Hyperacute ischemic stroke without lesions on diffusion-weighted imaging in a patient treated with rtPA thrombolysis. <i>Clinical Case Reports (discontinued)</i> , 2014 , 2, 70-3	0.7	1
26	The iScore predicts clinical response to tissue plasminogen activator in Korean stroke patients. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2014 , 23, 367-73	2.8	7
25	Comparison of fast MRI-based individual thrombolysis therapy for patients with superacute infarction. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2014 , 23, e263-9	2.8	3
24	Myelin basic protein and ischemia modified albumin levels in acute ischemic stroke cases. <i>Pakistan Journal of Medical Sciences</i> , 2015 , 31, 1110-4	2	11
23	Perfusion CT and acute stroke imaging: foundations, applications, and literature review. <i>Journal of Neuroradiology</i> , 2015 , 42, 21-9	3.1	54
22	Serum Aβs predictive for short-term neurological deficits after acute ischemic stroke. <i>Neurotoxicity Research</i> , 2015 , 27, 292-9	4.3	13
21	Dynamic Evolution of Diffusion-Weighted Imaging Lesions in Patients With Minor Ischemic Stroke. <i>Stroke</i> , 2015 , 46, 2318-21	6.7	10
20	Younger Stroke Patients With Large Pretreatment Diffusion-Weighted Imaging Lesions May Benefit From Endovascular Treatment. <i>Stroke</i> , 2015 , 46, 2510-6	6.7	66
19	Serum neuron specific enolase level as a predictor of prognosis in acute ischemic stroke patients after intravenous thrombolysis. <i>Journal of the Neurological Sciences</i> , 2015 , 359, 202-6	3.2	21
18	Infarct volume predicts critical care needs in stroke patients treated with intravenous thrombolysis. <i>Neuroradiology</i> , 2015 , 57, 171-8	3.2	9
17	Sensitivity of diffusion- and perfusion-weighted imaging for diagnosing acute ischemic stroke is 97.5%. <i>Stroke</i> , 2015 , 46, 98-101	6.7	67
16	Prediction of recanalization in acute stroke patients receiving intravenous and endovascular revascularization therapy. <i>International Journal of Stroke</i> , 2015 , 10, 28-36	6.3	14
15	Subacute lesion volume as a potential prognostic biomarker for acute ischemic stroke after intravenous thrombolysis. <i>Journal of the Neurological Sciences</i> , 2016 , 369, 77-81	3.2	2
14	White Matter Hyperintensity-Adjusted Critical Infarct Thresholds to Predict a Favorable 90-Day Outcome. <i>Stroke</i> , 2016 , 47, 2526-33	6.7	20
13	Imaging in acute stroke. <i>Expert Review of Cardiovascular Therapy</i> , 2016 , 14, 963-75	2.5	7

12	Prediction of Early Arterial Recanalization and Tissue Fate in the Selection of Patients With the Greatest Potential to Benefit From Intravenous Tissue-Type Plasminogen Activator. <i>Stroke</i> , 2016 , 47, 397-403	6.7	10
11	Ischemic stroke subtype is associated with outcome in thrombolized patients. <i>Acta Neurologica Scandinavica</i> , 2017 , 135, 176-182	3.8	6
10	Utility of perfusion imaging in acute stroke treatment: a systematic review and meta-analysis. <i>Journal of NeuroInterventional Surgery</i> , 2017 , 9, 1012-1016	7.8	23
9	Effect of glycated hemoglobin index and mean arterial pressure on acute ischemic stroke prognosis after intravenous thrombolysis with recombinant tissue plasminogen activator. <i>Medicine (United States)</i> , 2018 , 97, e13216	1.8	6
8	Relationship between normalized distributional pattern and functional outcome in patients with acute cardiogenic cerebral embolism. <i>PLoS ONE</i> , 2019 , 14, e0210709	3.7	1
7	Susceptibility-weighted imaging in post-treatment evaluation in the early stage in patients with acute ischemic stroke. <i>Journal of International Medical Research</i> , 2019 , 47, 196-205	1.4	6
6	Imaging evaluation of acute ischemic stroke. <i>Journal of International Medical Research</i> , 2020 , 48, 300060518802530	1.8	3
5	Impact of aging and comorbidities on ischemic stroke outcomes in preclinical animal models: A translational perspective. <i>Experimental Neurology</i> , 2021 , 335, 113494	5.7	12
4	Multiclass Support Vector Machine-Based Lesion Mapping Predicts Functional Outcome in Ischemic Stroke Patients. <i>PLoS ONE</i> , 2015 , 10, e0129569	3.7	31
3	Applications of diffusion-weighted imaging in diagnosis, evaluation, and treatment of acute ischemic stroke. <i>Precision and Future Medicine</i> , 2019 , 3, 69-76	1.1	7
2	Portable, low-field magnetic resonance imaging enables highly accessible and dynamic bedside evaluation of ischemic stroke.. <i>Science Advances</i> , 2022 , 8, eabm3952	14.3	2
1	Beyond Diffusion Tensor MRI Methods for Improved Characterization of the Brain after Ischemic Stroke: A Review.. <i>American Journal of Neuroradiology</i> , 2022 ,	4.4	0