

CITATION REPORT

List of articles citing

Trends in treatment strategies and the usage of different disease-modifying anti-rheumatic drugs in early rheumatoid arthritis in Finland. Results from a nationwide register in 2000-2007

DOI: 10.3109/03009742.2010.486768

Scandinavian Journal of Rheumatology, 2011, 40, 16-21.

Source: <https://exaly.com/paper-pdf/52154065/citation-report.pdf>

Version: 2024-04-24

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
20	[Combination therapy using methotrexate with DMARDs or biologics--current status]. <i>Zeitschrift Fur Rheumatologie</i> , 2011 , 70, 114-22	1.9	4
19	Cost-minimisation analysis of subcutaneous methotrexate versus biologic therapy for the treatment of patients with rheumatoid arthritis who have had an insufficient response or intolerance to oral methotrexate. <i>Clinical Rheumatology</i> , 2013 , 32, 1605-12	3.9	17
18	Patterns of disease-modifying antirheumatic drug use in rheumatoid arthritis patients after 2002: a systematic review. <i>Arthritis Care and Research</i> , 2013 , 65, 1927-35	4.7	15
17	A nationwide cross-sectional overview of patients with rheumatoid arthritis followed in outpatient specialty clinics in Finland. <i>Scandinavian Journal of Rheumatology</i> , 2014 , 43, 286-90	1.9	4
16	Antidepressant use among persons with recent-onset rheumatoid arthritis: a nationwide register-based study in Finland. <i>Scandinavian Journal of Rheumatology</i> , 2014 , 43, 364-70	1.9	6
15	[O'Dell reloaded: Do we use TNF inhibitors too early?]. <i>Zeitschrift Fur Rheumatologie</i> , 2014 , 73, 111-2	1.9	0
14	Targeted treatment with a combination of traditional DMARDs produces excellent clinical and radiographic long-term outcomes in early rheumatoid arthritis regardless of initial infliximab. The 5-year follow-up results of a randomised clinical trial, the NEO-RACo trial. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, 1954-61	2.4	31
13	Significantly impaired shoulder function in the first years of rheumatoid arthritis: a controlled study. <i>Arthritis Research and Therapy</i> , 2015 , 17, 261	5.7	9
12	Patients with rheumatoid arthritis have better functional and working ability but poorer general health and higher comorbidity rates today than in the late 1990s. <i>Scandinavian Journal of Rheumatology</i> , 2015 , 44, 173-81	1.9	8
11	Time trends in the incidence, prevalence, and severity of rheumatoid arthritis: A systematic literature review. <i>Joint Bone Spine</i> , 2016 , 83, 625-630	2.9	71
10	Drugs used in incident systemic lupus erythematosus - results from the Finnish nationwide register 2000-2007. <i>Lupus</i> , 2016 , 25, 666-70	2.6	6
9	Incidence, prevalence et sévérité de la polyarthrite rhumatoïde au XXI ^e siècle. <i>Revue Du Rhumatisme Monographies</i> , 2017 , 84, 303-310	0	0
8	Évolution dans le temps de la polyarthrite rhumatoïde : incidence, prévalence, gravité. Revue systématique de la littérature. <i>Revue Du Rhumatisme (Edition Francaise)</i> , 2017 , 84, 9-16	0.1	2
7	Burden of rheumatoid arthritis in the Nordic region, 1990-2015: a comparative analysis using the Global Burden of Disease Study 2015. <i>Scandinavian Journal of Rheumatology</i> , 2018 , 47, 1-101	1.9	11
6	A 4-year non-randomized comparative phase-IV study of early rheumatoid arthritis: integrative anthroposophic medicine for patients with preference against DMARDs versus conventional therapy including DMARDs for patients without preference. <i>Patient Preference and Adherence</i> , 2018 , 12, 275-287	2.4	5
5	High prevalence of widespread pain in women with early rheumatoid arthritis. <i>Scandinavian Journal of Rheumatology</i> , 2018 , 47, 447-454	1.9	12
4	First-year drug therapy of new-onset rheumatoid and undifferentiated arthritis: a nationwide register-based study. <i>BMC Rheumatology</i> , 2020 , 4, 34	2.9	1

- 3 Factors associated with osteoporosis medication use in Japanese patients with rheumatoid arthritis: Results from the Institute of Rheumatology Rheumatoid Arthritis cohort study. *Osteoporosis and Sarcopenia*, **2020**, 6, 82-87 2.3 3
- 2 First-year drug therapy of new-onset rheumatoid and undifferentiated arthritis: A nationwide register-based study..
- 1 First-year drug therapy of new-onset rheumatoid and undifferentiated arthritis: A nationwide register-based study..