

Effects of Drug Burden Index on Cognitive Function in O

Journal of Clinical Psychopharmacology

32, 273-277

DOI: [10.1097/jcp.0b013e3182487825](https://doi.org/10.1097/jcp.0b013e3182487825)

Citation Report

#	ARTICLE	IF	CITATIONS
1	Statin use and clinical outcomes in older men: a prospective population-based study. <i>BMJ Open</i> , 2013, 3, e002333.	0.8	44
2	Drug Burden Index in older adults: theoretical and practical issues. <i>Clinical Interventions in Aging</i> , 2014, 9, 1503.	1.3	113
3	Modifications to the Drug Burden Index Calculation May Limit Interpretation of its Association with Clinical Outcomes in Older Adults. <i>Neurocritical Care</i> , 2014, 21, 365-366.	1.2	0
4	Anticholinergic Drug Burden in Older People's Brain - How well is it Measured?. <i>Basic and Clinical Pharmacology and Toxicology</i> , 2014, 114, 151-159.	1.2	89
5	Exposure to Anticholinergic and Sedative Drugs, Risk of Falls, and Mortality. <i>Journal of Clinical Psychopharmacology</i> , 2014, 34, 565-570.	0.7	58
6	A longitudinal study of knee pain in older men: Concord Health and Ageing in Men Project. <i>Age and Ageing</i> , 2014, 43, 206-212.	0.7	30
7	Sedative load and functional outcomes in community-dwelling older Australian men: the <scp>CHAMP</scp> study. <i>Fundamental and Clinical Pharmacology</i> , 2014, 28, 10-19.	1.0	14
8	Drug Burden Index Score and Anticholinergic Risk Scale as Predictors of Readmission to the Hospital. <i>The Consultant Pharmacist</i> , 2014, 29, 158-168.	0.4	22
9	Associations of drug burden index with falls, general practitioner visits, and mortality in older people. <i>Pharmacoepidemiology and Drug Safety</i> , 2014, 23, 753-758.	0.9	110
10	Effect of medications with anti-cholinergic properties on cognitive function, delirium, physical function and mortality: a systematic review. <i>Age and Ageing</i> , 2014, 43, 604-615.	0.7	269
11	Nonsteroidal anti-inflammatory drugs (NSAIDs) in older people: Prescribing patterns according to pain prevalence and adherence to clinical guidelines. <i>Pain</i> , 2014, 155, 1814-1820.	2.0	20
12	Anticholinergic burden in older women: not seeing the wood for the trees?. <i>Medical Journal of Australia</i> , 2015, 202, 91-94.	0.8	25
13	Ischemic heart disease, prescription of optimal medical therapy and geriatric syndromes in community-dwelling older men: A population-based study. <i>International Journal of Cardiology</i> , 2015, 192, 49-55.	0.8	27
14	Effect of anticholinergic drugs on cognitive impairment in the elderly. <i>Revista De Psiquiatrãa Y Salud Mental (English Edition)</i> , 2015, 8, 35-43.	0.2	8
15	Different methods, different resultsâ€”how do available methods link a patientâ€™s anticholinergic load with adverse outcomes?. <i>European Journal of Clinical Pharmacology</i> , 2015, 71, 1299-1314.	0.8	47
17	Lower Urinary Tract Symptoms and Incident Falls in Community Dwelling Older Men: The Concord Health and Ageing in Men Project. <i>Journal of Urology</i> , 2016, 196, 1694-1699.	0.2	23
18	Anticholinergic medication use and dementia: latest evidence and clinical implications. <i>Therapeutic Advances in Drug Safety</i> , 2016, 7, 217-224.	1.0	53
19	Quantification of anticholinergic and sedative drug load with the Drug Burden Index: a review of outcomes and methodological quality of studies. <i>European Journal of Clinical Pharmacology</i> , 2017, 73, 257-266.	0.8	70

#	ARTICLE	IF	CITATIONS
20	An improved, high-efficiency assay for assessing serum anticholinergic activity using cultured cells stably expressing M1 receptors. <i>Journal of Pharmacological and Toxicological Methods</i> , 2017, 86, 28-33.	0.3	9
21	Drug Burden Index and change in cognition over time in community-dwelling older men: the CHAMP study. <i>Annals of Medicine</i> , 2017, 49, 157-164.	1.5	23
22	Polypharmacy in older adults: Association Rule and Frequent-Set Analysis to evaluate concomitant medication use. <i>Pharmacological Research</i> , 2017, 116, 39-44.	3.1	16
23	Relationship between drug burden and physical and cognitive functions in a sample of nursing home patients with dementia. <i>European Journal of Clinical Pharmacology</i> , 2017, 73, 1633-1642.	0.8	4
24	A multi-center, randomized, controlled trial to assess the efficacy of optimization of drug prescribing in an elderly population, at 18 months of follow-up, in the evolution of functional autonomy: the OPTIM study protocol. <i>BMC Geriatrics</i> , 2017, 17, 195.	1.1	10
25	Drug burden index to define the burden of medicines in older adults with intellectual disabilities: An observational cross-sectional study. <i>British Journal of Clinical Pharmacology</i> , 2018, 84, 553-567.	1.1	45
26	Anticholinergic Drug Burden Tools/Scales and Adverse Outcomes in Different Clinical Settings: A Systematic Review of Reviews. <i>Drugs and Aging</i> , 2018, 35, 523-538.	1.3	92
27	Association of Drug Burden Index with grip strength, timed up and go and Barthel index activities of daily living in older adults with intellectual disabilities: an observational cross-sectional study. <i>BMC Geriatrics</i> , 2019, 19, 173.	1.1	12
28	Analysis of anticholinergic and sedative medicine effects on physical function, cognitive function, appetite and frailty: a cross-sectional study in Australia. <i>BMJ Open</i> , 2019, 9, e029221.	0.8	23
29	Anticholinergic Drugs in Geriatric Psychopharmacology. <i>Frontiers in Neuroscience</i> , 2019, 13, 1309.	1.4	68
30	Using the Drug Burden Index to identify older adults at highest risk for medication-related falls. <i>BMC Geriatrics</i> , 2020, 20, 208.	1.1	9
31	Is Long-Term Benzodiazepine Use a Risk Factor for Cognitive Decline? Results of a Systematic Review. <i>Journal of Addiction</i> , 2020, 2020, 1-10.	0.9	22
32	Impact of <sc>STEADI&ER</sc>: A Community Pharmacy&EBased Fall Prevention Intervention. <i>Journal of the American Geriatrics Society</i> , 2020, 68, 1778-1786.	1.3	13
33	Prevalence and factors associated with frailty in hospitalized older patients. <i>BMC Geriatrics</i> , 2020, 20, 144.	1.1	32
34	Association between anticholinergic burden and anticholinergic adverse outcomes in the elderly: Pharmacological basis of their predictive value for adverse outcomes. <i>Pharmacological Research</i> , 2021, 163, 105306.	3.1	10
35	Size of the associations between anticholinergic burden tool scores and adverse outcomes in older patients. <i>International Journal of Clinical Pharmacy</i> , 2021, 43, 128-136.	1.0	10
36	Exposure to Anticholinergic and Sedative Drugs and Healthcare Costs in Older Patients with Neurocognitive Disorders. <i>Journal of Alzheimer's Disease</i> , 2021, 80, 1515-1524.	1.2	1
37	Anticholinergic burden (prognostic factor) for prediction of dementia or cognitive decline in older adults with no known cognitive syndrome. <i>The Cochrane Library</i> , 2021, 2021, CD013540.	1.5	32

#	ARTICLE	IF	CITATIONS
40	The Impact of Drug Burden Index on Unplanned Hospital Readmission and Length of Hospital Stay. , 0, , .		0
41	Drug Burden Index in Older Adults with Psychiatric Illnesses: A Cross-Sectional Study. Drugs - Real World Outcomes, 0, , .	0.7	0