## CITATION REPORT List of articles citing

Serum uric acid levels and incident chronic kidney disease in patients with type 2 diabetes and preserved kidney function

DOI: 10.2337/dc11-1346 Diabetes Care, 2012, 35, 99-104.

**Source:** https://exaly.com/paper-pdf/53112725/citation-report.pdf

Version: 2024-04-28

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
184	Correlation between plasma renalase level and coronary artery disease. <b>2014</b> , 30, 863-967		7
183	dido dico como factor de riesgo cardiovascular. <b>2012</b> , 29, 36-43		
182	Epidemiology of Chronic Kidney Disease in Diabetes. <b>2012</b> , 14-28		
181	Is hyperuricemia the missing piece in the metabolic syndrome puzzle?. <b>2012</b> , 14, 97-9		
180	Klinisch relevante Faktoren der Progressionshemmung. <b>2012</b> , 7, 40-45		2
179	Uric acid lowering to prevent kidney function loss in diabetes: the preventing early renal function loss (PERL) allopurinol study. <b>2013</b> , 13, 550-9		120
178	Uric acid and chronic kidney disease: which is chasing which?. <b>2013</b> , 28, 2221-8		351
177	Urate reduction and renal preservation: what is the evidence?. <b>2013</b> , 15, 386		5
176	Law, public health, and the diabetes epidemic. <b>2013</b> , 45, 486-93		5
175	Uric acid as a target of therapy in CKD. American Journal of Kidney Diseases, 2013, 61, 134-46	7.4	165
174	The effects of mammalian target of rapamycin inhibitors on serum uric acid levels in renal transplant patients. <b>2013</b> , 45, 547-52		2
173	Serum uric acid and cardio-renal diseases. <b>2013</b> , 29 Suppl 3, 25-31		6
172	Association of uric acid and left ventricular mass index with renal outcomes in chronic kidney disease. <b>2013</b> , 26, 243-9		11
171	Current world literature. 2013, 22, 238-50		
170	Hyperuricaemia: more than just a cause of gout?. <b>2013</b> , 14, 397-402		58
169	Uric acid and chronic kidney disease: A time to act?. <b>2013</b> , 2, 17-25		28
168	Uric acid promotes apoptosis in human proximal tubule cells by oxidative stress and the activation of NADPH oxidase NOX 4. <i>PLoS ONE</i> , <b>2014</b> , 9, e115210	3.7	78

167	The association of albuminuria with tubular reabsorption of uric acid: results from a general population cohort. <b>2014</b> , 3, e000613		15
166	Prevention of diabetic nephropathy by compound 21, selective agonist of angiotensin type 2 receptors, in Zucker diabetic fatty rats. <i>American Journal of Physiology - Renal Physiology</i> , <b>2014</b> , 307, F112	<del>2</del> 3 <sup>2</sup> 31	39
165	Hyperuricemia and chronic kidney disease: an enigma yet to be solved. <b>2014</b> , 36, 1351-9		27
164	Serum uric acid concentration is associated with worsening in severity of diabetic retinopathy among type 2 diabetic patients in Taiwana 3-year prospective study. <b>2014</b> , 106, 366-72		29
163	Salusin-alpha and -beta expression in heart and aorta with and without metabolic syndrome. <b>2014</b> , 89, 98-103		8
162	Serum uric acid, the metabolic syndrome, and the risk of chronic kidney disease in patients with type 2 diabetes. <b>2014</b> , 12, 102-9		20
161	Uric acid: association with rate of renal function decline and time until start of dialysis in incident pre-dialysis patients. <b>2014</b> , 15, 91		19
160	Is hyperuricemia an independent risk factor for new-onset chronic kidney disease?: A systematic review and meta-analysis based on observational cohort studies. <b>2014</b> , 15, 122		195
159	Markers of and risk factors for the development and progression of diabetic kidney disease.  American Journal of Kidney Diseases, <b>2014</b> , 63, S39-62	7.4	185
158	High-normal serum uric acid predicts the development of chronic kidney disease in patients with type 2 diabetes mellitus and preserved kidney function. <i>Journal of Diabetes and Its Complications</i> , <b>2014</b> , 28, 130-4	3.2	37
157	Relation of uric acid and contrast-induced nephropathy in patients undergoing primary percutaneous coronary intervention in the ED. <b>2014</b> , 32, 119-23		15
156	Uric acid, chronic kidney disease and type 2 diabetes: a cluster of vascular risk factors. <i>Journal of Diabetes and Its Complications</i> , <b>2014</b> , 28, 122-3	3.2	12
155	Comorbidities in patients with crystal diseases and hyperuricemia. <b>2014</b> , 40, 251-78		32
154	Serum uric acid and the risk of cardiovascular and renal disease. <b>2015</b> , 33, 1729-41; discussion 1741		267
153	Estimated glomerular filtration rate progression in UK primary care patients with type 2 diabetes and diabetic kidney disease: a retrospective cohort study. <b>2015</b> , 69, 871-82		10
152	Five-year incidence of chronic kidney disease (stage 3-5) and associated risk factors in a Spanish cohort: the MADIABETES Study. <i>PLoS ONE</i> , <b>2015</b> , 10, e0122030	3.7	20
151	Use of Contrast-Enhanced Ultrasound to Study Relationship between Serum Uric Acid and Renal Microvascular Perfusion in Diabetic Kidney Disease. <i>BioMed Research International</i> , <b>2015</b> , 2015, 732317	3	6
150	Urate lowering therapy to improve renal outcomes in patients with chronic kidney disease: systematic review and meta-analysis. <b>2015</b> , 16, 58		107

149	Uric Acid is independently associated with diabetic kidney disease: a cross-sectional study in a Chinese population. <i>PLoS ONE</i> , <b>2015</b> , 10, e0129797	3.7	35
148	Management of Chronic Kidney Disease: The Relationship Between Serum Uric Acid and Development of Nephropathy. <b>2015</b> , 32, 1177-91		36
147	Hyperuricemia-induced NLRP3 activation of macrophages contributes to the progression of diabetic nephropathy. <i>American Journal of Physiology - Renal Physiology</i> , <b>2015</b> , 308, F993-F1003	4.3	103
146	Metabonomics revealed xanthine oxidase-induced oxidative stress and inflammation in the pathogenesis of diabetic nephropathy. <i>Analytical and Bioanalytical Chemistry</i> , <b>2015</b> , 407, 2569-79	4.4	55
145	Uric acid as a biomarker and a therapeutic target in diabetes. <i>Canadian Journal of Diabetes</i> , <b>2015</b> , 39, 239-46	2.1	77
144	The molecular insight into the antihyperuricemic and renoprotective effect of Shuang Qi gout capsule in mice. <b>2015</b> , 163, 278-89		9
143	Uric acid is not associated with decline in renal function or time to renal replacement therapy initiation in a referred cohort of patients with Stage III, IV and V chronic kidney disease. <b>2015</b> , 30, 2039	-45	24
142	Efficacy of Febuxostat for Slowing the GFR Decline in Patients With CKD and Asymptomatic Hyperuricemia: A 6-Month, Double-Blind, Randomized, Placebo-Controlled Trial. <i>American Journal of Kidney Diseases</i> , <b>2015</b> , 66, 945-50	7.4	169
141	Prognostic clinical and molecular biomarkers of renal disease in type 2 diabetes. <b>2015</b> , 30 Suppl 4, iv86	-95	29
140	Fructose and uric acid in diabetic nephropathy. <b>2015</b> , 58, 1993-2002		73
140	Fructose and uric acid in diabetic nephropathy. <b>2015</b> , 58, 1993-2002  Decreased urine uric acid excretion is an independent risk factor for chronic kidney disease but not for carotid atherosclerosis in hospital-based patients with type 2 diabetes: a cross-sectional study. <b>2015</b> , 14, 36		73
	Decreased urine uric acid excretion is an independent risk factor for chronic kidney disease but not for carotid atherosclerosis in hospital-based patients with type 2 diabetes: a cross-sectional study.		
139	Decreased urine uric acid excretion is an independent risk factor for chronic kidney disease but not for carotid atherosclerosis in hospital-based patients with type 2 diabetes: a cross-sectional study. <b>2015</b> , 14, 36  Pterostilbene and allopurinol reduce fructose-induced podocyte oxidative stress and inflammation	3.2	13
139	Decreased urine uric acid excretion is an independent risk factor for chronic kidney disease but not for carotid atherosclerosis in hospital-based patients with type 2 diabetes: a cross-sectional study. 2015, 14, 36  Pterostilbene and allopurinol reduce fructose-induced podocyte oxidative stress and inflammation via microRNA-377. 2015, 83, 214-26  Gender differences in the association between hyperuricemia and diabetic kidney disease in	3.2	13
139 138 137	Decreased urine uric acid excretion is an independent risk factor for chronic kidney disease but not for carotid atherosclerosis in hospital-based patients with type 2 diabetes: a cross-sectional study. 2015, 14, 36  Pterostilbene and allopurinol reduce fructose-induced podocyte oxidative stress and inflammation via microRNA-377. 2015, 83, 214-26  Gender differences in the association between hyperuricemia and diabetic kidney disease in community elderly patients. <i>Journal of Diabetes and Its Complications</i> , 2015, 29, 1042-9	3.2	13 113 5
139 138 137	Decreased urine uric acid excretion is an independent risk factor for chronic kidney disease but not for carotid atherosclerosis in hospital-based patients with type 2 diabetes: a cross-sectional study. 2015, 14, 36  Pterostilbene and allopurinol reduce fructose-induced podocyte oxidative stress and inflammation via microRNA-377. 2015, 83, 214-26  Gender differences in the association between hyperuricemia and diabetic kidney disease in community elderly patients. <i>Journal of Diabetes and Its Complications</i> , 2015, 29, 1042-9  Detrimental role of hyperuricemia on the cardio-reno-vascular system. 2015, 31 Suppl 2, 21-6	3.2	13 113 5
139 138 137 136	Decreased urine uric acid excretion is an independent risk factor for chronic kidney disease but not for carotid atherosclerosis in hospital-based patients with type 2 diabetes: a cross-sectional study. 2015, 14, 36  Pterostilbene and allopurinol reduce fructose-induced podocyte oxidative stress and inflammation via microRNA-377. 2015, 83, 214-26  Gender differences in the association between hyperuricemia and diabetic kidney disease in community elderly patients. <i>Journal of Diabetes and Its Complications</i> , 2015, 29, 1042-9  Detrimental role of hyperuricemia on the cardio-reno-vascular system. 2015, 31 Suppl 2, 21-6  Therapies on the Horizon for Diabetic Kidney Disease. 2015, 15, 111	3.2	13 113 5 11 20

131	Uric Acid as a Marker of Mortality and Morbidity in Fabry Disease. <i>PLoS ONE</i> , <b>2016</b> , 11, e0166290	3.7	2
130	Effects of xanthine oxidase inhibition with febuxostat on the development of nephropathy in experimental type 2 diabetes. <b>2016</b> , 173, 2573-88		24
129	Thyroid hormones associate with risk of incident chronic kidney disease and rapid decline in renal function: a prospective investigation. <b>2016</b> , 14, 336		13
128	Association between serum uric acid related genetic loci and diabetic kidney disease in the Chinese type 2 diabetes patients. <i>Journal of Diabetes and Its Complications</i> , <b>2016</b> , 30, 798-802	3.2	5
127	Gout, hyperuricemia and chronic kidney disease: New treatment possibilities. <b>2016</b> , 23, 195-201		3
126	Inflammasomes in the urinary tract: a disease-based review. <i>American Journal of Physiology - Renal Physiology</i> , <b>2016</b> , 311, F653-F662	4.3	15
125	Hyperuricemia contributes to the faster progression of diabetic kidney disease in type 2 diabetes mellitus. <i>Journal of Diabetes and Its Complications</i> , <b>2016</b> , 30, 1300-7	3.2	36
124	Diabetic Kidney Disease in Adolescents With Type 2 Diabetes: New Insights and Potential Therapies. <b>2016</b> , 16, 11		22
123	Serum uric acid level as an indicator for CKD regression and progression in patients with type 2 diabetes mellitus-a 4.6-year cohort study. <b>2016</b> , 32, 557-64		38
122	Serum uric acid levels are associated with a high risk of rapid chronic kidney disease progression among patients with type 2 diabetes: a prospective cohort study [Diabetes Distress and Care Registry at Tenri (DDCRT 12)]. <b>2016</b> , 7, 352-360		6
122	among patients with type 2 diabetes: a prospective cohort study [Diabetes Distress and Care		6
	among patients with type 2 diabetes: a prospective cohort study [Diabetes Distress and Care Registry at Tenri (DDCRT 12)]. <b>2016</b> , 7, 352-360  Sociodemographic, behavioral and genetic determinants of allostatic load in a Swiss	193	
	among patients with type 2 diabetes: a prospective cohort study [Diabetes Distress and Care Registry at Tenri (DDCRT 12)]. <b>2016</b> , 7, 352-360  Sociodemographic, behavioral and genetic determinants of allostatic load in a Swiss population-based study. <b>2016</b> , 67, 76-85	<b>193</b>	34
121	among patients with type 2 diabetes: a prospective cohort study [Diabetes Distress and Care Registry at Tenri (DDCRT 12)]. <b>2016</b> , 7, 352-360  Sociodemographic, behavioral and genetic determinants of allostatic load in a Swiss population-based study. <b>2016</b> , 67, 76-85  Time to target uric acid to retard CKD progression. <i>Clinical and Experimental Nephrology</i> , <b>2017</b> , 21, 182-Sustained kidney biochemical derangement in treated experimental diabetes: a clue to metabolic		34 56
121 120 119	among patients with type 2 diabetes: a prospective cohort study [Diabetes Distress and Care Registry at Tenri (DDCRT 12)]. 2016, 7, 352-360  Sociodemographic, behavioral and genetic determinants of allostatic load in a Swiss population-based study. 2016, 67, 76-85  Time to target uric acid to retard CKD progression. Clinical and Experimental Nephrology, 2017, 21, 182-Sustained kidney biochemical derangement in treated experimental diabetes: a clue to metabolic memory. Scientific Reports, 2017, 7, 40544  Serum uric acid to creatinine ratio: A predictor of incident chronic kidney disease in type 2 diabetes		<ul><li>34</li><li>56</li><li>9</li></ul>
121 120 119 118	among patients with type 2 diabetes: a prospective cohort study [Diabetes Distress and Care Registry at Tenri (DDCRT 12)]. 2016, 7, 352-360  Sociodemographic, behavioral and genetic determinants of allostatic load in a Swiss population-based study. 2016, 67, 76-85  Time to target uric acid to retard CKD progression. Clinical and Experimental Nephrology, 2017, 21, 182-Sustained kidney biochemical derangement in treated experimental diabetes: a clue to metabolic memory. Scientific Reports, 2017, 7, 40544  Serum uric acid to creatinine ratio: A predictor of incident chronic kidney disease in type 2 diabetes mellitus patients with preserved kidney function. 2017, 14, 221-225  Incidence of chronic kidney disease among people with diabetes: a systematic review of		<ul><li>34</li><li>56</li><li>9</li><li>30</li></ul>
121 120 119 118	among patients with type 2 diabetes: a prospective cohort study [Diabetes Distress and Care Registry at Tenri (DDCRT 12)]. 2016, 7, 352-360  Sociodemographic, behavioral and genetic determinants of allostatic load in a Swiss population-based study. 2016, 67, 76-85  Time to target uric acid to retard CKD progression. Clinical and Experimental Nephrology, 2017, 21, 182-Sustained kidney biochemical derangement in treated experimental diabetes: a clue to metabolic memory. Scientific Reports, 2017, 7, 40544  Serum uric acid to creatinine ratio: A predictor of incident chronic kidney disease in type 2 diabetes mellitus patients with preserved kidney function. 2017, 14, 221-225  Incidence of chronic kidney disease among people with diabetes: a systematic review of observational studies. 2017, 34, 887-901  Uric Acid Is Associated With Inflammatory Biomarkers and Induces Inflammation Via Activating the		<ul><li>34</li><li>56</li><li>9</li><li>30</li><li>59</li></ul>

113	The problem of cardio-renal diseases in patients with gout. <b>2017</b> , 33, 9-13		4
112	Impact of Uric Acid Levels on Kidney Disease Progression. <b>2017</b> , 46, 315-322		15
111	Clinical predictive factors in diabetic kidney disease progression. <b>2017</b> , 8, 6-18		79
110	Hyperuricemia and Urologic Disease. <b>2017</b> , 12, 103		
109	Increased Serum Uric Acid Level Is a Risk Factor for Left Ventricular Hypertrophy but Not Independent of eGFR in Patients with Type 2 Diabetic Kidney Disease. <b>2017</b> , 2017, 5016093		2
108	Correlation between Serum Uric Acid Level and Microalbuminuria in Type-2 Diabetic Nephropathy. <b>2017</b> , 33, 1371-1375		8
107	Hyperuricemia as a Predictive Marker for Progression of Nephrosclerosis: Clinical Assessment of Prognostic Factors in Biopsy-Proven Arterial/Arteriolar Nephrosclerosis. <b>2017</b> , 24, 630-642		17
106	Does Altered Uric Acid Metabolism Contribute to Diabetic Kidney Disease Pathophysiology?. <b>2018</b> , 18, 18		9
105	Drug repurposing in kidney disease. <i>Kidney International</i> , <b>2018</b> , 94, 40-48	9.9	28
104	Association of serum uric acid with nephropathy in Iranian type 2 diabetic patients. <i>Journal of Diabetes and Metabolic Disorders</i> , <b>2018</b> , 17, 71-75	2.5	3
104		2.5	9
<u> </u>	Diabetes and Metabolic Disorders, <b>2018</b> , 17, 71-75	2.5	
103	Diabetes and Metabolic Disorders, 2018, 17, 71-75  Uric Acid: The Unknown Uremic Toxin. 2018, 192, 25-33	2.5	9
103	Diabetes and Metabolic Disorders, 2018, 17, 71-75  Uric Acid: The Unknown Uremic Toxin. 2018, 192, 25-33  Uric Acid and Diabetic Nephropathy Risk. 2018, 192, 103-109  The relationships among hyperuricemia, body mass index and impaired renal function in type 2	2.5	9
103	Diabetes and Metabolic Disorders, 2018, 17, 71-75  Uric Acid: The Unknown Uremic Toxin. 2018, 192, 25-33  Uric Acid and Diabetic Nephropathy Risk. 2018, 192, 103-109  The relationships among hyperuricemia, body mass index and impaired renal function in type 2 diabetic patients. 2018, 65, 281-290	2.5	9 19 3
103 102 101	Uric Acid: The Unknown Uremic Toxin. 2018, 192, 25-33  Uric Acid and Diabetic Nephropathy Risk. 2018, 192, 103-109  The relationships among hyperuricemia, body mass index and impaired renal function in type 2 diabetic patients. 2018, 65, 281-290  An association of smoking with serum urate and gout: A health paradox. 2018, 47, 825-842	2.5	9 19 3 16
103 102 101 100	Uric Acid: The Unknown Uremic Toxin. 2018, 17, 71-75  Uric Acid and Diabetic Nephropathy Risk. 2018, 192, 103-109  The relationships among hyperuricemia, body mass index and impaired renal function in type 2 diabetic patients. 2018, 65, 281-290  An association of smoking with serum urate and gout: A health paradox. 2018, 47, 825-842  Unterschiedliche Hyperurikihie-Philotypen rufen nach Differenzialtherapie. 2018, 12, 22-23  Scutellariae Radix and Coptidis Rhizoma Improve Glucose and Lipid Metabolism in T2DM Rats via	2.5	9 19 3 16

## (2020-2018)

95	Serum uric acid levels and decreased estimated glomerular filtration rate in patients with type 2 diabetes: A cohort study and meta-analysis. <b>2018</b> , 34, e3046		16
94	Incidence and predictors of chronic kidney diseases among type 2 diabetes mellitus patients at St. Paul's Hospital, Addis Ababa, Ethiopia. <b>2018</b> , 11, 532		12
93	Effects of uric acid on kidney function decline differ depending on baseline kidney function in type 2 diabetic patients. <b>2019</b> , 34, 1328-1335		8
92	Temporal trends in hyperuricaemia in the Irish health system from 2006-2014: A cohort study. <i>PLoS ONE</i> , <b>2018</b> , 13, e0198197	3.7	30
91	Dietary Antioxidant Supplements and Uric Acid in Chronic Kidney Disease: A Review. <b>2019</b> , 11,		33
90	Sodium-Glucose Cotransporter-2 (SGLT2) Inhibitors: A Clinician's Guide. <b>2019</b> , 12, 2125-2136		26
89	Diabetic kidney disease: New clinical and therapeutic issues. Joint position statement of the Italian Diabetes Society and the Italian Society of Nephrology on "The natural history of diabetic kidney disease and treatment of hyperglycemia in patients with type 2 diabetes and impaired renal		38
88	function". <b>2019</b> , 29, 1127-1150 High Serum Uric Acid Is Associated with Tubular Damage and Kidney Inflammation in Patients with Type 2 Diabetes. <b>2019</b> , 2019, 6025804		4
87	Association between serum 25-hydroxyvitamin D and diabetic kidney disease in Chinese patients with type 2 diabetes. <i>PLoS ONE</i> , <b>2019</b> , 14, e0214728	3.7	11
86	Triglyceride-glucose index is a predictor of incident chronic kidney disease: a population-based longitudinal study. <i>Clinical and Experimental Nephrology</i> , <b>2019</b> , 23, 948-955	2.5	12
85	Coexistence of hyper-uricaemia and low urinary uric acid excretion further increases risk of chronic kidney disease in type 2 diabetes. <b>2019</b> , 45, 557-563		2
84	Association of serum uric acid levels with the incident of kidney disease and rapid eGFR decline in Chinese individuals with eGFR > 60′mL/min/1.73′m and negative proteinuria. <i>Clinical and Experimental Nephrology</i> , <b>2019</b> , 23, 871-879	2.5	11
83	Uric acid and the cardio-renal effects of SGLT2 inhibitors. <i>Diabetes, Obesity and Metabolism</i> , <b>2019</b> , 21, 1291-1298	6.7	68
82	Effect of Stem Extract on Uric Acid Excretion in Hyperuricemia Mice. <i>Frontiers in Pharmacology</i> , <b>2019</b> , 10, 1464	5.6	15
81	The association between serum uric acid to creatinine ratio and renal disease progression in type 2 diabetic patients in Chinese communities. <i>Journal of Diabetes and Its Complications</i> , <b>2019</b> , 33, 473-476	3.2	2
80	Diabetic kidney disease: new clinical and therapeutic issues. Joint position statement of the Italian Diabetes Society and the Italian Society of Nephrology on "The natural history of diabetic kidney disease and treatment of hyperglycemia in patients with type 2 diabetes and impaired renal	4.8	38
79	Plasma uric acid and renal haemodynamics in type 2 diabetes patients. <i>Nephrology</i> , <b>2020</b> , 25, 290-297	2.2	
78	Association between serum uric acid and proteinuria in patients with type 2 diabetes and stages 1 and 2 chronic kidney disease. <i>Clinical Epidemiology and Global Health</i> , <b>2020</b> , 8, 166-169	1.8	O

77	Time-averaged serum uric acid and 10-year incident diabetic kidney disease: A prospective study from China. <i>Journal of Diabetes</i> , <b>2020</b> , 12, 169-178	3.8	1
76	Fructose tolerance test in obese people with and without type 2 diabetes. <i>Journal of Diabetes</i> , <b>2020</b> , 12, 197-204	3.8	2
75	Effects of dams and their environmental impacts on the genetic diversity and connectivity of freshwater mussel populations in Poyang Lake Basin, China. <i>Freshwater Biology</i> , <b>2020</b> , 65, 264-277	3.1	6
74	Lycium barbarum L. leaves ameliorate type 2 diabetes in rats by modulating metabolic profiles and gut microbiota composition. <i>Biomedicine and Pharmacotherapy</i> , <b>2020</b> , 121, 109559	7.5	32
73	Study on the Risk Factors for Hyperuricaemia and Related Vascular Complications in Patients with Type 2 Diabetes Mellitus. <i>Risk Management and Healthcare Policy</i> , <b>2020</b> , 13, 1661-1675	2.8	3
7 <del>2</del>	Uric acid and risk of diabetic kidney disease. <i>Journal of Nephrology</i> , <b>2020</b> , 33, 995-999	4.8	5
71	Hyperuricaemia and gout in cardiovascular, metabolic and kidney disease. <i>European Journal of Internal Medicine</i> , <b>2020</b> , 80, 1-11	3.9	46
70	An electrochemical biosensor based on multi-wall carbon nanotube-modified screen-printed electrode immobilized by uricase for the detection of salivary uric acid. <i>Analytical and Bioanalytical Chemistry</i> , <b>2020</b> , 412, 7275-7283	4.4	22
69	Association between Neck Circumference and the Risk of Decreased Estimated Glomerular Filtration Rate in the General Population of China: A Cross-Sectional Study. <i>BioMed Research International</i> , <b>2020</b> , 2020, 3496328	3	8
68	Uric acid and inflammation in kidney disease. <i>American Journal of Physiology - Renal Physiology</i> , <b>2020</b> , 318, F1327-F1340	4.3	22
67	Hyperuricemia as a trigger of immune response in hypertension and chronic kidney disease. <i>Kidney International</i> , <b>2020</b> , 98, 1149-1159	9.9	18
66	Empagliflozin Attenuates Hyperuricemia by Upregulation of ABCG2 via AMPK/AKT/CREB Signaling Pathway in Type 2 Diabetic Mice. <i>International Journal of Biological Sciences</i> , <b>2020</b> , 16, 529-542	11.2	20
65	Serum uric acid is independently associated with diabetic nephropathy but not diabetic retinopathy in patients with type 2 diabetes mellitus. <i>Journal of the Chinese Medical Association</i> , <b>2020</b> , 83, 350-356	2.8	7
64	Prevalence and Risk Factors of Chronic Kidney Disease among Type 2 Diabetes Patients: A Cross-Sectional Study in Primary Care Practice. <i>Scientific Reports</i> , <b>2020</b> , 10, 6205	4.9	26
63	Cardiovascular risk assessment after one-year acute ischemic stroke based on uric acid levels and renal dysfunction. A clinical study. <i>International Journal of Neuroscience</i> , <b>2021</b> , 131, 609-614	2	2
62	Association Between Serum Albumin Level and All-Cause Mortality in Patients With Chronic Kidney Disease: A Retrospective Cohort Study. <i>American Journal of the Medical Sciences</i> , <b>2021</b> , 361, 451-460	2.2	3
61	The Association between Urinary Glucose and Renal Uric Acid Excretion in Non-diabetic Patients with Stage 1-2 Chronic Kidney Disease. <i>Endocrine Research</i> , <b>2021</b> , 46, 28-36	1.9	1
60	SGLT2 Inhibition for CKD and Cardiovascular Disease in Type 2 Diabetes: Report of a Scientific Workshop Sponsored by the National Kidney Foundation. <i>American Journal of Kidney Diseases</i> , <b>2021</b> , 77, 94-109	7.4	27

59	SGLT2 Inhibition for CKD and Cardiovascular Disease in Type 2 Diabetes: Report of a Scientific Workshop Sponsored by the National Kidney Foundation. <i>Diabetes</i> , <b>2021</b> , 70, 1-16	0.9	18
58	The association between serum uric acid and diabetic complications in patients with type 2 diabetes mellitus by gender: a cross-sectional study. <i>Peer J.</i> <b>2021</b> , 9, e10691	3.1	3
57	Serum Uric acid is a better indicator of kidney impairment than serum uric acid to creatine ratio; a cross sectional study of type 2 diabetes mellitus patients. <i>Journal of Diabetes and Metabolic Disorders</i> , <b>2021</b> , 20, 313-320	2.5	1
56	Drug Repurposing: An Emerging Tool for Drug Reuse, Recycling and Discovery. <i>Current Drug Research Reviews</i> , <b>2021</b> , 13, 101-119	2	4
55	Novel biomarkers of diabetic kidney disease: current status and potential clinical application. <i>Acta Diabetologica</i> , <b>2021</b> , 58, 819-830	3.9	8
54	Association between uric acid lowering and renal function progression: a longitudinal study. <i>PeerJ</i> , <b>2021</b> , 9, e11073	3.1	
53	A kidney perspective on the mechanism of action of sodium glucose co-transporter 2 inhibitors. <i>Cell Metabolism</i> , <b>2021</b> , 33, 732-739	24.6	25
52	Designing a Strategy Trial for the Management of Gout: The Use of a Modified Delphi Panel. <i>ACR Open Rheumatology</i> , <b>2021</b> , 3, 341-348	3.5	1
51	Association Between Classical and Emerging Risk Factors for Diabetic Kidney Disease and Albuminuria in a Cohort of Type 2 Diabetes Mellitus Patients. <i>Medicina Interna (Bucharest, Romania: 1991)</i> , <b>2021</b> , 18, 17-25	0.1	
50	The Impact of Uric Acid and Hyperuricemia on Cardiovascular and Renal Systems. <i>Cardiology Clinics</i> , <b>2021</b> , 39, 365-376	2.5	6
49	Empagliflozin and uric acid metabolism in diabetes: A post hoc analysis of the EMPA-REG OUTCOME trial. <i>Diabetes, Obesity and Metabolism</i> , <b>2022</b> , 24, 135-141	6.7	6
48	Association between serum uric acid and renal outcome in patients with biopsy-confirmed diabetic nephropathy. <i>Endocrine Connections</i> , <b>2021</b> , 10, 1299-1306	3.5	2
47	New progress in drugs treatment of diabetic kidney disease. <i>Biomedicine and Pharmacotherapy</i> , <b>2021</b> , 141, 111918	7.5	3
46	Allopurinol and Renal Outcomes in Adults With and Without Type 2 Diabetes: A Retrospective, Population-Based Cohort Study and Propensity Score Analysis. <i>Canadian Journal of Diabetes</i> , <b>2021</b> , 45, 641-649.e4	2.1	1
45	Asymptomatic hyperuricemia and chronic kidney disease: Narrative review of a treatment controversial. <i>Journal of Advanced Research</i> , <b>2017</b> , 8, 555-560	13	38
44	Hyperuricemia as an independent predictor of vascular complications and mortality in type 2 diabetes patients: a meta-analysis. <i>PLoS ONE</i> , <b>2013</b> , 8, e78206	3.7	56
43	Uric Acid Control in Advanced Chronic Kidney Disease in a Southeastern US Urban Cohort. <i>Southern Medical Journal</i> , <b>2018</b> , 111, 549-555	0.6	2
42	Urate-Lowering Therapy Ameliorates Kidney Function in Type 2 Diabetes Patients With Hyperuricemia. <i>Journal of Clinical Medicine Research</i> , <b>2017</b> , 9, 1007-1012	2.9	1

41	Is Uric Acid elevation a random finding or a causative agent of diabetic nephropathy?. <i>Revista Da Associa</i> Mdica Brasileira, <b>2019</b> , 65, 1155-1160	1.4	15
40	Renal effects of uric acid: hyperuricemia and hypouricemia. <i>Korean Journal of Internal Medicine</i> , <b>2020</b> , 35, 1291-1304	2.5	8
39	Hyperuricemia as a Predictor of Progression of Chronic Kidney Disease: A Matched Cohort Analysis. <i>International Journal of Clinical Medicine</i> , <b>2017</b> , 08, 178-197	0.3	1
38	The association between elevated serum uric acid levels and islet Etell function indexes in newly diagnosed type 2 diabetes mellitus: a cross-sectional study. <i>PeerJ</i> , <b>2018</b> , 6, e4515	3.1	9
37	Estimation of the prevalence of type 2 diabetes in combination with diabetic kidney disease and identification of the associated factors in patients attending primary hospitals in Anhui Province, China. <i>Journal of International Medical Research</i> , <b>2021</b> , 49, 3000605211051225	1.4	О
36	Sex Difference in the Association Between Eating Away From Home and the Risk of High Serum Uric Acid in South China. <i>Frontiers in Nutrition</i> , <b>2021</b> , 8, 647287	6.2	O
35	Serum Uric Acid Levels and Risk of Rapid Decline of Estimated Glomerular Filtration Rate in Patients with Type 2 Diabetes: Findings from a 5-Year Prospective Cohort Study. <i>Healthcare</i> (Switzerland), 2021, 9,	3.4	2
34	Correlation between serum uric Acid and renal function in patients with stable coronary artery disease and type 2 diabetes. <i>Journal of Clinical Medicine Research</i> , <b>2014</b> , 6, 443-50	2.9	8
33	SERUM URIC ACID AND TYPE 2 DIABETES MELLITUS COMPLICATIONS. <i>Journal of Evidence Based Medicine and Healthcare</i> , <b>2018</b> , 5, 870-873	O	
32	Serum Uric Acid to Creatinine Ratio as a Predictor of Metabolic Syndrome in Healthy Adults Men. <i>Korean Journal of Clinical Laboratory Science</i> , <b>2019</b> , 51, 42-49	0.4	
31	POTENTIAL DUAL MECHANISM OF HYPOURICEMICACTIVITY OF DPP-4 INHIBITORS WITH PURINE-BASED SCAFFOLD. <i>Acta Medica Medianae</i> , <b>2019</b> , 58, 50-63	O	
30	STORIA NATURALE DELLA MALATTIA RENALE NEL DIABETE E TRATTAMENTO DELL <b>I</b> PERGLICEMIA NEI PAZIENTI CON DIABETE DI TIPO 2 E RIDOTTA FUNZIONE RENALE. <i>Il Diabete</i> , <b>2019</b> , 3,	Ο	
29	Cardiovascular disease-related mortality risk in end stage renal disease and type 2 diabetes: A systematic review. <i>Journal of Family Medicine and Primary Care</i> , <b>2020</b> , 9, 3195-3199	1.5	1
28	[Asymptomatic Hyperuricemia and Risk Of Cardiovascular and Renal Diseases]. <i>Kardiologiya</i> , <b>2020</b> , 60, 113-121	1.5	4
27	The effect of allopurinol on lowering blood pressure in hemodialysis patients with hyperuricemia. <i>Journal of Research in Medical Sciences</i> , <b>2012</b> , 17, 1039-46	1.6	10
26	Association between uric acid level and incidence of albuminuria in patients with type 2 diabetes mellitus: A 4.5-year cohort study. <i>Medicine (United States)</i> , <b>2021</b> , 100, e27496	1.8	
25	Re-thinking diabetic nephropathy: Microalbuminuria is just a piece of the diagnostic puzzle. <i>Clinica Chimica Acta</i> , <b>2021</b> , 524, 146-146	6.2	О
24	Association between uric acid level and incidence of albuminuria in patients with type 2 diabetes mellitus. <i>Medicine (United States)</i> , <b>2021</b> , 100, e27496	1.8	О

23	Uric Acid and Impairment of Renal Function in Non-diabetic Hypertensive Patients <i>Frontiers in Medicine</i> , <b>2021</b> , 8, 746886	4.9	O
22	Incidence and predictors of chronic kidney disease in type-II diabetes mellitus patients attending at the Amhara region referral hospitals, Ethiopia: A follow-up study <i>PLoS ONE</i> , <b>2022</b> , 17, e0263138	3.7	O
21	Phytomedicine as a source of SGLT2 inhibitors, GLP-1 secretagogues and DPP-IV inhibitors for mitigation of Diabetic Nephropathy. <i>Phytomedicine Plus</i> , <b>2022</b> , 2, 100225		
20	Diabetic Kidney Disease: From Pathogenesis to Novel Treatment Possibilities <i>Handbook of Experimental Pharmacology</i> , <b>2022</b> , 1	3.2	O
19	Effectiveness of Early Advanced Glycation End Product Accumulation Testing in the Diagnosis of Diabetes: A Health Risk Factor Analysis Using the Body Mass Index as a Moderator <i>Frontiers in Endocrinology</i> , <b>2021</b> , 12, 766778	5.7	О
18	Serum uric acid levels and diabetic kidney disease in patients with type 2 diabetes mellitus: A dose-response meta-analysis <i>Primary Care Diabetes</i> , <b>2022</b> ,	2.4	Ο
17	Genetic risk score for plasma uric acid levels is associated with early rapid kidney function decline in type 2 diabetes <i>Journal of Clinical Endocrinology and Metabolism</i> , <b>2022</b> ,	5.6	0
16	DataSheet_1.docx. <b>2019</b> ,		
15	Carvacrol Alleviates Hyperuricemia-Induced Oxidative Stress and Inflammation by Modulating the NLRP3/NF- <b>B</b> Pathwayt <i>Drug Design, Development and Therapy</i> , <b>2022</b> , 16, 1159-1170	4.4	2
14	Serum Uric Acid and Metabolic Markers in Diabetes. <i>Biomarkers in Disease</i> , <b>2022</b> , 1-21		
13	Mechanism of glycometabolism regulation by bioactive compounds from the fruits of Lycium barbarum: A review. <i>Food Research International</i> , <b>2022</b> , 111408	7	О
12	Uric acid and cardiometabolic risk by gender in youth with type 1 diabetes. <i>Scientific Reports</i> , <b>2022</b> , 12,	4.9	
11	Advanced glycation end products and diabetes and other metabolic indicators. 2022, 14,		1
10	Effects of allopurinol on renal function in patients with diabetes: a systematic review and meta-analysis. <b>2022</b> , 44, 806-814		O
9	Non-Immunologic Causes of Late Death-Censored Kidney Graft Failure: A Personalized Approach. <b>2022</b> , 12, 1271		
8	SGLT2 Inhibitors in Patients with Chronic Kidney Disease and Heart Disease: A Literature Review. <b>2022</b> , 18, 62-72		O
7	Sodium Clucose Cotransporter-2 Inhibitors Could Help Delay Renal Impairment in Patients with Type 2 Diabetes: A Real-World Clinical Setting. <b>2022</b> , 11, 5259		1
6	Serum Uric Acid and Metabolic Markers in Diabetes. <b>2023</b> , 239-259		О

5	Association between uric acid and renal impairment in non-albuminuric diabetes kidney disease of type 2 diabetes.	O
4	Efficacy and safety of switching from febuxostat to dotinurad, a novel selective urate reabsorption inhibitor, in hyperuricemic patients with type 2 diabetic kidney disease: Protocol for a single-arm, open-label, prospective, exploratory study. 13,	O
3	Oral sub-chronic treatment with Terminalia phaeocarpa Eichler (Combretaceae) reduces liver PTP1B activity in a murine model of diabetes. <b>2023</b> , 306, 116164	O
2	The Renal Effects of SGLT2 Inhibitors. 76-83	O
1	Association between hyperuricemia and chronic kidney disease: a cross-sectional study in Bangladeshi adults. <b>2023</b> , 23,	1