

# CITATION REPORT

List of articles citing

Effects of flavored mineral waters on AhR-CYP1A1 signaling pathway in primary human hepatocytes and in human hepatic and intestinal cancer cells

DOI: 10.1016/j.fct.2012.03.073

Food and Chemical Toxicology, 2012, 50, 1933-9.

**Source:** <https://exaly.com/paper-pdf/53637033/citation-report.pdf>

**Version:** 2024-04-28

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
8	Effects of artificial sweeteners on the AhR- and GR-dependent CYP1A1 expression in primary human hepatocytes and human cancer cells. <i>Toxicology in Vitro</i> , <b>2013</b> , 27, 2283-8	3.6	10
7	Pelargonidin activates the AhR and induces CYP1A1 in primary human hepatocytes and human cancer cell lines HepG2 and LS174T. <i>Toxicology Letters</i> , <b>2013</b> , 218, 253-9	4.4	42
6	AhR expression is increased in hepatocellular carcinoma. <i>Journal of Molecular Histology</i> , <b>2013</b> , 44, 455-61	3.3	40
5	Profiling of enantiopure drugs towards aryl hydrocarbon (AhR), glucocorticoid (GR) and pregnane X (PXR) receptors in human reporter cell lines. <i>Chemico-Biological Interactions</i> , <b>2014</b> , 208, 64-76	5	4
4	2,3,7,8-Tetrachlorodibenzo-p-dioxin Mediated Cleft palate by Mouse Embryonic Palate Mesenchymal Cells. <i>Archives of Oral Biology</i> , <b>2016</b> , 71, 150-154	2.8	4
3	Effects of Flavored Nonalcoholic Beverages on Transcriptional Activities of Nuclear and Steroid Hormone Receptors: Proof of Concept for Novel Reporter Cell Line PAZ-PPAR $\alpha$ . <i>Journal of Agricultural and Food Chemistry</i> , <b>2018</b> , 66, 12066-12078	5.7	2
2	Taxifolin suppresses the malignant progression of gastric cancer by regulating the AhR/CYP1A1 signaling pathway. <i>International Journal of Molecular Medicine</i> , <b>2021</b> , 48,	4.4	0
1	Flavoured water consumption alters pharmacokinetic parameters and increases exposure of erlotinib and gefitinib in a preclinical study using Wistar rats. <i>PeerJ</i> , <b>2020</b> , 8, e9881	3.1	3