

Blind Seed Disease of Rye-Grass

Nature

153, 287-288

DOI: [10.1038/153287a0](https://doi.org/10.1038/153287a0)

Citation Report

#	ARTICLE	IF	CITATIONS
1	Blind-seed disease of rye-grass (<i>Phiala temulenta</i> Prill. & Delacr.). <i>Annals of Applied Biology</i> , 1945, 32, 329-343.	2.5	18
2	Recent developments in seed technology. <i>Botanical Review, The</i> , 1949, 15, 283-344.	3.9	3
3	THE SUSCEPTIBILITY OF PERENNIAL RYEGRASS TO BLIND SEED DISEASE (<i>Gleotinia* temulenta</i> (Prill, and) Tj ETQq0.0.0 rgBT /Overlock 1	2.9	8
4	SUSCEPTIBILITY OF PERENNIAL RYEGRASS TO BLIND SEED DISEASE (<i>GLOEOTINIA TEMULENTA</i>). <i>Grass and Forage Science</i> , 1958, 13, 275-278.	2.9	4
5	FLOWERING BEHAVIOUR, COMPATIBILITY AND MAJOR GENE DIFFERENCES IN <i>HOLCUS LANATUS</i> L.. <i>New Phytologist</i> , 1961, 60, 312-324.	7.3	6
6	Blind seed disease of ryegrass. <i>Euphytica</i> , 1967, 16, 122-130.	1.2	5
7	I. Occurrence and evidence for the use of nitrogen as a control measure. <i>New Zealand Journal of Agricultural Research</i> , 1980, 23, 143-147.	1.6	14
8	Fungal Diversity in Permafrost and Tallgrass Prairie Soils under Experimental Warming Conditions. <i>Applied and Environmental Microbiology</i> , 2013, 79, 7063-7072.	3.1	66