

High-resolution record of the environmental response to the
Last Interglacialâ€“Glacial cycle in Central Europe: the
VÄ›stonice (Czech Republic)

Quaternary Science Reviews

67, 17-38

DOI: [10.1016/j.quascirev.2013.01.014](https://doi.org/10.1016/j.quascirev.2013.01.014)

Citation Report

#	ARTICLE	IF	CITATIONS
1	Grass firesâ€”an unlikely process to explain the magnetic properties of prairie soils. <i>Geophysical Journal International</i> , 2013, 195, 1566-1575.	1.0	17
2	Modeling dust emission response to North Atlantic millennial-scale climate variations from the perspective of East European MIS 3 loess deposits. <i>Climate of the Past</i> , 2013, 9, 1385-1402.	1.3	46
3	Major dust events in Europe during marine isotope stage 5 (130â€”74 ka): a climatic interpretation of the "markers". <i>Climate of the Past</i> , 2013, 9, 2213-2230.	1.3	23
4	Geochronological reconsideration of the eastern European key loess section at Stayky in Ukraine. <i>Climate of the Past</i> , 2014, 10, 783-796.	1.3	9
6	European glacial dust deposits: Geochemical constraints on atmospheric dust cycle modeling. <i>Geophysical Research Letters</i> , 2014, 41, 7666-7674.	1.5	38
7	A simplified permafrost-carbon model for long-term climate studies with the CLIMBER-2 coupled earth system model. <i>Geoscientific Model Development</i> , 2014, 7, 3111-3134.	1.3	9
8	Geoarchaeology of Upper Palaeolithic loess sites located within a transect through Moravian valleys, Czech Republic. <i>Quaternary International</i> , 2014, 351, 25-37.	0.7	15
9	From micromorphology to palaeoenvironment: The MIS 10 to MIS 5 record in Paudorf (Lower Austria). <i>Catena</i> , 2014, 117, 60-72.	2.2	31
10	Holocene sedimentary systems on continental shelves. <i>Marine Geology</i> , 2014, 352, 268-294.	0.9	175
11	Climate variability and associated vegetation response throughout Central and Eastern Europe (CEE) between 60 and 8Åka. <i>Quaternary Science Reviews</i> , 2014, 106, 206-224.	1.4	188
12	Introducing an improved multi-proxy approach for paleoenvironmental reconstruction of loessâ€”paleosol archives applied on the Late Pleistocene Nussloch sequence (SW Germany). <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2014, 410, 300-315.	1.0	53
13	Soil burial contributes to deep soil organic carbon storage. <i>Soil Biology and Biochemistry</i> , 2014, 69, 251-264.	4.2	141
14	Landscape aridification in Central Germany during the late Weichselian Pleniglacial - results from the Zauschwitz loess site in western Saxony. <i>Zeitschrift FÅ¼r Geomorphologie</i> , 2014, 58, 27-50.	0.3	17
15	Earthworm calcite granules: a new tracker of millennialâ€”timescale environmental changes in Last Glacial loess deposits. <i>Journal of Quaternary Science</i> , 2015, 30, 529-536.	1.1	19
16	Palaeoenvironmental Interpretation of Yedoma Silt (Ice Complex) Deposition as Coldâ€”Climate Loess, Duvanny Yar, Northeast Siberia. <i>Permafrost and Periglacial Processes</i> , 2015, 26, 208-288.	1.5	110
17	Human-induced landscape evolution in the loess areas of Lublin Upland, E Poland: evidence from pedosedimentary archives in closed depressions. <i>Zeitschrift FÅ¼r Geomorphologie</i> , 2015, 59, 155-175.	0.3	4
18	The loess-palaeosol sequence in the Upper Palaeolithic site at KrakÃ³w Spadzista: A palaeoenvironmental approach. <i>Quaternary International</i> , 2015, 365, 98-113.	0.7	14
19	To meat or not to meat? New perspectives on <sc>N</sc>eanderthal ecology. <i>American Journal of Physical Anthropology</i> , 2015, 156, 43-71.	2.1	79

#	ARTICLE	IF	CITATIONS
20	Investigation of vegetation history of buried chernozem soils using near-infrared spectroscopy (NIRS). <i>Quaternary International</i> , 2015, 365, 203-211.	0.7	23
21	Danube loess stratigraphy – Towards a pan-European loess stratigraphic model. <i>Earth-Science Reviews</i> , 2015, 148, 228-258.	4.0	241
22	DolnÄ-VÄstonice Ila: Gravettian microstratigraphy, environment, and the origin of baked clay production in Moravia. <i>Quaternary International</i> , 2015, 359-360, 195-210.	0.7	25
23	The early Gravettian in a marginal area: New evidence from SW Poland. <i>Quaternary International</i> , 2015, 359-360, 131-152.	0.7	11
24	OSL chronostratigraphy of a loess-palaeosol sequence in ZÄota using quartz and polymineral fine grains. <i>Radiation Measurements</i> , 2015, 81, 23-31.	0.7	18
25	The origin of grasslands in the temperate forest zone of east-central Europe: long-term legacy of climate and human impact. <i>Quaternary Science Reviews</i> , 2015, 116, 15-27.	1.4	104
26	An integrated rock-magnetic and geochemical approach to loess/palaeosol sequences from Bohemia and Moravia (Czech Republic): Implications for the Upper Pleistocene paleoenvironment in central Europe. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2015, 418, 344-358.	1.0	58
27	Austrian MIS 3/2 loess – palaeosol records – Key sites along a west – east transect. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2015, 418, 43-56.	1.0	53
28	Loess, flints and bones: Multidisciplinary research at Jaksice II Gravettian site (southern Poland). <i>Quaternary International</i> , 2015, 359-360, 114-130.	0.7	29
29	Natural environment of MIS 5 and soil catena sequence along a loess slope in the Seret River valley: Evidence from the Pronyatyn Palaeolithic site (Ukraine). <i>Quaternary International</i> , 2015, 365, 74-97.	0.7	16
30	Environmental controls on paleo-pH at mid-latitudes: A case study from Central and Eastern Europe. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2015, 417, 458-466.	1.0	5
31	Late Pleistocene aeolian dust provenances and wind direction changes reconstructed by heavy mineral analysis of the sediments of the Dehner dry maar (Eifel, Germany). <i>Global and Planetary Change</i> , 2016, 147, 25-39.	1.6	28
32	Upper Pleistocene loess-palaeosol records from Northern France in the European context: Environmental background and dating of the Middle Palaeolithic. <i>Quaternary International</i> , 2016, 411, 4-24.	0.7	72
33	Pavlov I: A large Gravettian site in space and time. <i>Quaternary International</i> , 2016, 406, 95-105.	0.7	24
34	Late Pleistocene loess of the Lower Rhine. <i>Quaternary International</i> , 2016, 411, 44-61.	0.7	46
35	Loess-palaeosol sequences at the northern European loess belt in Germany: Distribution, geomorphology and stratigraphy. <i>Quaternary Science Reviews</i> , 2016, 153, 11-30.	1.4	96
36	Loess in Armenia – stratigraphic findings and palaeoenvironmental indications. <i>Proceedings of the Geologists Association</i> , 2016, 127, 29-39.	0.6	26
37	First results from the Late Pleistocene palaeosols in northern Western Siberia: Implications for pedogenesis and landscape evolution at the end of MIS3. <i>Quaternary International</i> , 2016, 418, 132-146.	0.7	17

#	ARTICLE	IF	CITATIONS
38	The physics of wind-blown loess: Implications for grain size proxy interpretations in Quaternary paleoclimate studies. <i>Earth-Science Reviews</i> , 2016, 154, 247-278.	4.0	170
39	The history of Danube loess research. <i>Quaternary International</i> , 2016, 399, 86-99.	0.7	27
40	Loess in a cave: Lithostratigraphic and correlative value of loess and loess-like layers in caves from the Kraków-Częstochowa Upland (Poland). <i>Quaternary International</i> , 2016, 399, 13-30.	0.7	29
41	Loess: Rock, sediment or soil – What is missing for its definition?. <i>Quaternary International</i> , 2016, 399, 198-207.	0.7	86
42	Middle Paleolithic sites of Katta Sai in western Tian Shan piedmont, Central Asiatic loess zone: Geoarchaeological investigation of the site formation and the integrity of the lithic assemblages. <i>Quaternary International</i> , 2016, 399, 136-150.	0.7	22
43	Comparative ¹⁴ C and OSL dating of loess-paleosol sequences to evaluate post-depositional contamination of <i>n</i> -alkane biomarkers. <i>Quaternary Research</i> , 2017, 87, 180-189.	1.0	20
44	(MIS3 & 2) millennial oscillations in Greenland dust and Eurasian aeolian records – A paleosol perspective. <i>Quaternary Science Reviews</i> , 2017, 169, 99-113.	1.4	59
45	Hominid visitation of the Moravian Karst during the Middle-Upper Paleolithic transition: New results from Pod Hradem Cave (Czech Republic). <i>Journal of Human Evolution</i> , 2017, 108, 131-146.	1.3	15
46	The loess sequence at Münzenberg (Wetterau/Germany): A reinterpretation based on new luminescence dating results. <i>Zeitschrift für Geomorphologie</i> , 2017, 61, 101-120.	0.3	7
47	Micromorphology of an Upper Paleolithic cultural layer at Grub-Kranawetberg, Austria. <i>Journal of Archaeological Science: Reports</i> , 2017, 14, 152-162.	0.2	3
48	The impact of Last Glacial climate variability in west-European loess revealed by radiocarbon dating of fossil earthworm granules. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2017, 114, 6209-6214.	3.3	93
49	Middle Pleniglacial pedogenesis on the northwestern edge of the Carpathian basin: A multidisciplinary investigation of the Băca pedo-sedimentary section, SW Slovakia. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2017, 487, 321-339.	1.0	22
50	Middle to Late Pleistocene multi-proxy record of environmental response to climate change from the Vienna Basin, Central Europe (Austria). <i>Quaternary Science Reviews</i> , 2017, 173, 193-210.	1.4	7
51	History of Czech Vegetation Since the Late Pleistocene. <i>Plant and Vegetation</i> , 2017, , 193-227.	0.6	13
52	The impact of natural and anthropogenic processes on the evolution of closed depressions in loess areas. A multi-proxy case study from Nałęczów Plateau, Eastern Poland. <i>Catena</i> , 2017, 149, 1-18.	2.2	17
53	The Eltville Tephra (Western Europe) age revised: Integrating stratigraphic and dating information from different Last Glacial loess localities. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2017, 466, 240-251.	1.0	19
54	A New Tool for Separating the Magnetic Mineralogy of Complex Mineral Assemblages from Low Temperature Magnetic Behavior. <i>Frontiers in Earth Science</i> , 2017, 5, .	0.8	29
55	Cave deposits as a sedimentary trap for the Marine Isotope Stage 3 environmental record: The case study of Pod Hradem, Czech Republic. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2018, 497, 201-217.	1.0	9

#	ARTICLE	IF	CITATIONS
56	Podhradem Interstadial; A critical review of the middle and late MIS 3 (Denekamp, Hengelo) in Moravia, Czech Republic. <i>Quaternary Science Reviews</i> , 2018, 182, 191-201.	1.4	6
57	The MIS 3/2 transition in a new loess profile at Krems-Wachtberg East – A multi-methodological approach. <i>Quaternary International</i> , 2018, 464, 370-385.	0.7	15
58	Last Glacial to Holocene vegetation succession recorded in polyphase slope-failure deposits on the Malenák Ridge, Outer Western Carpathians. <i>Quaternary International</i> , 2018, 470, 38-52.	0.7	15
59	Patterns and timing of loess-paleosol transitions in Eurasia: Constraints for paleoclimate studies. <i>Global and Planetary Change</i> , 2018, 162, 1-7.	1.6	35
60	Loess correlations – Between myth and reality. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2018, 509, 4-23.	1.0	31
61	The Crvenka loess-paleosol sequence: A record of continuous grassland domination in the southern Carpathian Basin during the Late Pleistocene. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2018, 509, 33-46.	1.0	38
62	Loess accumulation in the Tian Shan piedmont: Implications for palaeoenvironmental change in arid Central Asia. <i>Quaternary International</i> , 2018, 469, 30-43.	0.7	42
63	OSL chronologies of paleoenvironmental dynamics recorded by loess-paleosol sequences from Europe: Case studies from the Rhine-Meuse area and the Neckar Basin. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2018, 509, 105-125.	1.0	22
64	The Last Glacial aeolian record of the Island of Susak (Croatia) as seen from a high-resolution grain-size and rock magnetic analysis. <i>Quaternary International</i> , 2018, 494, 211-224.	0.7	26
65	Reconstruction of Late Pleistocene paleoenvironments in southern Germany using two high-resolution loess-paleosol records. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2018, 509, 58-76.	1.0	10
66	Short-term soil formation events in last glacial east European loess, evidence from multi-method luminescence dating. <i>Quaternary Science Reviews</i> , 2018, 200, 34-51.	1.4	34
68	Loess and other Quaternary sediments in Germany. <i>Journal of Maps</i> , 2018, 14, 330-340.	1.0	18
69	Approaches and challenges to the study of loess – Introduction to the LoessFest Special Issue. <i>Quaternary Research</i> , 2018, 89, 563-618.	1.0	92
70	Last Glacial aeolian landforms and deposits in the Rhône Valley (SE France): Spatial distribution and grain-size characterization. <i>Geomorphology</i> , 2018, 318, 250-269.	1.1	20
71	Paleoenvironmental conditions and sedimentation dynamics in Central Europe inferred from geochemical data of the loess-paleosol sequence at SÁ1/4ttÁ (Hungary). <i>Quaternary Science Reviews</i> , 2018, 196, 21-37.	1.4	16
72	European Loess Records – , 2018, , .		4
73	Integration of Linearbandkeramik cattle husbandry in the forested landscape of the mid-Holocene climate optimum: Seasonal-scale investigations in Bohemia. <i>Journal of Anthropological Archaeology</i> , 2018, 51, 16-27.	0.7	24
74	Chronostratigraphy of a loess-paleosol sequence in BiaÅy KoÅciÅ3Å,, Poland using OSL and radiocarbon dating. <i>Quaternary International</i> , 2019, 502, 4-17.	0.7	25

#	ARTICLE	IF	CITATIONS
75	Atmospheric Dust Variations in the Ili Basin, Northwest China, During the Last Glacial Period as Revealed by a High Mountain Loess-Paleosol Sequence. <i>Journal of Geophysical Research D: Atmospheres</i> , 2019, 124, 8449-8466.	1.2	41
76	Pleistocene landslides and mammoth bone deposits: The case of Dolnı-Vıstonice II, Czech Republic. <i>Geoarchaeology - an International Journal</i> , 2019, 34, 745-758.	0.7	7
77	A remarkable Late Saalian (MIS 6) loess (dust) accumulation in the Lower Danube at Harletz (Bulgaria). <i>Quaternary Science Reviews</i> , 2019, 207, 80-100.	1.4	16
78	Clumped isotope paleotemperatures from MIS 5 soil carbonates in southern Hungary. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2019, 518, 72-81.	1.0	14
79	A luminescence-based chronology for the Harletz loess sequence, Bulgaria. <i>Boreas</i> , 2019, 48, 179-194.	1.2	19
80	Landscape instability at the end of MIS 3 in western Central Europe: evidence from a multi proxy study on a Loess-Palaeosol-Sequence from the eastern Lower Rhine Embayment, Germany. <i>Quaternary International</i> , 2019, 502, 119-136.	0.7	17
81	High resolution dating of loess profile from Strzyıw (Horodıo Plateau-Ridge, Volhynia Upland). <i>Quaternary International</i> , 2019, 502, 18-29.	0.7	15
82	The Middle and Late Pleniglacial (Weichselian) malacofauna of the Zemun loess-paleosol sequence, Serbia. <i>Palaontologische Zeitschrift</i> , 2020, 94, 519-531.	0.8	5
83	Multiproxy evidence of middle and Late Pleistocene environmental changes in the loess-paleosol sequence of Bıhzdaı (Czech Republic). <i>Quaternary International</i> , 2020, 552, 4-14.	0.7	8
84	Geochemical signature of sources, recycling and weathering in the Last Glacial loess from the Rhıne Valley (southeast France) and comparison with other European regions. <i>Aeolian Research</i> , 2020, 42, 100561.	1.1	30
85	Paleoenvironments from robust loess stratigraphy using high-resolution color and grain-size data of the last glacial Krems-Wachtberg record (NE Austria). <i>Quaternary Science Reviews</i> , 2020, 248, 106602.	1.4	35
86	Geoheritage of East Kazakhstan. <i>Geoheritage</i> , 2020, 12, 1.	1.5	9
87	Origin and ¹⁰ Be surface exposure dating of a coarse debris accumulation in the Hrubı Jesenık Mountains, Central Europe. <i>Geomorphology</i> , 2020, 365, 107292.	1.1	3
88	Extratropical cyclones over the North Atlantic and western Europe during the Last Glacial Maximum and implications for proxy interpretation. <i>Climate of the Past</i> , 2020, 16, 611-626.	1.3	33
89	Shelter in Smoleı, III - A unique example of stratified Holocene clastic cave sediments in Central Europe, a lithostratigraphic stratotype and a record of regional paleoecology. <i>PLoS ONE</i> , 2020, 15, e0228546.	1.1	7
90	A chronology of landsliding based on archaeological and documentary data: Pavlovskı vrchy Hills, Western Carpathian Flysch Belt. <i>Scientific Reports</i> , 2020, 10, 976.	1.6	5
91	Interpretation of sedimentary subpopulations extracted from grain size distributions in loess deposits at the Sea of Azov, Russia. <i>Aeolian Research</i> , 2020, 45, 100597.	1.1	13
92	Smoothed millennial-scale palaeoclimatic reference data as unconventional comparison targets: Application to European loess records. <i>Scientific Reports</i> , 2020, 10, 5455.	1.6	8

#	ARTICLE	IF	CITATIONS
93	Environments at the MIS 3/2 transition in the northern Alps and their foreland. <i>Quaternary International</i> , 2021, 581-582, 99-113.	0.7	9
94	¹⁰ Be exposure age for sorted polygons in the Sudetes Mountains. <i>Permafrost and Periglacial Processes</i> , 2021, 32, 154-168.	1.5	3
95	A review of past changes in extratropical cyclones in the northern hemisphere and what can be learned for the future. <i>Wiley Interdisciplinary Reviews: Climate Change</i> , 2021, 12, .	3.6	15
96	The ¹³ C, ¹⁸ O and ⁴⁷ records in biogenic, pedogenic and geogenic carbonate types from paleosol-loess sequence and their paleoenvironmental meaning. <i>Quaternary Research</i> , 2021, 101, 256-272.	1.0	5
97	Chronological Assessment of the Balta Alba Kurgan Loess-Paleosol Section (Romania) – A Comparative Study on Different Dating Methods for a Robust and Precise Age Model. <i>Frontiers in Earth Science</i> , 2021, 8, .	0.8	16
98	Pleistocene dynamics of dust accumulation and soil formation in the southern Caspian Lowlands - New insights from the loess-paleosol sequence at Neka-Abelou, northern Iran. <i>Quaternary Science Reviews</i> , 2021, 253, 106774.	1.4	19
99	How dusty was the last glacial maximum over Europe?. <i>Quaternary Science Reviews</i> , 2021, 254, 106775.	1.4	11
100	Constraining the timing and deposition pattern of loess-palaeosol sequences in Kashmir Valley, Western Himalaya: Implications to paleoenvironment studies. <i>Aeolian Research</i> , 2021, 49, 100660.	1.1	7
101	New results concerning the pedo- and chronostratigraphy of the loess-palaeosol sequence Attenfeld (Bavaria, Germany) derived from a multi-methodological approach. <i>Journal of Quaternary Science</i> , 2021, 36, 1382.	1.1	4
102	The last 750 ka in loess-palaeosol sequences from northern France: environmental background and dating of the western European Palaeolithic. <i>Journal of Quaternary Science</i> , 2021, 36, 1293-1310.	1.1	12
103	Respiratory adaptation to climate in modern humans and Upper Palaeolithic individuals from Sungir and Mladec. <i>Scientific Reports</i> , 2021, 11, 7997.	1.6	13
104	Intra-interstadial environmental changes in Last Glacial loess revealed by molluscan assemblages from the Upper Palaeolithic site of Amiens-Renancourt 1 (Somme, France). <i>Journal of Quaternary Science</i> , 2021, 36, 1322-1340.	1.1	6
105	Loess landscapes of Europe – Mapping, geomorphology, and zonal differentiation. <i>Earth-Science Reviews</i> , 2021, 215, 103496.	4.0	104
106	What are the main factors limiting the distribution of <i>Bembix rostrata</i> (Hymenoptera: Crabronidae) at early-succession sites?. <i>Journal of Insect Conservation</i> , 2021, 25, 571-583.	0.8	4
107	Pedosedimentary record of MIS 5 as an interplay of climatic trends and local conditions: Multi-proxy evidence from the Palaeolithic site of Moravské Krumlov IV (Moravia, Czech Republic). <i>Catena</i> , 2021, 200, 105174.	2.2	6
108	Structure of the Late Pleistocene Climate Rhythm Inferred from the Detailed Soil-Sedimentation Archive of the Extraglacial Region of the East European Plain (Aleksandrov Quarry). <i>Stratigraphy and Geological Correlation</i> , 2021, 29, 368-387.	0.2	2
109	Separating provenance and palaeoclimatic signals from particle size and geochemistry of loess-palaeosol sequences using log-ratio transformation: Central European loess belt, Czech Republic. <i>Sedimentary Geology</i> , 2021, 419, 105907.	1.0	0
110	Pedosedimentary environments in the Caspian Lowland during MIS5 (Srednaya Akhtuba reference) Tj ETQq1 1 0.784314 rgBT/Overlock	0.7	13

#	ARTICLE	IF	CITATIONS
111	Aeolian dust dynamics in the Fergana Valley, Central Asia, since ~30ka inferred from loess deposits. <i>Geoscience Frontiers</i> , 2021, 12, 101180.	4.3	22
112	The Epigravettian chronology and the human population of eastern Central Europe during MIS2. <i>Quaternary Science Reviews</i> , 2021, 271, 107187.	1.4	6
113	Limestone Klippen of the Pavlov Hills. <i>World Geomorphological Landscapes</i> , 2016, , 373-384.	0.1	2
114	Long-Term Geomorphological History of the Czech Republic. <i>World Geomorphological Landscapes</i> , 2016, , 29-39.	0.1	4
115	Pedostratigraphy and chronology of the Late Pleistocene for the extra glacial area in the Central Russian Upland (reference section Aleksandrov quarry). <i>Catena</i> , 2020, 194, 104689.	2.2	17
116	Geomorphological evolution of the Petrovaradin Fortress Palaeolithic site (Novi Sad, Serbia). <i>Quaternary Research</i> , 2021, 103, 21-34.	1.0	6
117	Clay mineralogical composition and chemical properties of Haplic Luvisol developed on loess in the protected landscape area Litovelská Pomoravá. <i>European Journal of Soil Science</i> , 2021, 72, 1128-1142.	1.8	6
118	Lithological indicators of loess sedimentation of SW Poland. <i>Contemporary Trends in Geoscience</i> , 2017, 6, 94-111.	0.5	9
119	Luminescence chronostratigraphy for the loess deposits in Żółta, Poland. <i>Geochronometria</i> , 2018, 45, 44-55.	0.2	20
120	OSL chronostratigraphy for loess deposits from Tyszowce " Poland. <i>Geochronometria</i> , 2017, 44, 307-318.	0.2	16
121	From Chernozem to Luvisol or from Luvisol to Chernozem? A discussion about the relationships and limits of the two types of soils. A case study of the soil catena of Hrušov, Czechia. <i>Geografie-Sborník CGS</i> , 2020, 125, 473-500.	0.3	7
122	La séquence de Gravelle-Autoroute (Somme, France): une référence pour l'enregistrement des variations climatiques Quaternaires depuis 1Ma à l'Ouest de l'Europe. <i>Quaternaire</i> , 2020, , 205-230.	0.1	4
123	The middle/upper Würmian alluvio-loessic sequences of Quincieux and Lyon (Rhône-Alpes, France): first palaeoenvironmental interpretations and correlations. <i>Quaternaire</i> , 2017, , 423-453.	0.1	3
125	Palaeoenvironments during MIS3 and MIS2 inferred from lacustrine intercalations in the loess " palaeosol sequence at Bobingen (southern Germany). <i>E&G Quaternary Science Journal</i> , 2017, 66, 73-89.	0.2	8
127	PETROFYZIKÁLNĚ-CHARAKTERISTIKA SPRAĚ A FOSILNĚ-PĚDY V HLINĚKU U LITOVLE. <i>Geological Research in Moravia and Silesia</i> , 2019, 25, .	0.1	0
128	A History Of Fire At Hitchcock Nature Center, Western Iowa, Reconstructed Through An Analysis Of Macroscopic Charcoal And Soil-Magnetic Properties. <i>Journal of the Iowa Academy of Science</i> , 2020, 127, 13-22.	0.5	1
129	Dust Deposits: Loess. , 2022, , 320-365.		1
130	A tale of two signals: Global and local influences on the Late Pleistocene loess sequences in Bulgarian Lower Danube. <i>Quaternary Science Reviews</i> , 2021, 274, 107264.	1.4	16

#	ARTICLE	IF	CITATIONS
131	Vegetation and land snail-based reconstruction of the palaeocological changes in the forest steppe eco-region of the Carpathian Basin during last glacial warming. <i>Global Ecology and Conservation</i> , 2022, 33, e01976.	1.0	7
132	Detection of occupational surface remnants at a heavily eroded site; case study of archaeological soils from La Terrasse, Bibracte oppidum. <i>Catena</i> , 2022, 210, 105911.	2.2	1
133	Rock-magnetic characteristics and mechanisms of the Titel loess-paleosol sequence in northern Serbia since late Middle Pleistocene. <i>Science China Earth Sciences</i> , 2022, 65, 503-517.	2.3	3
134	Evaluation of geochemical proxies and radiocarbon data from a loess record of the Upper Palaeolithic site Kammern-Grubgraben, Lower Austria. <i>E&G Quaternary Science Journal</i> , 2022, 71, 23-43.	0.2	2
135	Half-precession signals in Lake Ohrid (Balkan) and their spatio-temporal relations to climate records from the European realm. <i>Quaternary Science Reviews</i> , 2022, 280, 107413.	1.4	9
136	Neanderthals in changing environments from MIS 5 to early MIS 4 in northern Central Europe – Integrating archaeological, (chrono)stratigraphic and paleoenvironmental evidence at the site of Lichtenberg. <i>Quaternary Science Reviews</i> , 2022, 284, 107519.	1.4	5
137	The Pampean region (Argentina) underwent larger variation in aridity than in temperature during the late Pleistocene: New evidence from the isotopic analysis of mammalian taxa. <i>Quaternary Science Reviews</i> , 2022, 286, 107555.	1.4	9
138	Environmental factors controlling the Last Glacial multi-phase development of the Moravian Sahara dune field, Lower Moravian Basin, Central Europe. <i>Geomorphology</i> , 2022, 413, 108355.	1.1	7
139	The myth of “Bohunician soil”: A re-evaluation of the MIS 3 palaeosol record at the Brno-Bohunice site (Czechia). <i>Catena</i> , 2022, 217, 106510.	2.2	1
140	Investigating the loess “palaeosol sequence of Bahlingen-SchÄ¶nberg (Kaiserstuhl), southwestern Germany, using a multi-methodological approach. <i>E&G Quaternary Science Journal</i> , 2022, 71, 145-162.	0.2	4
141	Amplified and suppressed regional imprints of global warming events on the southeastern Tibetan Plateau during MIS 3 “2. <i>Quaternary Science Reviews</i> , 2022, 294, 107736.	1.4	5
142	A pedogenically-informed chronostratigraphic model elucidates the geochronology at the type site of the Bohunician technocomplex. <i>Quaternary Science Reviews</i> , 2022, 297, 107827.	1.4	2