

CITATION REPORT

List of articles citing

miR-210 is a target of hypoxia-inducible factors 1 and 2 in renal cancer, regulates ISCU and correlates with good prognosis

DOI: 10.1038/bjc.2013.56

British Journal of Cancer, 2013, 108, 1133-42.

Source: <https://exaly.com/paper-pdf/55895680/citation-report.pdf>

Version: 2024-04-28

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
121	Identification of somatic VHL gene mutations in sporadic head and neck paragangliomas in association with activation of the HIF-1 α /miR-210 signaling pathway. 2013 , 98, E1661-6		15
120	The Crosstalk between Micro RNA and Iron Homeostasis. 2013 , 01,		2
119	Elevated expression of miR-210 predicts poor survival of cancer patients: a systematic review and meta-analysis. 2014 , 9, e89223		51
118	HIF-1 had pivotal effects on downregulation of miR-210 decreasing viability and inducing apoptosis in hypoxic chondrocytes. 2014 , 2014, 876363		13
117	MiR-210 expression reverses radioresistance of stem-like cells of oesophageal squamous cell carcinoma. 2014 , 5, 1068-77		10
116	Differential expression of microRNA-210 in gliomas of variable cell origin and correlation between increased expression levels and disease progression in astrocytic tumours. 2014 , 52, 79-85		3
115	18. Iron-sulfur proteins and human diseases.		
114	Elevated microRNA-185 is associated with high vascular endothelial growth factor receptor 2 expression levels and high microvessel density in clear cell renal cell carcinoma. 2014 , 35, 12757-63		10
113	Tumor miR-125b predicts recurrence and survival of patients with clear-cell renal cell carcinoma after surgical resection. 2014 , 105, 1427-34		21
112	miR-210 over-expression enhances mesenchymal stem cell survival in an oxidative stress environment through antioxidation and c-Met pathway activation. 2014 , 57, 989-97		37
111	MicroRNA-382 induced by HIF-1 α is an angiogenic miR targeting the tumor suppressor phosphatase and tensin homolog. 2014 , 42, 8062-72		100
110	Overexpression of miR-210 is associated with SDH-related pheochromocytomas, paragangliomas, and gastrointestinal stromal tumours. 2014 , 21, 415-26		27
109	miR-210 promotes IPF fibroblast proliferation in response to hypoxia. 2014 , 307, L283-94		59
108	Positive prognostic impact of miR-210 in non-small cell lung cancer. 2014 , 83, 272-8		47
107	MicroRNA-210 overexpression predicts poorer prognosis in glioma patients. 2014 , 21, 755-60		32
106	Predicting distant metastasis and chemoresistance using plasma miRNAs. 2014 , 31, 799		56
105	Methods of quantifying microRNAs for hypoxia research: classic and next generation. 2014 , 21, 1239-48		4

104	Effect of microRNA-210 on prognosis and response to chemotherapeutic drugs in pediatric acute lymphoblastic leukemia. 2014 , 105, 463-72	45
103	MicroRNAs in hypoxia and acidic tumor microenvironment. 2014 , 59, 2223-2231	5
102	Multiple functions of hypoxia-regulated miR-210 in cancer. 2014 , 33, 50	112
101	Vascular endothelial growth factor signaling in hypoxia and inflammation. 2014 , 9, 142-60	185
100	HypoxamiRs and cancer: from biology to targeted therapy. 2014 , 21, 1220-38	74
99	Oncogenic micro-RNAs and Renal Cell Carcinoma. 2014 , 4, 49	43
98	Downregulation of microRNA-210 inhibits osteosarcoma growth in vitro and in vivo. 2015 , 12, 3674-3680	9
97	MicroRNAs in Renal Cell Carcinoma. 2015 , 4, 26-35	27
96	MicroRNA-210 and Endoplasmic Reticulum Chaperones in the Regulation of Chemoresistance in Glioblastoma. 2015 , 6, 227-32	27
95	Prognostic Impact of Hypoxia-Inducible miRNA-210 in Patients with Lung Adenocarcinoma. 2015 , 2015, 316745	29
94	MicroRNAs as prognostic molecular signatures in renal cell carcinoma: a systematic review and meta-analysis. 2015 , 6, 32545-60	89
93	Prognostic value of meta-signature miRNAs in renal cell carcinoma: an integrated miRNA expression profiling analysis. 2015 , 5, 10272	59
92	Interactional role of microRNAs and bHLH-PAS proteins in cancer (Review). 2015 , 47, 25-34	4
91	miR-210 is a prognostic marker in clear cell renal cell carcinoma. 2015 , 17, 136-44	49
90	miR-133a mediates the hypoxia-induced apoptosis by inhibiting TAGLN2 expression in cardiac myocytes. 2015 , 400, 173-81	42
89	The administration of erythropoietin attenuates kidney injury induced by ischemia/reperfusion with increased activation of Wnt/ β -catenin signaling. 2015 , 114, 430-7	38
88	Personalized radiotherapy: concepts, biomarkers and trial design. 2015 , 88, 20150009	20
87	Age and sex differences in kidney microRNA expression during the life span of F344 rats. 2015 , 6, 1	47

86	Epigenetic programming of hypoxic-ischemic encephalopathy in response to fetal hypoxia. 2015 , 124, 28-48		33
85	Targeting tumour hypoxia to prevent cancer metastasis. From biology, biosensing and technology to drug development: the METOXIA consortium. 2015 , 30, 689-721		79
84	Integrative analysis of protein-coding and non-coding RNAs identifies clinically relevant subtypes of clear cell renal cell carcinoma. 2016 , 7, 82671-82685		11
83	MicroRNA-421 regulated by HIF-1 β promotes metastasis, inhibits apoptosis, and induces cisplatin resistance by targeting E-cadherin and caspase-3 in gastric cancer. 2016 , 7, 24466-82		86
82	MicroRNAs in the pathogenesis of renal cell carcinoma and their diagnostic and prognostic utility as cancer biomarkers. 2016 , 31, e26-37		17
81	MicroRNA-210 and its theranostic potential. 2016 , 20, 1325-1338		12
80	Neurotensin Promotes the Development of Colitis and Intestinal Angiogenesis via Hif-1 β miR-210 Signaling. 2016 , 196, 4311-21		27
79	Expression of miR-210 in relation to other measures of hypoxia and prediction of benefit from hypoxia modification in patients with bladder cancer. <i>British Journal of Cancer</i> , 2016 , 115, 571-8	8.7	16
78	MicroRNA-210 induces apoptosis in colorectal cancer via induction of reactive oxygen. 2016 , 16, 42		34
77	Hypoxic regulation of the noncoding genome and NEAT1. 2016 , 15, 174-85		36
76	The role of miRNAs in the pheochromocytomas. 2016 , 37, 4235-9		5
75	Hypoxia, Hypoxia-inducible Transcription Factors, and Renal Cancer. 2016 , 69, 646-657		158
74	Mitochondrial composition and function under the control of hypoxia. 2017 , 12, 208-215		241
73	The MicroRNA miR-210 Is Expressed by Cancer Cells but Also by the Tumor Microenvironment in Triple-Negative Breast Cancer. 2017 , 65, 335-346		21
72	Prognostic evaluation of microRNA-210 in various carcinomas: Evidence from 19 studies. 2017 , 96, e8113		8
71	HIF-1 β Promotes Breast Cancer Cell MCF-7 Proliferation and Invasion Through Regulating miR-210. 2017 , 32, 297-301		24
70	The epigenetic landscape of renal cancer. 2017 , 13, 47-60		70
69	10 Iron-sulfur proteins and human diseases. 2017 ,		

68	Evaluation of Plasma MicroRNAs as Diagnostic and Prognostic Biomarkers in Pancreatic Adenocarcinoma: miR-196a and miR-210 Could Be Negative and Positive Prognostic Markers, Respectively. 2017 , 2017, 6495867		21
67	Pheochromocytoma and paraganglioma: genotype versus anatomic location as determinants of tumor phenotype. 2018 , 372, 347-365		11
66	miRNA in tumour metabolism and why could it be the preferred pathway for energy reprogramming. 2018 , 17, 157-169		15
65	The biological roles and clinical implications of microRNAs in clear cell renal cell carcinoma. 2018 , 233, 4458-4465		31
64	miR-4326 promotes lung cancer cell proliferation through targeting tumor suppressor APC2. 2018 , 443, 151-157		13
63	miR-30a-5p together with miR-210-3p as a promising biomarker for non-small cell lung cancer: A preliminary study. 2018 , 21, 479-488		27
62	Hypoxia-induced miR-210 contributes to apoptosis of mouse spermatocyte GC-2 cells by targeting Kruppel-like factor 7. 2019 , 19, 271-279		9
61	Non-Coding Micro RNAs and Hypoxia-Inducible Factors Are Selenium Targets for Development of a Mechanism-Based Combination Strategy in Clear-Cell Renal Cell Carcinoma-Bench-to-Bedside Therapy. <i>International Journal of Molecular Sciences</i> , 2018 , 19,	6.3	15
60	Clinical utility of microRNAs in renal cell carcinoma: current evidence and future perspectives. 2018 , 18, 981-991		13
59	Genome-scale analysis to identify potential prognostic microRNA biomarkers for predicting overall survival in patients with colon adenocarcinoma. 2018 , 40, 1947-1958		12
58	Identification of cancer-related miRNA-lncRNA biomarkers using a basic miRNA-lncRNA network. 2018 , 13, e0196681		42
57	Alterations in Cellular Iron Metabolism Provide More Therapeutic Opportunities for Cancer. <i>International Journal of Molecular Sciences</i> , 2018 , 19,	6.3	47
56	Bioinformatic identification of key genes and analysis of prognostic values in clear cell renal cell carcinoma. 2018 , 16, 1747-1757		18
55	Epigenetics in renal cell cancer: mechanisms and clinical applications. 2018 , 15, 430-451		67
54	Dysfunction in the mitochondrial Fe-S assembly machinery leads to formation of the chemoresistant truncated VDAC1 isoform without HIF-1 α activation. 2018 , 13, e0194782		19
53	Discovery and Validation of Circulating Hsa-miR-210-3p as a Potential Biomarker for Primary Open-Angle Glaucoma. 2019 , 60, 2925-2934		10
52	miR-22 Regulates Invasion, Gene Expression and Predicts Overall Survival in Patients with Clear Cell Renal Cell Carcinoma. 2019 , 3, 119-132		5
51	MicroRNA-210-3p Targets RGMA to Enhance the Angiogenic Functions of Endothelial Progenitor Cells Under Hypoxic Conditions. 2019 , 13, 223		9

50	Molecular Mechanisms in Clear Cell Renal Cell Carcinoma: Role of miRNAs and Hypermethylated miRNA Genes in Crucial Oncogenic Pathways and Processes. 2019 , 10, 320	39
49	Iron regulatory protein 2 modulates the switch from aerobic glycolysis to oxidative phosphorylation in mouse embryonic fibroblasts. 2019 , 116, 9871-9876	16
48	Biomarkers of Prognosis and Efficacy of Anti-angiogenic Therapy in Metastatic Clear Cell Renal Cancer. 2019 , 9, 1400	21
47	A Five-MicroRNA Signature Predicts Survival and Disease Control of Patients with Head and Neck Cancer Negative for HPV Infection. 2019 , 25, 1505-1516	35
46	Roles of miR-210 in the pathogenesis of pre-eclampsia. 2019 , 15, 183-190	12
45	miRNA networks modulate human endothelial cell adaptation to cyclic hypoxia. 2019 , 54, 150-160	19
44	Emerging role of secreted miR-210-3p as potential biomarker for clear cell Renal Cell Carcinoma metastasis. 2020 , 27, 181-188	15
43	Circulating Non-coding RNAs in Renal Cell Carcinoma-Pathogenesis and Potential Implications as Clinical Biomarkers. 2020 , 8, 828	15
42	Morphing projections: a new visual technique for fast and interactive large-scale analysis of biomedical datasets. 2021 , 37, 1571-1580	0
41	Renal Cell Tumors: Uncovering the Biomarker Potential of ncRNAs. 2020 , 12,	9
40	Crosstalk between noncoding RNAs and ferroptosis: new dawn for overcoming cancer progression. 2020 , 11, 580	15
39	Adipose-derived exosomal miR-210/92a cluster inhibits adipose browning via the FGFR-1 signaling pathway in high-altitude hypoxia. 2020 , 10, 14390	6
38	MiR-210-3p Inhibits Proliferation and Migration of C6 Cells by Targeting Iscu. 2020 , 45, 1813-1824	1
37	Deletion of hypoxia-responsive microRNA-210 results in a sex-specific decrease in nephron number. <i>FASEB Journal</i> , 2020 , 34, 5782-5799	0.9 2
36	The interplay between HIF-1 α and noncoding RNAs in cancer. 2020 , 39, 27	24
35	Identification of RCC Subtype-Specific microRNAs-Meta-Analysis of High-Throughput RCC Tumor microRNA Expression Data. 2021 , 13,	4
34	Analysis of differential expression of hypoxia-inducible microRNA-210 gene targets in mild and severe preeclamptic patients. 2021 , 6, 51-57	2
33	The Ambivalent Role of miRNAs in Carcinogenesis: Involvement in Renal Cell Carcinoma and Their Clinical Applications. 2021 , 14,	4

32	Iron-sulphur cluster biogenesis factor LYRM4 is a novel prognostic biomarker associated with immune infiltrates in hepatocellular carcinoma. 2021 , 21, 463		1
31	miR-210 hypoxamiR in angiogenesis and diabetes. 2021 ,		1
30	Prognostic value of carbonic anhydrase IX immunohistochemical expression in renal cell carcinoma: a meta-analysis of the literature. 2014 , 9, e114096		23
29	MicroRNAs in clear cell renal cell carcinoma: biological functions and applications. 2015 , 2, 140-152		13
28	Role of VHL, HIF1A and SDH on the expression of miR-210: Implications for tumoral pseudo-hypoxic fate. 2017 , 8, 6700-6717		11
27	Clinically relevant HIF-1 β -dependent metabolic reprogramming in oropharyngeal squamous cell carcinomas includes coordinated activation of CAIX and the miR-210/ISCU signaling axis, but not MCT1 and MCT4 upregulation. 2017 , 8, 13730-13746		20
26	microRNA-210-3p depletion by CRISPR/Cas9 promoted tumorigenesis through revival of TWIST1 in renal cell carcinoma. 2017 , 8, 20881-20894		45
25	Suppression of homology-dependent DNA double-strand break repair induces PARP inhibitor sensitivity in -deficient human renal cell carcinoma. 2018 , 9, 4647-4660		14
24	Mitochondrial respiratory chain composition and organization in response to changing oxygen levels. 2020 , 2,		1
23	Epigenomics of clear cell renal cell carcinoma: mechanisms and potential use in molecular pathology. 2016 , 28, 80-91		39
22	MicroRNAs in Body Fluids: A More Promising Biomarker for Clear Cell Renal Cell Carcinoma. 2021 , 13, 7663-7675		2
21	miR-210 as a Biomarker in Renal Carcinoma. 2015 , 1-16		
20	miR-210 as a Biomarker in Renal Carcinoma. 2016 , 895-910		
19	Hypoxia-Regulated MicroRNAs in the Retina. <i>Advances in Experimental Medicine and Biology</i> , 2019 , 1185, 413-417	3.6	1
18	Noncoding RNAs and Its Implication as Biomarkers in Renal Cell Carcinoma: A Systematic Analysis. <i>Annals of Urologic Oncology</i> , 2019 , 1-11	0.1	1
17	An RNA-sequencing-based transcriptome for a significantly prognostic novel driver signature identification in bladder urothelial carcinoma. <i>PeerJ</i> , 2020 , 8, e9422	3.1	2
16	Can miRNAs be useful biomarkers in improving prognostic stratification in endometrial cancer patients? An update review. <i>International Journal of Cancer</i> , 2021 ,	7.5	0
15	Signaling Pathways Involved in Kidney and Urinary Tract Physiology and Pathology. 2020 , 163-193		

14	Prognostic role of microRNA-210 in various carcinomas: a meta-analysis. <i>International Journal of Clinical and Experimental Medicine</i> , 2015 , 8, 15283-9		8
13	miR-210 promotes progression of endometrial carcinoma by regulating the expression of NFIX. <i>International Journal of Clinical and Experimental Pathology</i> , 2018 , 11, 5213-5222	1.4	4
12	Nickel nanoparticle-induced cell transformation: involvement of DNA damage and DNA repair defect through HIF-1 α /miR-210/Rad52 pathway. <i>Journal of Nanobiotechnology</i> , 2021 , 19, 370	9.4	4
11	Circulating miR-210 and miR-23b in bladder Cancer. <i>Urological Science</i> , 2021 , 32, 64	0.3	0
10	Applications of noncoding RNAs in renal cancer patients. 2022 , 211-284		0
9	PRP4 Induces Epithelial-Mesenchymal Transition and Drug Resistance in Colon Cancer Cells via Activation of p53.. <i>International Journal of Molecular Sciences</i> , 2022 , 23,	6.3	2
8	Image_1.tif. 2019 ,		
7	Mitochondrial-Related Transcriptome Feature Correlates with Prognosis, Vascular Invasion, Tumor Microenvironment, and Treatment Response in Hepatocellular Carcinoma.. <i>Oxidative Medicine and Cellular Longevity</i> , 2022 , 2022, 1592905	6.7	0
6	miRNA Pattern in Hypoxic Microenvironment of Kidney Cancer Role of PTEN. <i>Biomolecules</i> , 2022 , 12, 686	5.9	0
5	The hypoxia-induced changes in miRNA-mRNA in RNA -induced silencing complexes and HIF -2 induced miRNAs in human endothelial cells. <i>FASEB Journal</i> , 2022 , 36,	0.9	1
4	Development of lactate-related gene signature and prediction of overall survival and chemosensitivity in patients with colorectal cancer.		0
3	Emerging role of miRNAs in the regulation of ferroptosis. 10,		0
2	Hypoxia-induced autophagy in triple negative breast cancer: association with prognostic variables, patients survival and response to neoadjuvant chemotherapy.		0
1	Targeting RNA with Small Molecules. 2023 , 1-33		0