

CITATION REPORT

List of articles citing

PDE5 inhibitors blunt inflammation in human BPH: a potential mechanism of action for PDE5 inhibitors in LUTS

DOI: 10.1002/pros.22686
Prostate, 2013, 73, 1391-402.

Source: <https://exaly.com/paper-pdf/56699908/citation-report.pdf>

Version: 2024-04-25

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
94	Risks and benefits of late onset hypogonadism treatment: an expert opinion. <i>World Journal of Men's Health</i> , 2013 , 31, 103-25	6.8	69
93	Benign prostatic hyperplasia: a new metabolic disease of the aging male and its correlation with sexual dysfunctions. <i>International Journal of Endocrinology</i> , 2014 , 2014, 329456	2.7	77
92	Editorial comment on Zhang et al.: Impact of metabolic syndrome on benign prostatic hyperplasia in elderly Chinese men. <i>Urologia Internationalis</i> , 2014 , 93, 247-8	1.9	
91	Microvascular dysfunction and efficacy of PDE5 inhibitors in BPH-LUTS. <i>Nature Reviews Urology</i> , 2014 , 11, 231-41	5.5	26
90	Acute effects of physical exercise and phosphodiesterase type 5 inhibition on serum 11 β -hydroxysteroid dehydrogenase related glucocorticoids metabolites: a pilot study. <i>Endocrine</i> , 2014 , 47, 952-8	4	7
89	Benign prostatic hyperplasia: a new metabolic disease?. <i>Journal of Endocrinological Investigation</i> , 2014 , 37, 313-22	5.2	104
88	Insulin resistance is an independent predictor of severe lower urinary tract symptoms and of erectile dysfunction: results from a cross-sectional study. <i>Journal of Sexual Medicine</i> , 2014 , 11, 2074-82	1.1	34
87	Effect of chronic Sildenafil treatment on the prostate of C57Bl/6 mice. <i>Tissue and Cell</i> , 2014 , 46, 439-49	2.7	6
86	Daily phosphodiesterase type 5 inhibitor therapy: a new treatment option for prostatitis/prostatodynia?. <i>BJU International</i> , 2014 , 113, 694-5	5.6	5
85	Management of benign prostatic hyperplasia: role of phosphodiesterase-5 inhibitors. <i>Drugs and Aging</i> , 2014 , 31, 425-39	4.7	10
84	Protective effect of tadalafil on the functional and structural changes of the rat ventral prostate caused by chronic pelvic ischemia. <i>Prostate</i> , 2015 , 75, 233-41	4.2	28
83	The role of phosphodiesterase-5 inhibitors in prostatic inflammation: a review. <i>Journal of Inflammation</i> , 2015 , 12, 54	6.7	23
82	Impact of cardiovascular risk factors and related comorbid conditions and medical therapy reported at baseline on the treatment response to tadalafil 5 mg once-daily in men with lower urinary tract symptoms associated with benign prostatic hyperplasia: an integrated analysis of four randomised, double-blind, placebo-controlled, clinical trials. <i>International Journal of Clinical Practice</i> , 2015 , 69, 1496-507	2.9	11
81	Recent advances in treatment for Benign Prostatic Hyperplasia. <i>F1000Research</i> , 2015 , 4,	3.6	8
80	Testosterone and benign prostatic hyperplasia. <i>Asian Journal of Andrology</i> , 2015 , 17, 212-6	2.8	33
79	A synopsis of drugs currently in preclinical and early clinical development for the treatment of benign prostatic hyperplasia. <i>Expert Opinion on Investigational Drugs</i> , 2015 , 24, 1059-73	5.9	7
78	Emerging links between non-neurogenic lower urinary tract symptoms secondary to benign prostatic obstruction, metabolic syndrome and its components: A systematic review. <i>International Journal of Urology</i> , 2015 , 22, 982-90	2.3	29

77	Chronic Pelvic Ischemia: Contribution to the Pathogenesis of Lower Urinary Tract Symptoms (LUTS): A New Target for Pharmacological Treatment?. <i>LUTS: Lower Urinary Tract Symptoms</i> , 2015 , 7, 1-8	1.9	26
76	Effect of a single treatment with tadalafil on blood flow in lower urinary tract tissues in rat models of bladder overdistension/emptying and abdominal aorta clamping/release. <i>European Journal of Pharmacology</i> , 2015 , 754, 92-7	5.3	13
75	Relationship between non-alcoholic fatty liver disease and benign prostatic hyperplasia/lower urinary tract symptoms: new insights from an Italian cross-sectional study. <i>World Journal of Urology</i> , 2015 , 33, 743-51	4	19
74	Connections between lower urinary tract symptoms related to benign prostatic enlargement and metabolic syndrome with its components: a systematic review and meta-analysis. <i>Aging Male</i> , 2015 , 18, 207-16	2.1	24
73	Benign prostatic hyperplasia and metabolic syndrome: the expanding evidences of a new disease of aging male. <i>Aging Male</i> , 2015 , 18, 133-4	2.1	10
72	Long-term phosphodiesterase 5 inhibitor administration reduces inflammatory markers and heat-shock proteins in cavernous tissue of Zucker diabetic fatty rat (ZDF/fa/fa). <i>International Journal of Impotence Research</i> , 2015 , 27, 182-90	2.3	10
71	Metabolic syndrome and benign prostatic enlargement: a systematic review and meta-analysis. <i>BJU International</i> , 2015 , 115, 24-31	5.6	141
70	Ejaculatory dysfunction in the treatment of lower urinary tract symptoms. <i>Translational Andrology and Urology</i> , 2016 , 5, 450-9	2.3	11
69	Inflammatory mediators in the development and progression of benign prostatic hyperplasia. <i>Nature Reviews Urology</i> , 2016 , 13, 613-26	5.5	109
68	Antimuscarinics, β Agonists, and Phosphodiesterase Inhibitors in the Treatment of Male Lower Urinary Tract Symptoms: An Evolving Paradigm. <i>Urologic Clinics of North America</i> , 2016 , 43, 337-49	2.9	3
67	The role of prostate inflammation and fibrosis in lower urinary tract symptoms. <i>American Journal of Physiology - Renal Physiology</i> , 2016 , 311, F817-F821	4.3	27
66	Latest Evidence on the Use of Phosphodiesterase Type 5 Inhibitors for the Treatment of Lower Urinary Tract Symptoms Secondary to Benign Prostatic Hyperplasia. <i>European Urology</i> , 2016 , 70, 124-133 ^{10.2}		85
65	Tadalafil reduces visceral adipose tissue accumulation by promoting preadipocytes differentiation towards a metabolically healthy phenotype: Studies in rabbits. <i>Molecular and Cellular Endocrinology</i> , 2016 , 424, 50-70	4.4	19
64	Heme oxygenase levels and metaflammation in benign prostatic hyperplasia patients. <i>World Journal of Urology</i> , 2016 , 34, 1183-92	4	13
63	Antioxidant Potential and Ability of Phloroglucinol to Decrease Formation of Advanced Glycation End Products Increase Efficacy of Sildenafil in Diabetes-Induced Sexual Dysfunction of Rats. <i>Sexual Medicine</i> , 2016 , 4, e104-12	2.7	23
62	Phosphodiesterase type 5 inhibitors: back and forward from cardiac indications. <i>Journal of Endocrinological Investigation</i> , 2016 , 39, 143-51	5.2	22
61	The regulation of transient receptor potential canonical 4 (TRPC4) channel by phosphodiesterase 5 inhibitor via the cyclic guanosine 3',5'-monophosphate. <i>Pflugers Archiv European Journal of Physiology</i> , 2017 , 469, 693-702	4.6	4
60	[Bladder storage and voiding dysfunctions : Side effects of drug therapy]. <i>Der Urologe</i> , 2017 , 56, 456-464		2

59	Relation between histological prostatitis and lower urinary tract symptoms and erectile function. <i>Prostate International</i> , 2017 , 5, 119-123	3.4	5
58	Lower urinary tract symptoms/benign prostatic hypertrophy and vascular function: Role of the nitric oxide-phosphodiesterase type 5-cyclic guanosine 3'5'Smonophosphate pathway. <i>International Journal of Urology</i> , 2017 , 24, 412-424	2.3	6
57	Benign prostatic enlargement can be influenced by metabolic profile: results of a multicenter prospective study. <i>BMC Urology</i> , 2017 , 17, 22	2.2	19
56	Emerging drugs for the treatment of benign prostatic hyperplasia. <i>Expert Opinion on Emerging Drugs</i> , 2017 , 22, 201-212	3.7	11
55	The influence of asymptomatic inflammatory prostatitis on the onset and progression of lower urinary tract symptoms in men with histologic benign prostatic hyperplasia. <i>Asian Journal of Urology</i> , 2017 , 4, 158-163	2.7	13
54	The effect of doxazosin and sildenafil citrate combination on bladder tissue contractility, alpha adrenergic receptor, and iNOS subtype expression in a male rat model of partially bladder outlet obstruction. <i>Neurourology and Urodynamics</i> , 2017 , 36, 1479-1487	2.3	3
53	The link between vascular dysfunction, bladder ischemia, and aging bladder dysfunction. <i>Therapeutic Advances in Urology</i> , 2017 , 9, 11-27	3.2	61
52	The effect of tadalafil therapy on kidney damage caused by sepsis in a polymicrobial septic model induced in rats: a biochemical and histopathological study. <i>International Braz J Urol: Official Journal of the Brazilian Society of Urology</i> , 2017 , 43, 345-355	2	9
51	Singapore Urological Association Clinical Guidelines for Male Lower Urinary Tract Symptoms/Benign Prostatic Hyperplasia. <i>Singapore Medical Journal</i> , 2017 , 58, 473-480	1.9	8
50	Serenoa repens + selenium + lycopene vs tadalafil 5 mg for the treatment of lower urinary tract symptoms secondary to benign prostatic obstruction: a Phase IV, non-inferiority, open-label, clinical study (SPRITE study). <i>BJU International</i> , 2018 , 122, 317-325	5.6	7
49	Effect of tadalafil on chronic pelvic pain and prostatic inflammation in a rat model of experimental autoimmune prostatitis. <i>Prostate</i> , 2018 , 78, 707-713	4.2	18
48	Urine miR-21-5p as a potential biomarker for predicting effectiveness of tadalafil in benign prostatic hyperplasia. <i>Future Science OA</i> , 2018 , 4, FSO304	2.7	3
47	Prostatic inflammation: a potential treatment target for male LUTS due to benign prostatic obstruction. <i>Prostate Cancer and Prostatic Diseases</i> , 2018 , 21, 161-167	6.2	6
46	Tadalafil Improves Symptoms, Erectile Function and Quality of Life in Patients with Lower Urinary Tract Symptoms Suggestive of Benign Prostatic Hyperplasia (KYU-PRO Study). <i>LUTS: Lower Urinary Tract Symptoms</i> , 2018 , 10, 76-83	1.9	4
45	Association between metabolic syndrome and intravesical prostatic protrusion in patients with benign prostatic enlargement and lower urinary tract symptoms (MIPS Study). <i>BJU International</i> , 2018 , 121, 799-804	5.6	11
44	Metabolic Syndrome and LUTS/BPH. 2018 , 89-111		0
43	Efficacy and safety of PDE5-Is and α_1 blockers for treating lower ureteric stones or LUTS: a meta-analysis of RCTs. <i>BMC Urology</i> , 2018 , 18, 30	2.2	5
42	Oxidative stress marker 8-hydroxyguanosine is more highly expressed in prostate cancer than in benign prostatic hyperplasia. <i>Molecular and Clinical Oncology</i> , 2018 , 9, 302-304	1.6	7

41	Objective impacts of tadalafil on storage and voiding function in male patients with benign prostatic hyperplasia: 1-year outcomes from a prospective urodynamic study. <i>World Journal of Urology</i> , 2019 , 37, 867-872	4	12
40	Benign Prostatic Hyperplasia and Lower Urinary Tract Symptoms: What Is the Role and Significance of Inflammation?. <i>Current Urology Reports</i> , 2019 , 20, 54	2.9	19
39	Tadalafil 5 mg Alone or in Combination with Tamsulosin 0.4 mg for the Management of Men with Lower Urinary Tract Symptoms and Erectile Dysfunction: Results of a Prospective Observational Trial. <i>Journal of Clinical Medicine</i> , 2019 , 8,	5.1	7
38	A phosphodiesterase 5 inhibitor, tadalafil, suppresses stromal predominance and inflammation in a rat model of nonbacterial prostatitis. <i>BMC Urology</i> , 2019 , 19, 99	2.2	8
37	The Complex Relationship Between Lower Urinary Tract Symptoms and Sexual Health. <i>Current Urology Reports</i> , 2019 , 20, 58	2.9	4
36	Tadalafil for the treatment of benign prostatic hyperplasia. <i>Expert Opinion on Pharmacotherapy</i> , 2019 , 20, 929-937	4	12
35	Toward a Scientific Nutritional Supplement Combination for Prostatism and Erectile Dysfunction I: From Known Pharmacology to Clinical Testing. <i>Journal of Medicinal Food</i> , 2019 , 22, 529-537	2.8	1
34	The association between local arteriosclerosis of the prostatic arteries and chronic inflammation in human benign prostatic enlargement. <i>Prostate</i> , 2019 , 79, 574-582	4.2	5
33	Activation of cGMP/PKG/p65 signaling associated with PDE5-Is downregulates CCL5 secretion by CD8 T cells in benign prostatic hyperplasia. <i>Prostate</i> , 2019 , 79, 909-919	4.2	10
32	Tadalafil Treatment Improves Inflammation, Cognitive Function, And Mismatch Negativity Of Patients With Low Urinary Tract Symptoms And Erectile Dysfunction. <i>Scientific Reports</i> , 2019 , 9, 17119	4.9	11
31	Attenuation of Benign Prostatic Hyperplasia by Optimized Tadalafil Loaded Pumpkin Seed Oil-Based Self Nanoemulsion: In Vitro and In Vivo Evaluation. <i>Pharmaceutics</i> , 2019 , 11,	6.4	14
30	Efficacy of tadalafil against lower urinary tract symptoms after low-dose-rate brachytherapy in prostate cancer patients. <i>Journal of Clinical Urology</i> , 2019 , 12, 223-227	0.2	1
29	Sildenafil, a phosphodiesterase type 5 inhibitor, augments sphincter bursting and bladder afferent activity to enhance storage function and voiding efficiency in mice. <i>BJU International</i> , 2019 , 124, 163-173	5.6	6
28	Editorial Comment from Dr Tanaka to Tadalafil is sufficiently effective for severe chronic prostatitis/chronic pelvic pain syndrome in patients with benign prostatic hyperplasia. <i>International Journal of Urology</i> , 2020 , 27, 58-59	2.3	
27	Tadalafil is sufficiently effective for severe chronic prostatitis/chronic pelvic pain syndrome in patients with benign prostatic hyperplasia. <i>International Journal of Urology</i> , 2020 , 27, 53-57	2.3	11
26	Add-on effects of tadalafil in tamsulosin-treated patients with small benign prostatic enlargement: A randomized, placebo-controlled, double-blind, crossover study. <i>Neurourology and Urodynamics</i> , 2020 , 39, 237-242	2.3	2
25	Changes in the expression and function of the PDE5 pathway in the obstructed urinary bladder. <i>Journal of Cellular and Molecular Medicine</i> , 2020 , 24, 13181-13195	5.6	2
24	Effect and mechanism of prophylactic use of tadalafil during pregnancy on l-NAME-induced preeclampsia-like rats. <i>Placenta</i> , 2020 , 99, 35-44	3.4	3

23	Tadalafil Alone or in Combination with Tamsulosin for the Management for LUTS/BPH and ED. <i>Current Urology Reports</i> , 2020 , 21, 56	2.9	11
22	Tadalafil Improves Nocturia and Nocturia-Related Quality of Life in Patients with Benign Prostatic Hyperplasia (KYU-PRO Study). <i>Urologia Internationalis</i> , 2020 , 104, 587-593	1.9	4
21	Inflammation is a target of medical treatment for lower urinary tract symptoms associated with benign prostatic hyperplasia. <i>World Journal of Urology</i> , 2020 , 38, 2771-2779	4	16
20	Higher expression of phosphodiesterase type 5 in the anterior fibromuscular stroma of the human prostate. <i>World Journal of Urology</i> , 2020 , 38, 2915-2921	4	0
19	Phosphodiesterase 5 (PDE5): Structure-function regulation and therapeutic applications of inhibitors. <i>Biomedicine and Pharmacotherapy</i> , 2021 , 134, 111128	7.5	15
18	Which Drug to Discontinue 3 Months After Combination Therapy of Tadalafil plus Tamsulosin for Men with Lower Urinary Tract Symptom and Erectile Dysfunction? Results of a Prospective Observational Trial. <i>European Urology Focus</i> , 2021 , 7, 432-439	5.1	2
17	Bladder Outlet Obstruction and Overactive Bladder in Males. <i>Urodynamics, Neurourology and Pelvic Floor Dysfunctions</i> , 2021 , 67-85	0.1	
16	Impact of low-dose tadalafil on adverse events after low-dose-rate brachytherapy for prostate cancer: A bi-center randomized open-label trial. <i>International Journal of Urology</i> , 2021 , 28, 432-438	2.3	1
15	Lower urinary tract symptoms, erectile dysfunction, and their correlation in men aged 50 years and above: a cross-sectional survey in Beijing, China. <i>Medical Science Monitor</i> , 2014 , 20, 2806-10	3.2	12
14	Influence of metabolic syndrome on the relationship between fatty acids and the selected parameters in men with benign prostatic hyperplasia. <i>Aging</i> , 2019 , 11, 1524-1536	5.6	3
13	Clinical correlates of enlarged prostate size in subjects with sexual dysfunction. <i>Asian Journal of Andrology</i> , 2014 , 16, 767-73	2.8	8
12	Clinical and preclinical treatment of urologic diseases with phosphodiesterase isoenzymes 5 inhibitors: an update. <i>Asian Journal of Andrology</i> , 2016 , 18, 723-31	2.8	8
11	Non-Hormonal treatment of BPH/BOO. <i>Indian Journal of Urology</i> , 2014 , 30, 194-201	0.8	1
10	Antinociceptive Effect of Vardenafil on Carrageenan-Induced Hyperalgesia in Rat: involvement of Nitric Oxide/Cyclic Guanosine Monophosphate/Calcium Channels Pathway. <i>Iranian Journal of Pharmaceutical Research</i> , 2015 , 14, 1137-43	1.1	5
9	Tadalafil treatment improves cardiac, renal and lower urinary tract dysfunctions in rats with heart failure.. <i>Life Sciences</i> , 2021 , 289, 120237	6.8	1
8	Use of a Fission Yeast Platform to Identify and Characterize Small Molecule PDE Inhibitors.. <i>Frontiers in Pharmacology</i> , 2021 , 12, 833156	5.6	
7	Outcomes of combination therapy with daily tadalafil 5 mg plus tamsulosin 0.4 mg to treat lower urinary tract symptoms and erectile dysfunction in men with or without metabolic syndrome. <i>Minerva Urology and Nephrology</i> , 2022 , 73,	2.3	0
6	Benign Prostatic Hyperplasia/Obstruction Ameliorated Using a Soluble Guanylate Cyclase Activator.. <i>Journal of Pathology</i> , 2021 ,	9.4	0

5	Testosterone does not affect lower urinary tract symptoms while improving markers of prostatitis in men with benign prostatic hyperplasia: a randomized clinical trial.. <i>Journal of Endocrinological Investigation</i> , 2022 ,	5.2	
4	Efficacy Evaluation of Plant Products in the Treatment of Erectile Dysfunction Related to Diabetes.. <i>Nutrients</i> , 2021 , 13,	6.7	1
3	Ligand-Enhanced Negative Images Optimized for Docking Rescoring. <i>International Journal of Molecular Sciences</i> , 2022 , 23, 7871	6.3	1
2	Efficacy of Combination Treatment with Tadalafil and Mirabegron in Patients with Benign Prostatic Hyperplasia Who Presented with Persistent Storage Symptoms After Tadalafil Monotreatment: A Prospective, Multicenter, Open-Labeled Study. 2022 , 65, 231-237		0
1	A Novel Insight into the Immune-Related Interaction of Inflammatory Cytokines in Benign Prostatic Hyperplasia. 2023 , 12, 1821		0