

CITATION REPORT

List of articles citing

The Efficacy and Safety of Tadalafil 5 mg Once Daily in Korean Men with Lower Urinary Tract Symptoms Suggestive of Benign Prostatic Hyperplasia: An Integrated Analysis

DOI: 10.5534/wjmh.2014.32.1.28

World Journal of Men's Health, 2014, 32, 28-35.

Source: <https://exaly.com/paper-pdf/57563450/citation-report.pdf>

Version: 2024-04-23

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
11	The efficacy of mirodenafil for chronic prostatitis/chronic pelvic pain syndrome in middle-aged males. <i>World Journal of Men's Health</i> , 2014 , 32, 145-50	6.8	11
10	Impact of cardiovascular risk factors and related comorbid conditions and medical therapy reported at baseline on the treatment response to tadalafil 5 mg once-daily in men with lower urinary tract symptoms associated with benign prostatic hyperplasia: an integrated analysis of four randomised, double-blind, placebo-controlled clinical trials. <i>International Journal of Clinical Practice</i> 2015 , 69, 1496-507	2.9	11
9	Tadalafil 5 mg once daily for the treatment of Asian men with lower urinary tract symptoms secondary to benign prostatic hyperplasia: analyses of data pooled from three randomized, double-blind, placebo-controlled studies. <i>International Journal of Urology</i> , 2015 , 22, 378-84	2.3	21
8	Tadalafil for lower urinary tract symptoms secondary to benign prostatic hyperplasia: a review of clinical data in Asian men and an update on the mechanism of action. <i>Therapeutic Advances in Urology</i> , 2015 , 7, 249-64	3.2	17
7	Sexual Dysfunction Related to Drugs: a Critical Review. Part V: β -Blocker and 5-ARI Drugs. <i>Pharmacopsychiatry</i> , 2016 , 49, 3-13	2	29
6	Evaluation of hyoscine N-butyl bromide efficacy on the prevention of catheter-related bladder discomfort after transurethral resection of prostate: a randomized, double-blind control trial. <i>International Urology and Nephrology</i> , 2017 , 49, 1907-1913	2.3	4
5	Phosphodiesterase inhibitors for lower urinary tract symptoms consistent with benign prostatic hyperplasia. <i>The Cochrane Library</i> , 2018 , 11, CD010060	5.2	8
4	Additive effect of oral LDD175 to tamsulosin and finasteride in a benign prostate hyperplasia rat model. <i>Drug Design, Development and Therapy</i> , 2018 , 12, 1855-1863	4.4	5
3	Safety and Effectiveness of Once-Daily Tadalafil (5 mg) Therapy in Korean Men with Benign Prostatic Hyperplasia/Lower Urinary Tract Symptoms in a Real-World Clinical Setting: Results from a Post-Marketing Surveillance Study. <i>World Journal of Men's Health</i> , 2018 , 36, 161-170	6.8	9
2	Efficacy and safety of tadalafil 5mg once-daily in Asian men with both lower urinary tract symptoms associated with benign prostatic hyperplasia and erectile dysfunction: A phase III, randomized, double-blind, parallel, placebo- and tamsulosin-controlled study. <i>International Journal of Urology</i> , 2019 , 26, 192-200	2.3	11
1	Evaluation of the effectiveness and safety of a daily dose of 5 mg of tadalafil, over an 8-week period, for improving quality of life among Korean men with andropause symptoms, including erectile dysfunction: A pilot study. <i>Medicine (United States)</i> , 2018 , 97, e13827	1.8	4