

Changes of ventricular and peripheral performance in patients with normal ejection fraction: insights from ergometry stress

European Journal of Heart Failure

16, 888-897

DOI: [10.1002/ejhf.124](https://doi.org/10.1002/ejhf.124)

Citation Report

#	ARTICLE	IF	CITATIONS
1	Association of impaired left ventricular twisting—untwisting with vascular dysfunction, neurohumoral activation and impaired exercise capacity in hypertensive heart disease. <i>European Journal of Heart Failure</i> , 2015, 17, 1240-1251.	2.9	63
2	Clinical perspectives and evidence of diastolic stress test in heart failure with preserved ejection fraction. <i>Egyptian Heart Journal</i> , 2015, 67, 279-288.	0.4	1
3	Left ventricular long-axis performance during exercise is an important prognosticator in patients with heart failure and preserved ejection fraction. <i>International Journal of Cardiology</i> , 2015, 178, 131-135.	0.8	46
4	Importance of chronotropic response and left ventricular long-axis function for exercise performance in patients with heart failure and preserved ejection fraction. <i>International Journal of Cardiology</i> , 2016, 202, 339-343.	0.8	9
5	Impaired left ventricular global longitudinal strain in patients with heart failure with preserved ejection fraction: insights from the <sc>RELAX</sc> trial. <i>European Journal of Heart Failure</i> , 2017, 19, 893-900.	2.9	123
6	Contractile reserve and cardiopulmonary exercise parameters in patients with dilated cardiomyopathy, the two dimensions of exercise testing. <i>Echocardiography</i> , 2017, 34, 1179-1186.	0.3	8
7	Value of Strain Imaging and Maximal Oxygen Consumption in Patients With Hypertrophic Cardiomyopathy. <i>American Journal of Cardiology</i> , 2017, 120, 1203-1208.	0.7	10
8	The fallacy of resting echocardiographic parameters of cardiac function in heart failure with preserved ejection fraction. <i>European Journal of Heart Failure</i> , 2018, 20, 619-619.	2.9	2
9	Saudi Heart Association (SHA) guidelines for the management of heart failure. <i>Journal of the Saudi Heart Association</i> , 2019, 31, 204-253.	0.2	9
10	Obstructive sleep apnoea, intermittent hypoxia and heart failure with a preserved ejection fraction. <i>Heart</i> , 2021, 107, 190-194.	1.2	26
11	The Relationship Between Oxygen Uptake and the Rate of Myocardial Deformation During Exercise. <i>Bioengineered</i> , 2021, 10, 85-93.	1.4	0
12	Determining the thresholds for abnormal left ventricular strains in healthy subjects by echocardiography: a meta-analysis. <i>Cardiovascular Diagnosis and Therapy</i> , 2020, 10, 1858-1873.	0.7	7
14	Exercise Systolic Reserve and Exercise Pulmonary Hypertension Improve Diagnosis of Heart Failure With Preserved Ejection Fraction. <i>Frontiers in Cardiovascular Medicine</i> , 2022, 9, 814601.	1.1	1
15	Hemodynamic Changes During Physiological and Pharmacological Stress Testing in Patients With Heart Failure: A Systematic Review and Meta-Analysis. <i>Frontiers in Cardiovascular Medicine</i> , 2022, 9, 718114.	1.1	0
16	Additive effects of type 2 diabetes and metabolic syndrome on left ventricular torsion and linear deformation abnormalities during dobutamine stress echocardiography. <i>Frontiers in Cardiovascular Medicine</i> , 0, 9, .	1.1	1
17	Does global longitudinal strain improve stratification risk in heart failure with preserved ejection fraction?. <i>Journal of Cardiology and Cardiovascular Medicine</i> , 2022, 7, 074-080.	0.1	0