

CITATION REPORT

List of articles citing

Tadalafil 5 mg once-daily therapy for men with lower urinary tract symptoms suggestive of benign prostatic hyperplasia: results from a randomized, double-blind, placebo-controlled trial carried out in Japan and Korea

DOI: 10.1111/iju.12410

International Journal of Urology, 2014, 21, 670-5.

Source: <https://exaly.com/paper-pdf/58289891/citation-report.pdf>

Version: 2024-04-27

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
38	The Efficacy and Safety of Tadalafil 5 mg Once Daily in Korean Men with Lower Urinary Tract Symptoms Suggestive of Benign Prostatic Hyperplasia: An Integrated Analysis. <i>World Journal of Men's Health</i> , 2014 , 32, 28-35	6.8	11
37	Editorial comment to Tadalafil 5 mg once-daily therapy for men with lower urinary tract symptoms suggestive of benign prostatic hyperplasia: results from a randomized, double-blind, placebo-controlled trial carried out in Japan and Korea. <i>International Journal of Urology</i> , 2014 , 21, 676	2.3	
36	Treatment satisfaction and clinically meaningful symptom improvement in men with lower urinary tract symptoms and prostatic enlargement secondary to benign prostatic hyperplasia: Secondary results from a 6-month, randomized, double-blind study comparing finasteride plus tadalafil with	2.3	20
35	Editorial Comment from Dr Masumori to Treatment satisfaction and clinically meaningful symptom improvement in men with lower urinary tract symptoms and prostatic enlargement secondary to benign prostatic hyperplasia: Secondary results from a 6-month, randomized, double-blind study	2.3	
34	Comparing finasteride plus tadalafil with finasteride plus placebo. <i>International Journal of Urology</i> , 2015 , 22, 589 Tadalafil 5 mg once daily for the treatment of Asian men with lower urinary tract symptoms secondary to benign prostatic hyperplasia: analyses of data pooled from three randomized, double-blind, placebo-controlled studies. <i>International Journal of Urology</i> , 2015 , 22, 378-84	2.3	21
33	Tadalafil for lower urinary tract symptoms secondary to benign prostatic hyperplasia: a review of clinical data in Asian men and an update on the mechanism of action. <i>Therapeutic Advances in Urology</i> , 2015 , 7, 249-64	3.2	17
32	EFFECT OF TADALAFIL THAT WAS ADDITIONALLY ADMINISTERED TO PATIENTS RECEIVING AN ALPHA1-BLOCKER IN JAPANESE MEN WITH LOWER URINARY TRACT SYMPTOMS SUGGESTIVE OF BENIGN PROSTATIC HYPERPLASIA. <i>Japanese Journal of Urology</i> , 2016 , 107, 28-33		2
31	Geriatric Sexuality. 2016 , 163-227		
30	Effects of Sildenafil, a Phosphodiesterase Type 5 Inhibitor, on the Primary Single Afferent Activity of the Rat Bladder. <i>LUTS: Lower Urinary Tract Symptoms</i> , 2017 , 9, 57-61	1.9	2
29	Comparison of Silodosin Monotherapy vs Silodosin With Tadalafil Add-on Therapy in Patients With Benign Prostatic Hyperplasia. <i>Urology</i> , 2017 , 106, 153-159	1.6	3
28	Comparison of Silodosin versus Tadalafil in Patients with Lower Urinary Tract Symptoms Associated with Benign Prostatic Hyperplasia. <i>LUTS: Lower Urinary Tract Symptoms</i> , 2017 , 9, 176-186	1.9	6
27	Safety and efficacy of the combination of once-daily tadalafil and alpha-1 blocker in Japanese men with lower urinary tract symptoms suggestive of benign prostatic hyperplasia: A randomized, placebo-controlled, cross-over study. <i>International Journal of Urology</i> , 2017 , 24, 539-547	2.3	14
26	Efficacy and safety of tadalafil 5 mg once daily in the treatment of lower urinary tract symptoms associated with benign prostatic hyperplasia in men aged ≥5 years: integrated analyses of pooled data from multinational, randomized, placebo-controlled clinical studies. <i>BJU International</i> , 2017 , 119, 793-803	5.6	19
25	Clinical guidelines for male lower urinary tract symptoms and benign prostatic hyperplasia. <i>International Journal of Urology</i> , 2017 , 24, 716-729	2.3	63
24	Tadalafil 5 mg Once Daily Improves Lower Urinary Tract Symptoms and Erectile Dysfunction: A Systematic Review and Meta-analysis. <i>LUTS: Lower Urinary Tract Symptoms</i> , 2018 , 10, 84-92	1.9	17
23	Tadalafil Improves Symptoms, Erectile Function and Quality of Life in Patients with Lower Urinary Tract Symptoms Suggestive of Benign Prostatic Hyperplasia (KYU-PRO Study). <i>LUTS: Lower Urinary Tract Symptoms</i> , 2018 , 10, 76-83	1.9	4
22	Early improvement of storage or voiding symptoms by tadalafil predicts treatment outcomes in patients with lower urinary tract symptoms from benign prostatic hyperplasia. <i>International Journal of Urology</i> , 2018 , 25, 240-245	2.3	6

21	Safety and Effectiveness of Once-Daily Tadalafil (5 mg) Therapy in Korean Men with Benign Prostatic Hyperplasia/Lower Urinary Tract Symptoms in a Real-World Clinical Setting: Results from a Post-Marketing Surveillance Study. <i>World Journal of Men's Health</i> , 2018 , 36, 161-170	6.8	9
20	Tadalafil, a phosphodiesterase type 5 inhibitor, restores urethra and detrusor function in the initial phase of diabetes in rats. <i>LUTS: Lower Urinary Tract Symptoms</i> , 2019 , 11, 241-247	1.9	4
19	Effect of tadalafil 5mg on post-micturition dribble in men with lower urinary tract symptoms: a multicentre, double-blind, randomized, placebo-controlled trial. <i>BJU International</i> , 2019 , 124, 862-869	5.6	4
18	[EFFECTIVENESS AND SAFETY OF TADALAFIL IN VERY ELDERLY PATIENTS WITH LOWER URINARY TRACT SYMPTOMS]. <i>Japanese Journal of Urology</i> , 2019 , 110, 106-111		
17	Efficacy and safety of tadalafil 5mg once-daily in Asian men with both lower urinary tract symptoms associated with benign prostatic hyperplasia and erectile dysfunction: A phase III, randomized, double-blind, parallel, placebo- and tamsulosin-controlled study. <i>International Journal of Urology</i> , 2019 , 26, 192-200	2.3	11
16	Efficacy of newer medications for lower urinary tract symptoms attributed to benign prostatic hyperplasia: a systematic review. <i>Aging Male</i> , 2019 , 22, 1-11	2.1	21
15	Treatment of experimentally induced benign prostatic hyperplasia with Tadalafil and castration in dogs. <i>Theriogenology</i> , 2020 , 142, 236-245	2.8	4
14	Add-on effects of tadalafil in tamsulosin-treated patients with small benign prostatic enlargement: A randomized, placebo-controlled, double-blind, crossover study. <i>Neurourology and Urodynamics</i> , 2020 , 39, 237-242	2.3	2
13	Tadalafil Alone or in Combination with Tamsulosin for the Management for LUTS/BPH and ED. <i>Current Urology Reports</i> , 2020 , 21, 56	2.9	11
12	Tadalafil Improves Nocturia and Nocturia-Related Quality of Life in Patients with Benign Prostatic Hyperplasia (KYU-PRO Study). <i>Urologia Internationalis</i> , 2020 , 104, 587-593	1.9	4
11	Efficacy and safety of tadalafil vs tamsulosin in lower urinary tract symptoms (LUTS) as a result of benign prostate hyperplasia (BPH)-open label randomised controlled study. <i>International Journal of Clinical Practice</i> , 2020 , 74, e13530	2.9	3
10	Efficacy of testosterone replacement therapy plus alternate-day tadalafil for patients with late-onset hypogonadism: An open-label, randomized, crossover study. <i>International Journal of Urology</i> , 2021 , 28, 376-381	2.3	1
9	Sexual Dysfunctions Related to Drugs Used in the Management of Lower Urinary Tract Symptoms Due to Benign Prostatic Hyperplasia: A Narrative Review on α -Blockers and 5-Alpha Reductase Inhibitors. <i>Uro</i> , 2021 , 1, 82-98		
8	Are phosphodiesterase type 5 inhibitors effective for the management of lower urinary symptoms suggestive of benign prostatic hyperplasia?. <i>World Journal of Nephrology</i> , 2015 , 4, 138-47	3.6	9
7	Efficacy and Safety of 12-week Monotherapy With Once Daily 5 mg Tadalafil for Lower Urinary Tract Symptoms of Benign Prostatic Hyperplasia: Evidence-based Analysis. <i>Frontiers in Medicine</i> , 2021 , 8, 744012	4.9	0
6	New approaches to management of erectile dysfunction and lower urinary tract symptoms associated with benign prostatic hyperplasia. <i>Andrologia I Genitalsnaa Hirurgia</i> , 2019 , 20, 21-25	0.5	
5	Real-World Safety and Effectiveness of Tadalafil in Patients with Lower Urinary Tract Symptoms Secondary to Benign Prostatic Hyperplasia: A Japanese Post-Marketing Surveillance Study. <i>Journal of Pragmatic and Observational Research</i> , 2020 , 11, 45-54	7.4	2
4	Management for lower urinary tract dysfunction in the elderly according to guidelines. <i>Japanese Journal of Geriatrics</i> , 2022 , 59, 115-130	0	

- 3 Efficacy of Combination Treatment with Tadalafil and Mirabegron in Patients with Benign Prostatic Hyperplasia Who Presented with Persistent Storage Symptoms After Tadalafil Monotreatment: A Prospective, Multicenter, Open-Labeled Study. **2022**, 65, 231-237 ○
- 2 Discontinuation Rates of Tadalafil Alone and in Combination with α -Blockers in the Treatment of Male Lower Urinary Tract Symptoms with or without Coexisting Erectile Dysfunction: A Systematic Review and Meta-Analysis. **2022**, 2022, 1-14 ○
- 1 Effectiveness and Safety of a Mixture of Nobiletin and Tangeretin in Nocturia Patients: A Randomized, Placebo-Controlled, Double-Blind, Crossover Study. **2023**, 12, 2757 ○