

CITATION REPORT

List of articles citing

ENMeval: An R package for conducting spatially independent evaluations and estimating optimal model complexity for Maxent ecological niche models

DOI: 10.1111/2041-210x.12261

Methods in Ecology and Evolution, 2014, 5, 1198-1205.

Source: <https://exaly.com/paper-pdf/59919646/citation-report.pdf>

Version: 2024-04-27

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

| # | Paper | IF | Citations |
|------|--|-----|-----------|
| 1059 | Erratum. <i>Methods in Ecology and Evolution</i> , 2015 , 6, 119-120 | 7.7 | |
| 1058 | Lichen-Forming Fungi of the Genus <i>Montanelia</i> in Poland and Their Potential Distribution in Central Europe. 2015 , 28, 697-712 | | 4 |
| 1057 | Impact of model complexity on cross-temporal transferability in Maxent species distribution models: An assessment using paleobotanical data. 2015 , 312, 308-317 | | 95 |
| 1056 | Range increase of a Neotropical orchid bee under future scenarios of climate change. 2015 , 19, 901-910 | | 21 |
| 1055 | Fuzzy set theory for predicting the potential distribution and cost-effective monitoring of invasive species. 2015 , 316, 122-132 | | 11 |
| 1054 | Potential Distribution of Chagas Disease Vectors (Hemiptera, Reduviidae, Triatominae) in Colombia, Based on Ecological Niche Modeling. 2016 , 2016, 1439090 | | 22 |
| 1053 | Advances and Limitations of Disease Biogeography Using Ecological Niche Modeling. 2016 , 7, 1174 | | 67 |
| 1052 | Global Potential Distribution of <i>Bactrocera carambolae</i> and the Risks for Fruit Production in Brazil. 2016 , 11, e0166142 | | 13 |
| 1051 | Life History Traits and Niche Instability Impact Accuracy and Temporal Transferability for Historically Calibrated Distribution Models of North American Birds. 2016 , 11, e0151024 | | 24 |
| 1050 | Glacial refugia, recolonization patterns and diversification forces in Alpine-endemic <i>Megabunus</i> harvestmen. 2016 , 25, 2904-19 | | 27 |
| 1049 | Using spatiotemporal correlative niche models for evaluating the effects of climate change on mountain pine beetle. 2016 , 7, e01396 | | 13 |
| 1048 | Do community-weighted mean functional traits reflect optimal strategies?. 2016 , 283, 20152434 | | 81 |
| 1047 | Predicting the potential environmental suitability for <i>Theileria orientalis</i> transmission in New Zealand cattle using maximum entropy niche modelling. 2016 , 224, 82-91 | | 5 |
| 1046 | Summer-Habitat Suitability Modeling of <i>Myotis sodalis</i> (Indiana Bat) in the Eastern Mountains of West Virginia. 2016 , 23, 100-117 | | 4 |
| 1045 | Quantifying the impacts of sea-level rise on coastal biodiversity: A case study on lichens in the mid-Atlantic Coast of eastern North America. 2016 , 202, 119-126 | | 17 |
| 1044 | Landscape and flow metrics affecting the distribution of a federally-threatened fish: Improving management, model fit, and model transferability. 2016 , 342, 1-18 | | 14 |
| 1043 | Novel methods to select environmental variables in MaxEnt: A case study using invasive crayfish. 2016 , 341, 5-13 | | 67 |

| | | |
|------|--|-----|
| 1042 | Predicted Pleistocene-Holocene range shifts of the tiger (<i>Panthera tigris</i>). 2016 , 22, 1199-1211 | 21 |
| 1041 | Taxonomy and biogeography of <i>Bunopus spatulurus</i> (Reptilia; Gekkonidae) from the Arabian Peninsula. 2016 , 54, 67-81 | 17 |
| 1040 | Introducing BioSARN - an ecological niche model refinement tool. 2016 , 6, 5808-16 | 4 |
| 1039 | Do Ecological Niche Models Accurately Identify Climatic Determinants of Species Ranges?. 2016 , 187, 423-35 | 59 |
| 1038 | Realized climate niche breadth varies with population trend and distribution in North American birds. 2016 , 25, 1173-1180 | 11 |
| 1037 | Field validation of an invasive species Maxent model. 2016 , 36, 126-134 | 102 |
| 1036 | Predicting the distributions of predator (snow leopard) and prey (blue sheep) under climate change in the Himalaya. 2016 , 6, 4065-75 | 67 |
| 1035 | Regional modeling of large wildfires under current and potential future climates in Colorado and Wyoming, USA. 2016 , 134, 565-577 | 14 |
| 1034 | How important are choice of model selection method and spatial autocorrelation of presence data for distribution modelling by MaxEnt?. 2016 , 328, 108-118 | 44 |
| 1033 | Forest-structure data improve distribution models of threatened habitat specialists: Implications for conservation of epiphytic lichens in forest landscapes. 2016 , 196, 31-38 | 21 |
| 1032 | Modeling habitat distribution of <i>Cornus officinalis</i> with Maxent modeling and fuzzy logics in China. 2016 , 9, 742-751 | 32 |
| 1031 | Climate change impacts on endemic, high-elevation lichens in a biodiversity hotspot. 2016 , 25, 555-568 | 58 |
| 1030 | Distributional modeling of Mantophasmatodea (Insecta: Notoptera): a preliminary application and the need for future sampling. 2016 , 16, 259-268 | 3 |
| 1029 | Climatic Similarity of Extant and Extinct <i>Dasypus</i> Armadillos. 2017 , 24, 193-206 | 5 |
| 1028 | The role of abiotic and biotic factors in determining coexistence of multiple pollinators in the yucca-yucca moth mutualism. 2017 , 40, 511-520 | 4 |
| 1027 | Landslide susceptibility assessment using maximum entropy model with two different data sampling methods. 2017 , 152, 144-162 | 119 |
| 1026 | Palaeodistribution modelling of European vegetation types at the Last Glacial Maximum using modern analogues from Siberia: Prospects and limitations. 2017 , 159, 103-115 | 40 |
| 1025 | Lazarus ecology: Recovering the distribution and migratory patterns of the extinct Carolina parakeet. 2017 , 7, 5467-5475 | 11 |

| | | |
|------|--|-----|
| 1024 | On the need to consider multiphasic sensitivity of marine organisms to climate change: a case study of the Antarctic acorn barnacle. 2017 , 44, 2165-2175 | 10 |
| 1023 | Multilocus phylogeny of East African gerbils (Rodentia, Gerbilliscus) illuminates the history of the Somali-Masai savanna. 2017 , 44, 2295-2307 | 22 |
| 1022 | Reintroduced Eurasian beavers (<i>Castor fiber</i>): colonization and range expansion across human-dominated landscapes. 2017 , 26, 1863-1876 | 17 |
| 1021 | Hiding in a Cool Climatic Niche in the Tropics? An Assessment of the Ecological Biogeography of Hairy Long-Nosed Armadillos (<i>Dasypus pilosus</i>). 2017 , 10, 194008291769724 | 3 |
| 1020 | Opening the black box: an open-source release of Maxent. 2017 , 40, 887-893 | 762 |
| 1019 | Cryptic diversity in a fig wasp community-morphologically differentiated species are sympatric but cryptic species exhibit competitive exclusion. 2017 , 26, 937-950 | 24 |
| 1018 | The importance of herbivore density and management as determinants of the distribution of rare plant species. 2017 , 205, 77-84 | 10 |
| 1017 | Distribution and habitat associations of the critically endangered bird species of S Tom Island (Gulf of Guinea). 2017 , 27, 455-469 | 12 |
| 1016 | Spatial correspondence between the specialist galling insect genus <i>Kladothrips</i> (Thysanoptera: Tubulifera: Phlaeothripidae) and its <i>Acacia</i> host: ecological and evolutionary constraints on host use. 2017 , 56, 412-423 | |
| 1015 | The relative contributions of competition and abiotic tolerances in determining the geographical distributions of four closely related <i>Yucca</i> species in Texas. 2017 , 44, 1373-1382 | 3 |
| 1014 | Parasite biodiversity faces extinction and redistribution in a changing climate. 2017 , 3, e1602422 | 125 |
| 1013 | Idiosyncratic responses to climate-driven forest fragmentation and marine incursions in reed frogs from Central Africa and the Gulf of Guinea Islands. 2017 , 26, 5223-5244 | 34 |
| 1012 | Impact of climate and host availability on future distribution of Colorado potato beetle. 2017 , 7, 4489 | 20 |
| 1011 | Armadillo Mapper: A Case Study of an Online Application to Update Estimates of Species Potential Distributions. 2017 , 10, 194008291772413 | 1 |
| 1010 | Temperature drives abundance fluctuations, but spatial dynamics is constrained by landscape configuration: Implications for climate-driven range shift in a butterfly. 2017 , 86, 1339-1351 | 15 |
| 1009 | Can Niche Modeling and Geometric Morphometrics Document Competitive Exclusion in a Pair of Subterranean Rodents (Genus <i>Ctenomys</i>) with Tiny Parapatric Distributions?. 2017 , 7, 16283 | 15 |
| 1008 | The latitudinal diversity gradient in New World swallowtail butterflies is caused by contrasting patterns of out-of- and into-the-tropics dispersal. 2017 , 26, 1447-1458 | 14 |
| 1007 | Common garden test of range limits as predicted by a species distribution model in the annual plant. 2017 , 104, 817-827 | 7 |

| | | |
|------|--|-----|
| 1006 | Wild Felid Range Shift Due to Climatic Constraints in the Americas: a Bottleneck Explanation for Extinct Felids?. 2017 , 24, 427-438 | 7 |
| 1005 | A practical overview of transferability in species distribution modeling. 2017 , 25, 127-133 | 37 |
| 1004 | A single-algorithm ensemble approach to estimating suitability and uncertainty: cross-time projections for four Malagasy tenrecs. 2017 , 23, 196-208 | 10 |
| 1003 | Applying the One-Class Classification Method of Maxent to Detect an Invasive Plant <i>Spartina alterniflora</i> with Time-Series Analysis. 2017 , 9, 1120 | 14 |
| 1002 | Continental divide: Predicting climate-mediated fragmentation and biodiversity loss in the boreal forest. 2017 , 12, e0176706 | 18 |
| 1001 | Evidence of niche shift and invasion potential of <i>Lithobates catesbeianus</i> in the habitat of Mexican endemic frogs. 2017 , 12, e0185086 | 17 |
| 1000 | Using worldwide edaphic data to model plant species niches: An assessment at a continental extent. 2017 , 12, e0186025 | 46 |
| 999 | Forecasting distributions of an aquatic invasive species (<i>Nitellopsis obtusa</i>) under future climate scenarios. 2017 , 12, e0180930 | 20 |
| 998 | How many species of mammals are there in Brazil? New records of rare rodents (Rodentia: Cricetidae: Sigmodontinae) from Amazonia raise the current known diversity. 2017 , 5, e4071 | 12 |
| 997 | Identifying potential areas for an expanding wolf population in Sweden. 2018 , 220, 170-181 | 12 |
| 996 | Dryland photoautotrophic soil surface communities endangered by global change. 2018 , 11, 185-189 | 158 |
| 995 | Comparison of model selection technique performance in predicting the spread of newly invasive species: a case study with <i>Batrachochytrium salamandrivorans</i> . 2018 , 20, 2107-2119 | 18 |
| 994 | Host plant distributions and climate interact to affect the predicted geographic distribution of a Neotropical termite. 2018 , 50, 625-632 | 10 |
| 993 | Ignoring seasonal changes in the ecological niche of non-migratory species may lead to biases in potential distribution models: lessons from bats. 2018 , 27, 2425-2441 | 42 |
| 992 | Ecological niche modeling re-examined: A case study with the Darwin's fox. 2018 , 8, 4757-4770 | 29 |
| 991 | Revised distributional estimates for the recently discovered olinguito (<i>Bassaricyon neblina</i>), with comments on natural and taxonomic history. 2018 , 99, 321-332 | 11 |
| 990 | Potential global distribution of <i>Diabrotica</i> species and the risks for agricultural production. 2018 , 74, 2100 | 17 |
| 989 | Higher spring temperatures increase food scarcity and limit the current and future distributions of crossbills. 2018 , 24, 473-484 | 11 |

| | | | |
|-----|--|-----|----|
| 988 | Transferability of species distribution models for the detection of an invasive alien bryophyte using imaging spectroscopy data. 2018 , 68, 61-72 | | 11 |
| 987 | What are the best predictors for invasive potential of weeds? Transferability evaluations of model predictions based on diverse environmental data sets for <i>Flaveria bidentis</i> . 2018 , 58, 141-149 | | 13 |
| 986 | Wrong, but useful: regional species distribution models may not be improved by range-wide data under biased sampling. 2018 , 8, 2196-2206 | | 37 |
| 985 | Areas of endemism persist through time: A palaeoclimatic analysis in the Mexican Transition Zone. 2018 , 45, 952-961 | | 11 |
| 984 | Snub-nosed monkeys (): potential distribution and its implication for conservation. 2018 , 27, 1517-1538 | | 26 |
| 983 | Optimizing ensembles of small models for predicting the distribution of species with few occurrences. <i>Methods in Ecology and Evolution</i> , 2018 , 9, 802-808 | 7.7 | 59 |
| 982 | Species distribution models for Peruvian plantcutter improve with consideration of biotic interactions. 2018 , 49, jav-01617 | | 29 |
| 981 | Testing Range-Limit Hypotheses Using Range-Wide Habitat Suitability and Occupancy for the Scarlet Monkeyflower (<i>Erythranthe cardinalis</i>). 2018 , 191, E76-E89 | | 19 |
| 980 | Assessing the aggregated risk of invasive crayfish and climate change to freshwater crabs: A Southeast Asian case study. 2018 , 223, 58-67 | | 11 |
| 979 | Predicting the distribution of <i>Stipa purpurea</i> across the Tibetan Plateau via the MaxEnt model. 2018 , 18, 10 | | 59 |
| 978 | New insights into the distribution and conservation status of the Golden-White Tassel-Ear Marmoset <i>Mico chrysoleucos</i> (Primates, Callitrichidae). 2018 , 59, 347-353 | | 1 |
| 977 | Predicting the effects of future climate change on the distribution of an endemic damselfly (Odonata, Coenagrionidae) in subtropical South American grasslands. 2018 , 22, 303-319 | | 5 |
| 976 | Ecological niche modeling for a cultivated plant species: a case study on taro (<i>Colocasia esculenta</i>) in Hawaii. 2018 , 28, 967-977 | | 2 |
| 975 | A climate-associated multispecies cryptic cline in the northwest Atlantic. 2018 , 4, eaaq0929 | | 49 |
| 974 | Spiny mice of the Zambezian bioregion [phylogeny, biogeography and ecological differentiation within the <i>Acomys spinosissimus</i> complex. 2018 , 91, 79-90 | | 11 |
| 973 | The critical role of local refugia in postglacial colonization of Chinese pine: joint inferences from DNA analyses, pollen records, and species distribution modeling. 2018 , 41, 592-606 | | 13 |
| 972 | Maximum entropy modeling for orogenic gold prospectivity mapping in the Tangbale-Hatu belt, western Junggar, China. 2018 , 100, 133-147 | | 19 |
| 971 | Environmental marginality and geographic range limits: a case study with <i>Arabidopsis lyrata</i> ssp. <i>lyrata</i> . 2018 , 41, 622-634 | | 13 |

| | | | |
|-----|--|-----|-----|
| 970 | ENVIREM: an expanded set of bioclimatic and topographic variables increases flexibility and improves performance of ecological niche modeling. 2018 , 41, 291-307 | | 198 |
| 969 | Contribution of spatially explicit models to climate change adaptation and mitigation plans for a priority forest habitat. 2018 , 23, 371-386 | | 17 |
| 968 | The challenge of modeling niches and distributions for data-poor species: a comprehensive approach to model complexity. 2018 , 41, 726-736 | | 71 |
| 967 | Used-habitat calibration plots: a new procedure for validating species distribution, resource selection, and step-selection models. 2018 , 41, 737-752 | | 20 |
| 966 | Niche dynamics of two cryptic <i>Prosopis</i> invading South American drylands. 2018 , 20, 181-194 | | 6 |
| 965 | Phylogeography of a widespread sub-Saharan murid rodent <i>Aethomys chrysophilus</i> : the role of geographic barriers and paleoclimate in the Zambezian bioregion. 2018 , 82, 373-387 | | 17 |
| 964 | Landscape epidemiology in urban environments: The example of rodent-borne <i>Trypanosoma</i> in Niamey, Niger. 2018 , 63, 307-315 | | 3 |
| 963 | Ensemble species distribution modelling with transformed suitability values. 2018 , 100, 136-145 | | 32 |
| 962 | Wallace: A flexible platform for reproducible modeling of species niches and distributions built for community expansion. <i>Methods in Ecology and Evolution</i> , 2018 , 9, 1151-1156 | 7.7 | 85 |
| 961 | Modeling relative habitat suitability of southern Florida for invasive Burmese pythons (<i>Python molurus bivittatus</i>). 2018 , 33, 257-274 | | 11 |
| 960 | Toward ecologically realistic predictions of species distributions: A cross-time example from tropical montane cloud forests. 2018 , 24, 1511-1522 | | 74 |
| 959 | Paintings predict the distribution of species, or the challenge of selecting environmental predictors and evaluation statistics. 2018 , 27, 245-256 | | 198 |
| 958 | Isolation by distance, not incipient ecological speciation, explains genetic differentiation in an Andean songbird (Aves: Furnariidae: <i>Cranioleuca antisensis</i> , Line-cheeked Spinetail) despite near threefold body size change across an environmental gradient. 2018 , 27, 279-296 | | 20 |
| 957 | Quantifying the climatic niche of symbiont partners in a lichen symbiosis indicates mutualist-mediated niche expansions. 2018 , 41, 1380-1392 | | 28 |
| 956 | A MaxEnt Model for Mineral Prospectivity Mapping. 2018 , 27, 299-313 | | 17 |
| 955 | Climatic Niche Model for Overwintering Monarch Butterflies in a Topographically Complex Region of California. 2018 , 9, | | 5 |
| 954 | Mapping Fishing Activities and Suitable Fishing Grounds Using Nighttime Satellite Images and Maximum Entropy Modelling. 2018 , 10, 1604 | | 24 |
| 953 | Resolving distribution and population fragmentation in two leaf-tailed gecko species of north-east Australia: key steps in the conservation of microendemic species. 2018 , 66, 152 | | 3 |

| | | |
|-----|--|----|
| 952 | Glacial Refugia and the Postglacial Migration of Dominant Tree Species in Northern China. 2018 , 31-56 | |
| 951 | Direct modelling of limited migration improves projected distributions of Himalayan amphibians under climate change. 2018 , 227, 352-360 | 28 |
| 950 | Towards integrative taxonomy in Neotropical botany: disentangling the <i>Pagamea guianensis</i> species complex (Rubiaceae). 2018 , 188, 213-231 | 25 |
| 949 | Improved species-occurrence predictions in data-poor regions: using large-scale data and bias correction with down-weighted Poisson regression and Maxent. 2018 , 41, 1161-1172 | 36 |
| 948 | Genomic data reject the hypothesis of sympatric ecological speciation in a clade of <i>Desmognathus</i> salamanders. 2018 , 72, 2378-2393 | 9 |
| 947 | Aerial surveys and distribution models enable monitoring of fishing in Marine Protected Areas. 2018 , 165, 298-306 | 5 |
| 946 | Range dynamics of Palaearctic steppe species under glacial cycles: the phylogeography of <i>Proterebia afra</i> (Lepidoptera: Nymphalidae: Satyrinae). 2018 , | 2 |
| 945 | Using insects to detect, monitor and predict the distribution of <i>Xylella fastidiosa</i> : a case study in Corsica. 2018 , 8, 15628 | 38 |
| 944 | Using species distribution modelling to determine opportunities for trophic rewilding under future scenarios of climate change. 2018 , 373, | 25 |
| 943 | Predicting the potential distribution of the endangered red panda across its entire range using MaxEnt modeling. 2018 , 8, 10542-10554 | 51 |
| 942 | New State Record and Potential Distribution of the Snake <i>Sibon nebulatus</i> (Dipsadidae) from Mexico. 2018 , 78, 242-246 | |
| 941 | The effect of large sample sizes on ecological niche models: Analysis using a North American rodent, <i>Peromyscus maniculatus</i> . 2018 , 386, 83-88 | 12 |
| 940 | Modeling the potential habitats of dusky, commons and bottlenose dolphins in the Humboldt Current System off Peru: The influence of non-El Niño vs. El Niño 1997-98 conditions and potential prey availability. 2018 , 168, 169-181 | 2 |
| 939 | Inferring space from time: On the relationship between demography and environmental suitability in the desert plant <i>O. rastrera</i> . 2018 , 13, e0201543 | 2 |
| 938 | Climatic niche and potential distribution of <i>Tithonia diversifolia</i> (Hemsl.) A. Gray in Africa. 2018 , 13, e0202421 | 12 |
| 937 | First genome-wide analysis of the endangered, endemic lichen <i>Cetradonia linearis</i> reveals isolation by distance and strong population structure. 2018 , 105, 1556-1567 | 23 |
| 936 | Estimating the population size of lemurs based on their mutualistic food trees. 2018 , 45, 2546-2563 | 7 |
| 935 | Random subset feature selection for ecological niche models of wildfire activity in Western North America. 2018 , 383, 52-68 | 11 |

| | | |
|-----|--|----|
| 934 | A Framework for Simultaneous Tests of Abiotic, Biotic, and Historical Drivers of Species Distributions: Empirical Tests for North American Wood Warblers Based on Climate and Pollen. 2018 , 192, E48-E61 | 10 |
| 933 | Niche conservatism and phylogenetic clustering in a tribe of arid-adapted marsupial mice, the Sminthopsini. 2018 , 31, 1204-1215 | 4 |
| 932 | Isolation by marine barriers and climate explain areas of endemism in an island rodent. 2018 , 45, 2053-2066 | 6 |
| 931 | Niche models inform the effects of climate change on the endangered Nilgiri Tahr (<i>Nilgiritragus hylocrius</i>) populations in the southern Western Ghats, India. 2018 , 120, 355-363 | 25 |
| 930 | Predicting spatial factors associated with cattle depredations by the Mexican wolf (<i>Canis lupus baileyi</i>) with recommendations for depredation risk modeling. 2018 , 224, 327-335 | 8 |
| 929 | Predicting environmental suitability for key benthic species in an ecologically and economically important deep-sea environment. 2018 , 157-158, 121-133 | 7 |
| 928 | Relative importance of environmental variables for the distribution of the invasive marsh species <i>Spartina alterniflora</i> across different spatial scales. 2018 , 69, 790 | 5 |
| 927 | Land Cover and Climate Change May Limit Invasiveness of in Wales. 2018 , 9, 664 | 18 |
| 926 | Invasion of a Legume Ecosystem Engineer in a Cold Biome Alters Plant Biodiversity. 2018 , 9, 715 | 11 |
| 925 | Systematics of the broad-nosed bats, <i>Platyrrhinus umbratus</i> (Lyon, 1902) and <i>P. nigellus</i> (Gardner and Carter, 1972) (Chiroptera: Phyllostomidae), based on genetic, morphometric, and ecological niche analyses. 2018 , 4, 119-133 | 4 |
| 924 | No deaths in the desert: predicted responses of an arid-adapted bee and its two nesting trees suggest resilience in the face of warming climates. 2018 , 11, 449-463 | 6 |
| 923 | Distribution of the meadow lizard in Europe and its realized ecological niche model. 2018 , 52, 1909-1925 | 6 |
| 922 | Kelp Long-Distance Dispersal: Role of Ecological/Oceanographic Processes and Implications to Marine Forest Conservation. 2018 , 10, 11 | 14 |
| 921 | Mapping Disease Transmission Risk of Nipah Virus in South and Southeast Asia. 2018 , 3, | 18 |
| 920 | Multi-criteria spatial identification of carnivore conservation areas under data scarcity and conflict: a jaguar case study in Sierra Nevada de Santa Marta, Colombia. 2018 , 27, 3373-3392 | 4 |
| 919 | Biotic interactions in species distribution models enhance model performance and shed light on natural history of rare birds: a case study using the straight-billed reedhaunter <i>Limnortyx rectirostris</i> . 2018 , 49, e01743 | 17 |
| 918 | Phylogeography of an endangered disjunct herb: long-distance dispersal, refugia and colonization routes. 2018 , 10, ply047 | 7 |
| 917 | Enhancing conservation network design with graph-theory and a measure of protected area effectiveness: Refining wildlife corridors in Belize, Central America. 2018 , 178, 51-59 | 23 |

| | | |
|-----|--|-----|
| 916 | Cryptic genetic diversity, population structure, and gene flow in the Mojave rattlesnake (<i>Crotalus scutulatus</i>). 2018 , 127, 669-681 | 19 |
| 915 | Major challenges for correlational ecological niche model projections to future climate conditions. 2018 , 1429, 66-77 | 55 |
| 914 | Characterizing the cultural evolutionary process from eco-cultural niche models: niche construction during the Neolithic of the Struma River Valley (c. 6200-900 BC). 2019 , 11, 2181-2200 | 6 |
| 913 | Natural re-colonization and admixture of wolves (<i>Canis lupus</i>) in the US Pacific Northwest: challenges for the protection and management of rare and endangered taxa. 2019 , 122, 133-149 | 7 |
| 912 | An evaluation of transferability of ecological niche models. 2019 , 42, 521-534 | 41 |
| 911 | Spatial variability in species' potential distributions during the Last Glacial Maximum under different Global Circulation Models: Relevance in evolutionary biology. 2019 , 57, 113-126 | 9 |
| 910 | Predicted effects of climate factors on mountain species are not uniform over different spatial scales. 2019 , 50, | 7 |
| 909 | Cryptic phylogeographic history sheds light on the generation of species diversity in sky-island mountains. 2019 , 46, 2232-2247 | 6 |
| 908 | Ensemble Modeling of Antarctic Macroalgal Habitats Exposed to Glacial Melt in a Polar Fjord. 2019 , 7, | 14 |
| 907 | Islands in the desert: environmental distribution modelling of endemic flora reveals the extent of Pleistocene tropical relict vegetation in southern Arabia. 2019 , 124, 411-422 | 5 |
| 906 | An exhaustive analysis of heuristic methods for variable selection in ecological niche modeling and species distribution modeling. 2019 , 53, 100983 | 32 |
| 905 | Back home? Uncertainties for returning seized animals to the source-areas under climate change. 2019 , 25, 3242-3253 | 4 |
| 904 | kuenm: an R package for detailed development of ecological niche models using Maxent. 2019 , 7, e6281 | 185 |
| 903 | Plant phylogenetic diversity stabilizes large-scale ecosystem productivity. 2019 , 28, 1430-1439 | 11 |
| 902 | Current and future climatic regions favourable for a globally introduced wild carnivore, the raccoon <i>Procyon lotor</i> . 2019 , 9, 9174 | 16 |
| 901 | Phylogeography of the Tibetan hamster in response to uplift and environmental change in the Qinghai-Tibet Plateau. 2019 , 9, 7291-7306 | 6 |
| 900 | Effects of climate change on the distributional potential of three range-restricted West African bird species. 2019 , 121, | 6 |
| 899 | Recalculating route: dispersal constraints will drive the redistribution of Amazon primates in the Anthropocene. 2019 , 42, 1789-1801 | 27 |

| | | |
|-----|--|----|
| 898 | Spatial distribution and spread potential of sixteen <i>Leptospira</i> serovars in a subtropical region of Brazil. 2019 , 66, 2482-2495 | 8 |
| 897 | Exploring the Interplay Between Local and Regional Drivers of Distribution of a Subterranean Organism. 2019 , 11, 119 | 3 |
| 896 | Mapping parasite transmission risk from white-tailed deer to a declining moose population. 2019 , 65, 1 | 6 |
| 895 | Future distribution of wild boar in a highly anthropogenic landscape: Models combining hunting bag and citizen science data. 2019 , 411, 108804 | 10 |
| 894 | Spatio-Temporal Distribution of Monarch Butterflies Along Their Migratory Route. 2019 , 7, | 2 |
| 893 | The MIAMaxent R package: Variable transformation and model selection for species distribution models. 2019 , 9, 12051-12068 | 15 |
| 892 | Evaluating the boundaries of marine biogeographic regions of the Southwestern Atlantic using halacarid mites (Halacaridae), meiobenthic organisms with a low dispersal potential. 2019 , 9, 13359-13374 | 2 |
| 891 | Impact of Habitat Loss and Mining on the Distribution of Endemic Species of Amphibians and Reptiles in Mexico. 2019 , 11, 210 | 6 |
| 890 | Collinearity in ecological niche modeling: Confusions and challenges. 2019 , 9, 10365-10376 | 76 |
| 889 | Marine protected areas modulate habitat suitability of the invasive round goby (<i>Neogobius melanostomus</i>) in the Baltic Sea. 2019 , 229, 106380 | 1 |
| 888 | Predicting the risk of illegal activity and evaluating law enforcement interventions in the western Serengeti. 2019 , 1, e81 | 3 |
| 887 | Known and estimated distribution in Mexico of <i>Batrachochytrium dendrobatidis</i> , a pathogenic fungus of amphibians. 2019 , 51, 731-746 | 8 |
| 886 | Testing the role of ecological selection on colour pattern variation in the butterfly <i>Parnassius clodius</i> . 2019 , 28, 5086-5102 | 2 |
| 885 | Impacts of 21st-century climate change on montane habitat in the Madiran Sky Island Archipelago. 2019 , 25, 1625-1638 | 8 |
| 884 | Tracking seasonal activity of the western blacklegged tick across California. 2019 , 56, 2562-2573 | 8 |
| 883 | Species' range model metadata standards: RMMS. 2019 , 28, 1912-1924 | 8 |
| 882 | Climate change and its potential impact on the conservation of the Hoary Fox, <i>Lycalopex vetulus</i> (Mammalia: Canidae). 2019 , 98, 91-101 | 3 |
| 881 | Predicting Optimal Sites for Ecosystem Restoration Using Stacked-Species Distribution Modeling. 2019 , 6, | 17 |

| | | |
|-----|---|----|
| 880 | Landscape Features and Climatic Forces Shape the Genetic Structure and Evolutionary History of an Oak Species () in East China. 2019 , 10, 1060 | 14 |
| 879 | Modeling the Potential Global Distribution of <i>Phenacoccus madeirensis</i> Green under Various Climate Change Scenarios. 2019 , 10, 773 | 11 |
| 878 | Expansion of the agricultural frontier in the largest South American Dry Forest: Identifying priority conservation areas for snakes before everything is lost. 2019 , 14, e0221901 | 12 |
| 877 | Where will it cross next? Optimal management of road collision risk for otters in Italy. 2019 , 251, 109609 | 8 |
| 876 | State of knowledge and potential distribution of the Colombian endemic brown hairy dwarf porcupine <i>Coendou vestitus</i> (Rodentia). 2019 , 99, 1-11 | 3 |
| 875 | A checklist for maximizing reproducibility of ecological niche models. 2019 , 3, 1382-1395 | 56 |
| 874 | Ecological Niche Modeling of Three Species of <i>Stenella</i> Dolphins in the Caribbean Basin, With Application to the Seaflower Biosphere Reserve. 2019 , 6, | 18 |
| 873 | Polyploidy in creosote bush () shapes the biogeography of specialist herbivores. 2019 , 46, 597-610 | 4 |
| 872 | Exploring potential establishment of marine rafting species after transoceanic long-distance dispersal. 2019 , 28, 588-600 | 3 |
| 871 | Potential migratory routes of <i>Urania boisduvalii</i> (Lepidoptera: Uraniidae) among host plant populations. 2019 , 25, 478-488 | 2 |
| 870 | Current and future ranges of an elusive North American insect using species distribution models. 2019 , 23, 175-186 | 5 |
| 869 | Ecological and phenotypic divergence in Iberian shrews (Soricidae). 2019 , 57, 642 | 1 |
| 868 | Mapping changes in the spatiotemporal distribution of lumpy skin disease virus. 2019 , 66, 2045-2057 | 11 |
| 867 | Associations between habitat quality, body size and reproductive fitness in the alpine endemic spider <i>Vesubia jugorum</i> . 2019 , 28, 1325-1335 | 18 |
| 866 | Scots pine (<i>Pinus sylvestris</i> L.) mortality is explained by the climatic suitability of both host tree and bark beetle populations. 2019 , 448, 119-129 | 19 |
| 865 | Scale-dependent estimates of niche overlap and environmental effects on two sister species of Neotropical snakes. 2019 , 54, 121-132 | 3 |
| 864 | Genes on the edge: A framework to detect genetic diversity imperiled by climate change. 2019 , 25, 4034-4047 | 12 |
| 863 | Allopatric divergence and secondary contact with gene flow: a recurring theme in rattlesnake speciation. 2019 , 128, 149-169 | 10 |

- 862 Characterization and Mapping of a Deep-Sea Sponge Ground on the Tropic Seamount (Northeast Tropical Atlantic): Implications for Spatial Management in the High Seas. **2019**, 6, 18
- 861 Climatic conditions and human activities shape diploid-tetraploid coexistence at different spatial scales in the common weed *Tripleurospermum inodorum* (Asteraceae). **2019**, 46, 1355 4
- 860 Species Distribution Modeling in Latin America: A 25-Year Retrospective Review. **2019**, 12, 194008291985405 8
- 859 Spatiotemporal dynamics in habitat suitability of a large Arctic herbivore: Environmental heterogeneity is key to a sedentary lifestyle. **2019**, 18, e00647 13
- 858 The importance of biological plausibility for data poor models in the face of an immediate threat by an emerging infectious disease: a reply to Katz and Zellmer (2018). **2019**, 21, 2789-2793 3
- 857 Spatial ecology and conservation of the microendemic ovenbird Cipo *Cinclodes* (*Cinclodes espinhacensis*) from the Brazilian highlands. **2019**, 90, 128-142 6
- 856 Maximizing habitat connectivity in the mitigation hierarchy. A case study on three terrestrial mammals in an urban environment. **2019**, 243, 340-349 18
- 855 Modeling algorithm influence on the success of predicting new populations of rare species: ground-truthing models for the Pale-Belly Frost Lichen (*Physconia subpallida*) in Ontario. **2019**, 28, 1853-1862 3
- 854 Modeling the potential range expansion of larger grain borer, *Prostephanus truncatus* (Coleoptera: Bostrichidae). **2019**, 9, 6862 19
- 853 A new null model approach to quantify performance and significance for ecological niche models of species distributions. **2019**, 46, 1101-1111 18
- 852 Pure, shared, and coupling effects of climate change and sea level rise on the future distribution of along the Chinese coast. **2019**, 9, 5380-5391 1
- 851 Broad-scale species distribution models applied to data-poor areas. **2019**, 175, 198-207 10
- 850 Duck production systems and highly pathogenic avian influenza H5N8 in France, 2016-2017. **2019**, 9, 6177 19
- 849 Evaluating current and future range limits of an endangered, keystone rodent (*Dipodomys ingens*). **2019**, 25, 1074-1087 6
- 848 Differences between ecological niches in northern and southern populations of Angolan black and white colobus monkeys (*Colobus angolensis palliatus* and *Colobus angolensis sharpei*) throughout Kenya and Tanzania. **2019**, 81, e22975 4
- 847 Distribution models of estuarine fish species: The effect of sampling bias, species ecology and threshold selection on models' accuracy. **2019**, 51, 168-176 5
- 846 How to survive a glaciation: the challenge of estimating biologically realistic potential distributions under freezing conditions. **2019**, 42, 1237-1245 5
- 845 Cross-continental phylogeography of two Holarctic Nymphalid butterflies, *Boloria eunomia* and *Boloria selene*. **2019**, 14, e0214483 13

| | | |
|-----|--|----|
| 844 | BioModelos: A collaborative online system to map species distributions. 2019 , 14, e0214522 | 9 |
| 843 | Range expansion of an already widespread bee under climate change. 2019 , 17, e00584 | 8 |
| 842 | The role of protected areas in preserving habitat and functional connectivity for mobile flying vertebrates: the common noctule bat (<i>Nyctalus noctula</i>) in Tuscany (Italy) as a case study. 2019 , 28, 1569-1592 ¹ | 1 |
| 841 | Spatiotemporally explicit demographic modelling supports a joint effect of historical barriers to dispersal and contemporary landscape composition on structuring genomic variation in a red-listed grasshopper. 2019 , 28, 2155-2172 | 9 |
| 840 | Environmental impact assessment of development projects improved by merging species distribution and habitat connectivity modelling. 2019 , 241, 439-449 | 20 |
| 839 | Niche conservatism predominates in adaptive radiation: comparing the diversification of Hawaiian arthropods using ecological niche modelling. 2019 , 127, 479-492 | 5 |
| 838 | Tabula rasa in the Patagonian Channels II: The cushion peat bog species <i>Astelia pumila</i> (Asteliaceae) and <i>Donatia fascicularis</i> (Stylidiaceae) survived the last glacial in south-central Chile. 2019 , 46, 899-914 | 2 |
| 837 | Assessing effects of genetic, environmental, and biotic gradients in species distribution modelling. 2019 , 76, 1762-1775 | 5 |
| 836 | Identifying cloud forest conservation areas in Mexico from the potential distribution of 19 representative species. 2019 , 5, e01423 | 2 |
| 835 | Diverse Mycorrhizal Associations Enhance Terrestrial C Storage in a Global Model. 2019 , 33, 501-523 | 51 |
| 834 | Integrating climate and land-use change scenarios in modelling the future spread of invasive squirrels in Italy. 2019 , 25, 644-659 | 38 |
| 833 | Shifts in Climatic Niche Occupation in <i>Astrophytum Coahuilense</i> (H. Müller) Kayser and Its Potential Distribution in Mexico. 2019 , 11, 1138 | 3 |
| 832 | Predicting range shifts of Asian elephants under global change. 2019 , 25, 822-838 | 28 |
| 831 | Retrospective and Predictive Investigation of Fish Kill Events. 2019 , 31, 61-70 | 1 |
| 830 | Akaike information criterion should not be a best of geographical prediction accuracy in ecological niche modelling. 2019 , 51, 25-32 | 32 |
| 829 | Similar but different: Revealing the relative roles of species-traits versus biome properties structuring genetic variation in South American marsh rats. 2019 , 46, 770-783 | 5 |
| 828 | Modeling fall migration pathways and spatially identifying potential migratory hazards for the eastern monarch butterfly. 2019 , 34, 443-458 | 16 |
| 827 | Potential risk of <i>Batrachochytrium</i> salamandrivorans in Mexico. 2019 , 14, e0211960 | 8 |

| | | |
|-----|--|--------|
| 826 | Potential Distribution of Six North American Higher-Attine Fungus-Farming Ant (Hymenoptera: Formicidae) Species. 2019 , 19, | 5 |
| 825 | Stability and changes in the distribution of Pipiza hoverflies (Diptera, Syrphidae) in Europe under projected future climate conditions. 2019 , 14, e0221934 | 4 |
| 824 | Strategies to optimize modeling habitat suitability of in the Pan-Amazonia. 2019 , 9, 12623-12638 | 7 |
| 823 | Biodiversity can benefit from climate stabilization despite adverse side effects of land-based mitigation. 2019 , 10, 5240 | 24 |
| 822 | The effect of local species composition on the distribution of an avian invader. 2019 , 9, 15861 | 5 |
| 821 | Links between prey assemblages and poison frog toxins: A landscape ecology approach to assess how biotic interactions affect species phenotypes. 2019 , 9, 14317-14329 | 6 |
| 820 | Sufficient versus optimal climatic stability during the Late Quaternary: using environmental quality to guide phylogeographic inferences in a Neotropical montane system. 2019 , 100, 1783-1807 | 5 |
| 819 | Disentangling the genetic effects of refugial isolation and range expansion in a trans-continently distributed species. 2019 , 122, 441-457 | 6 |
| 818 | Predicting Yellow Fever Through Species Distribution Modeling of Virus, Vector, and Monkeys. 2019 , 16, 95-108 | 26 |
| 817 | Modeling endangered mammal species distributions and forest connectivity across the humid Upper Guinea lowland rainforest of West Africa. 2019 , 28, 671-685 | 12 |
| 816 | Global invasion in progress: modeling the past, current and potential global distribution of the common myna. 2019 , 21, 1295-1309 | 18 |
| 815 | Searching for a needle in a haystack: Where to survey for climatically-matched biological control agents for two grasses (<i>Sporobolus</i> spp.) invading Australia. 2019 , 129, 37-44 | 7 |
| 814 | Use of geographic information system tools to predict animal breed suitability for different agro-ecological zones. 2019 , 13, 1536-1543 | 8 |
| 813 | The contribution of fog to water and nutrient supply to <i>Arthraerua leubnitziae</i> in the central Namib Desert, Namibia. 2019 , 161, 35-46 | 14 |
| 812 | Environmental filtering improves ecological niche models across multiple scales. <i>Methods in Ecology and Evolution</i> , 2019 , 10, 481-492 | 7.7 18 |
| 811 | Identifying potential areas of expansion for the endangered brown bear (<i>Ursus arctos</i>) population in the Cantabrian Mountains (NW Spain). 2019 , 14, e0209972 | 50 |
| 810 | Climate and land use changes will degrade the distribution of <i>Rhododendrons</i> in China. 2019 , 659, 515-528 | 27 |
| 809 | Current and Future Distribution of the Lone Star Tick, <i>Amblyomma americanum</i> (L.) (Acari: Ixodidae) in North America. 2019 , 14, e0209082 | 77 |

| | | | |
|-----|---|-----|-----|
| 808 | Phylogeography of Atlantic Forest glassfrogs (<i>Vitreorana</i>): when geography, climate dynamics and rivers matter. 2019 , 122, 545-557 | | 17 |
| 807 | Quaternary range-shifts of arboreal rodents of the genus <i>Habromys</i> (Cricetidae, Neotominae) in Mesoamerica and their evolutionary consequences. 2019 , 94, 4-10 | | 1 |
| 806 | blockCV: An r package for generating spatially or environmentally separated folds for k-fold cross-validation of species distribution models. <i>Methods in Ecology and Evolution</i> , 2019 , 10, 225-232 | 7.7 | 125 |
| 805 | Evolution of the Grey-bellied pygmy mouse group: Highly structured molecular diversity with predictable geographic ranges but morphological crypsis. 2019 , 130, 143-155 | | 12 |
| 804 | Conventional methods for enhancing connectivity in conservation planning do not always maintain gene flow. 2019 , 56, 913-922 | | 6 |
| 803 | Modelling climate change effects on Zagros forests in Iran using individual and ensemble forecasting approaches. 2019 , 137, 1015-1025 | | 9 |
| 802 | Noteworthy records and ecological niche modeling of the rare and endangered Goldman's diminutive woodrat <i>Nelsonia goldmani</i> (Rodentia: Cricetidae) endemic to central Mexican highlands. 2019 , 83, 330-342 | | 4 |
| 801 | Combined Effects of Global Climate Suitability and Regional Environmental Variables on the Distribution of an Invasive Marsh Species <i>Spartina alterniflora</i> . 2019 , 42, 99-111 | | 9 |
| 800 | Climatic niche of the Saker Falcon <i>Falco cherrug</i> : predicted new areas to direct population surveys in Central Asia. 2020 , 162, 27-41 | | 8 |
| 799 | The importance of soils in predicting the future of plant habitat suitability in a tropical forest. 2020 , 450, 151-170 | | 19 |
| 798 | Spatial priorities for agricultural development in the Brazilian Cerrado: may economy and conservation coexist?. 2020 , 29, 1683-1700 | | 9 |
| 797 | Current and future distribution of the invasive oak processionary moth. 2020 , 22, 523-534 | | 11 |
| 796 | Range and conservation updates for the Critically Endangered blue-eyed black lemur <i>Eulemur flavifrons</i> and the Vulnerable black lemur <i>Eulemur macaco</i> . 2020 , 54, 819-827 | | 2 |
| 795 | Predicting the current and future distribution of three <i>Coptis</i> herbs in China under climate change conditions, using the MaxEnt model and chemical analysis. 2020 , 698, 134141 | | 52 |
| 794 | Evaluating presence-only species distribution models with discrimination accuracy is uninformative for many applications. 2020 , 47, 167-180 | | 35 |
| 793 | Potential distribution of two invasive pineapple pests under climate change. 2020 , 76, 1652-1663 | | 9 |
| 792 | Assessing the status of critically endangered Kondana soft-furred rat (<i>Millardia kondana</i>) using integrative taxonomy: combining evidence from morphological, molecular and environmental niche modeling. 2020 , 84, 392-406 | | |
| 791 | Spatial Risk Distribution of Dengue Based on the Ecological Niche Model of <i>Aedes aegypti</i> (Diptera: Culicidae) in the Central Mexican Highlands. 2020 , 57, 728-737 | | 3 |

| | | |
|-----|--|----|
| 790 | Defining priorities for global snow leopard conservation landscapes. 2020 , 241, 108387 | 20 |
| 789 | Insight into intraspecific niche divergence and conservatism in American horseshoe crabs (<i>Limulus polyphemus</i>). 2020 , 22, e00896 | 2 |
| 788 | Ecological niche models and species distribution models in marine environments: A literature review and spatial analysis of evidence. 2020 , 415, 108837 | 79 |
| 787 | Niches and radiations: a case study on the Andean sapphire-vented puffleg <i>Eriocnemis luciani</i> and coppery-naped puffleg <i>E. sapphiropygia</i> (Aves, Trochilidae). 2020 , 51, | |
| 786 | ENMTML: An R package for a straightforward construction of complex ecological niche models. 2020 , 125, 104615 | 38 |
| 785 | Species interactions and climate change: How the disruption of species co-occurrence will impact on an avian forest guild. 2020 , 26, 1212-1224 | 17 |
| 784 | Biotic predictors with phenological information improve range estimates for migrating monarch butterflies in Mexico. 2020 , 43, 341-352 | 16 |
| 783 | Climate change and the future restructuring of Neotropical anuran biodiversity. 2020 , 43, 222-235 | 15 |
| 782 | Colonizing the east and the west: distribution and niche properties of a dwarf Asian honey bee invading Africa, the Middle East, the Malay Peninsula, and Taiwan. 2020 , 51, 75-87 | 6 |
| 781 | Environmental ranges estimated from species distribution models are not good predictors of lizard and frog physiological tolerances. 2020 , 34, 89-99 | 2 |
| 780 | Coexistence barriers confine the poleward range of a globally distributed plant. 2020 , 23, 1838-1848 | 5 |
| 779 | Are population isolations and declines a threat to island endemic water striders? A lesson from demographic and niche modelling of <i>Metrocoris esakii</i> (Hemiptera: Gerridae). 2020 , 29, 4573-4587 | 0 |
| 778 | Species Distribution Modeling Predicts Significant Declines in Coralline Algae Populations Under Projected Climate Change With Implications for Conservation Policy. 2020 , 7, | 4 |
| 777 | Assessing geographic and climatic variables to predict the potential distribution of the visceral leishmaniasis vector <i>Lutzomyia longipalpis</i> in the state of Espírito Santo, Brazil. 2020 , 15, e0238198 | 0 |
| 776 | Habitat suitability for primate conservation in north-east Brazil. 2020 , 54, 803-813 | 6 |
| 775 | Biotic interactions govern the distribution of coexisting ungulates in the Arctic Archipelago [A case for conservation planning. 2020 , 24, e01239 | 4 |
| 774 | Predicting range expansion of invasive species: Pitfalls and best practices for obtaining biologically realistic projections. 2020 , 26, 1767-1779 | 6 |
| 773 | : An R package to tune and evaluate species distribution models. 2020 , 10, 11488-11506 | 24 |

| | | | |
|-----|--|-----|----|
| 772 | Habitat suitability model of endangered and the probable consequences of global warming. 2020 , 61, 1-13 | | 6 |
| 771 | Modelling the geographical distributions of one native and two introduced species of crayfish in the French Alps. 2020 , 60, 101172 | | 3 |
| 770 | Identifying suitable reintroduction sites for the White-rumped Vulture (<i>Gyps bengalensis</i>) in India's Western Ghats using niche models and habitat requirements. 2020 , 158, 106034 | | 3 |
| 769 | Climate suitability as indicative of invasion potential for the most seized bird species in Brazil. 2020 , 58, 125890 | | 0 |
| 768 | Extrapolation in species distribution modelling. Application to Southern Ocean marine species. 2020 , 188, 102438 | | 6 |
| 767 | Projected climate and land use change alter western blacklegged tick phenology, seasonal host-seeking suitability and human encounter risk in California. 2020 , 26, 5459-5474 | | 12 |
| 766 | ntbox: An r package with graphical user interface for modelling and evaluating multidimensional ecological niches. <i>Methods in Ecology and Evolution</i> , 2020 , 11, 1199-1206 | 7.7 | 59 |
| 765 | Incorporating interspecific interactions into phylogeographic models: A case study with Californian oaks. 2020 , 29, 4510-4524 | | 6 |
| 764 | Comparison of Habitat Suitability Models for Neumann in North America to Determine Its Potential Geographic Range. 2020 , 17, | | 3 |
| 763 | Bioclimatic Modelling Identifies Suitable Habitat for the Establishment of the Invasive European Paper Wasp (Hymenoptera: Vespidae) across the Southern Hemisphere. 2020 , 11, | | 2 |
| 762 | Phylogenetic conservatism of abiotic niche in sympatric Southwestern Atlantic skates. 2020 , 16, 458-473 | | 1 |
| 761 | Toward reliable habitat suitability and accessibility models in an era of multiple environmental stressors. 2020 , 10, 10937-10952 | | 7 |
| 760 | The fate of endemic birds of eastern Brazilian mountaintops in the face of climate change. 2020 , 18, 257-266 | | 0 |
| 759 | Assessing Niche Shifts and Conservatism by Comparing the Native and Post-Invasion Niches of Major Forest Invasive Species. 2020 , 11, | | 2 |
| 758 | The roles of climate, geography and natural selection as drivers of genetic and phenotypic differentiation in a widespread amphibian <i>Hyla annectans</i> (Anura: Hylidae). 2020 , 29, 3667-3683 | | 4 |
| 757 | Pre-colonial Amerindian legacies in forest composition of southern Brazil. 2020 , 15, e0235819 | | 5 |
| 756 | Conservation planning for adaptive and neutral evolutionary processes. 2020 , 57, 2159-2169 | | 6 |
| 755 | Phylogenomic approaches reveal how climate shapes patterns of genetic diversity in an African rain forest tree species. 2020 , 29, 3560-3573 | | 4 |

| | | |
|-----|--|----|
| 754 | Influence of climate change on the predicted distributions of the genus <i>Tympanoctomys</i> (Rodentia, Hystricomorpha, Octodontidae), and their conservation implications. 2020 , 101, 1364-1379 | 1 |
| 753 | Integrative taxonomy identifies a new stingray species of the genus <i>Hypanus</i> Rafinesque, 1818 (<i>Dasyatidae</i> , <i>Myliobatiformes</i>), from the Tropical Southwestern Atlantic. 2020 , 97, 1120-1142 | 4 |
| 752 | Species Distribution Models and Niche Partitioning among Unisexual <i>Darevskia dahli</i> and Its Parental Bisexual (<i>D. portschinskii</i> , <i>D. mixta</i>) Rock Lizards in the Caucasus. 2020 , 8, 1329 | 3 |
| 751 | Predicting distribution of <i>Zanthoxylum bungeanum</i> Maxim. in China. 2020 , 20, 46 | 4 |
| 750 | How Climate Change Will Alter the Distribution of Suitable <i>Dendrobium</i> Habitats. 2020 , 8, | 7 |
| 749 | Ecological Niche Modeling: An Introduction for Veterinarians and Epidemiologists. 2020 , 7, 519059 | 7 |
| 748 | Fine-scale landscape genetics unveiling contemporary asymmetric movement of red panda (<i>Ailurus fulgens</i>) in Kangchenjunga landscape, India. 2020 , 10, 15446 | 11 |
| 747 | Closely related species show species-specific environmental responses and different spatial conservation needs: <i>Prionailurus</i> cats in the Indian subcontinent. 2020 , 10, 18705 | 5 |
| 746 | Identifying marine invasion hotspots using stacked species distribution models. 2020 , 22, 3403-3423 | 0 |
| 745 | Perceptions of Similarity Can Mislead Provenancing Strategies—An Example from Five Co-Distributed <i>Acacia</i> Species. 2020 , 12, 306 | 4 |
| 744 | A Model to Predict the Expansion of throughout the Iberian Peninsula Using a Pest Risk Analysis Approach. 2020 , 11, | 5 |
| 743 | High-resolution habitat suitability model for <i>Phlebotomus pedifer</i> , the vector of cutaneous leishmaniasis in southwestern Ethiopia. 2020 , 13, 467 | 1 |
| 742 | Quantifying range decline and remaining populations of the large marsupial carnivore of Australia's tropical rainforest. 2020 , 101, 1021-1034 | 2 |
| 741 | Evaluating the capacity of species distribution modeling to predict the geographic distribution of the mangrove community in Mexico. 2020 , 15, e0237701 | 4 |
| 740 | Patterns of niche contraction identify vital refuge areas for declining mammals. 2020 , 26, 1467-1482 | 10 |
| 739 | Gene reuse facilitates rapid radiation and independent adaptation to diverse habitats in the Asian honeybee. 2020 , 6, | 14 |
| 738 | Niche determinants in a salamander complex: Does hybridism or reproductive parasitism explain patterns of distribution?. 2020 , 11, e03265 | |
| 737 | Citizen science and habitat modelling facilitates conservation planning for crabeater seals in the Weddell Sea. 2020 , 26, 1291-1304 | 12 |

| | | |
|-----|--|-----|
| 736 | In or Out of the Checklist? DNA Barcoding and Distribution Modelling Unveil a New Species of <i>Crocidura</i> Shrew for Italy. 2020 , 12, 380 | 2 |
| 735 | Modelling the habitat of the endangered Carpentarian Grasswren (<i>Amytornis dorotheae</i>): The importance of spatio-temporal habitat availability in a fire prone landscape. 2020 , 24, e01341 | 1 |
| 734 | MaxEnt Modeling of <i>Dermacentor marginatus</i> (Acari: Ixodidae) Distribution in Xinjiang, China. 2020 , 57, 1659-1667 | 2 |
| 733 | Multiple dimensions of climate change on the distribution of Amazon primates. 2020 , 18, 83-90 | 10 |
| 732 | A standard protocol for reporting species distribution models. 2020 , 43, 1261-1277 | 141 |
| 731 | Effects of different variable sets on the potential distribution of fish species in the Amazon Basin. 2020 , 29, 764-778 | 2 |
| 730 | Potential Distribution and the Risks of and Its Associated Plant Pathogen <i>Liberibacter Solanacearum</i> for Global Potato Production. 2020 , 11, | 8 |
| 729 | Niche overlap between two sympatric frugivorous Neotropical primates: improving ecological niche models using closely-related taxa. 2020 , 29, 2749-2763 | 3 |
| 728 | Climate niche mismatch and the collapse of primate seed dispersal services in the Amazon. 2020 , 247, 108628 | 10 |
| 727 | Global analysis of ecological niche conservation and niche shift in exotic populations of monkeyflowers (<i>Mimulus guttatus</i> , <i>M. luteus</i>) and their hybrid (<i>M. robertsii</i>). 2020 , 13, 133-146 | 5 |
| 726 | Where and why? Bees, snail shells and climate: Distribution of <i>Rhodanthidium</i> (Hymenoptera: Megachilidae) in the Iberian Peninsula. 2020 , 23, 256-270 | 0 |
| 725 | Effect of El Niño Southern Oscillation cycle on the potential distribution of cutaneous leishmaniasis vector species in Colombia. 2020 , 14, e0008324 | 3 |
| 724 | Finding complexity in complexes: Assessing the causes of mitonuclear discordance in a problematic species complex of Mesoamerican toads. 2020 , 29, 3543-3559 | 11 |
| 723 | An ensemble high-resolution projection of changes in the future habitat of American lobster and sea scallop in the Northeast US continental shelf. 2020 , 26, 987-1001 | 11 |
| 722 | Altitudinal, latitudinal and longitudinal responses of cloud forest species to Quaternary glaciations in the northern Neotropics. 2020 , 130, 615-625 | 4 |
| 721 | Predicting the Potential Global Geographical Distribution of Two <i>Icerya</i> Species under Climate Change. 2020 , 11, 684 | 12 |
| 720 | Predicting the potential distribution of the vine mealybug, <i>Planococcus ficus</i> under climate change by MaxEnt. 2020 , 137, 105268 | 9 |
| 719 | Predicting the distribution of a rare chipmunk (<i></i>): comparing MaxEnt and occupancy models. 2020 , 101, 1035-1048 | 5 |

| | | |
|-----------------|--|----|
| 7 ¹⁸ | Potential distributions of <i>Bacillus anthracis</i> and <i>Bacillus cereus</i> biovar anthracis causing anthrax in Africa. 2020 , 14, e0008131 | 10 |
| 7 ¹⁷ | Optimized Maxent Model Predictions of Climate Change Impacts on the Suitable Distribution of <i>Cunninghamia lanceolata</i> in China. 2020 , 11, 302 | 32 |
| 7 ¹⁶ | Modelling risks posed by wind turbines and power lines to soaring birds: the black stork (<i>Ciconia nigra</i>) in Italy as a case study. 2020 , 29, 1959-1976 | 45 |
| 7 ¹⁵ | Potential impact of climate change on the distribution of the Eurasian Lynx (<i>Lynx lynx</i>) in Iran (Mammalia: Felidae). 2020 , 66, 107-117 | 5 |
| 7 ¹⁴ | A sequential multi-level framework to improve habitat suitability modelling. 2020 , 35, 1001-1020 | 8 |
| 7 ¹³ | Fragmenting fragments: landscape genetics of a subterranean rodent (Mammalia, Ctenomyidae) living in a human-impacted wetland. 2020 , 35, 1089-1106 | 5 |
| 7 ¹² | Modelling species distributions in dynamic landscapes: The importance of the temporal dimension. 2020 , 47, 1510-1529 | 8 |
| 7 ¹¹ | Prioritising search effort to locate previously unknown populations of endangered marine reptiles. 2020 , 22, e01013 | 4 |
| 7 ¹⁰ | Distribution of <i>Bactrocera oleae</i> (Rossi, 1790) throughout the Iberian Peninsula based on a maximum entropy modelling approach. 2020 , 177, 112-120 | 2 |
| 7 ⁰⁹ | Spatiotemporal Distribution of Human-Elephant Conflict in Eastern Thailand: A Model-Based Assessment Using News Reports and Remotely Sensed Data. 2020 , 12, 90 | 5 |
| 7 ⁰⁸ | Invasive fountain grass (<i>Pennisetum setaceum</i> (Forssk.) Chiov.) increases its potential area of distribution in Tenerife island under future climatic scenarios. 2020 , 221, 867-882 | 4 |
| 7 ⁰⁷ | Niche divergence and paleo-distributions of <i>Lutzomyia longipalpis</i> mitochondrial haplogroups (Diptera: Psychodidae). 2020 , 211, 105607 | 5 |
| 7 ⁰⁶ | Phylogeography and Genetic Diversity in a Southern North American Desert: From the Tehuac  -Cuicat  n Valley, Mexico. 2020 , 11, 863 | 4 |
| 7 ⁰⁵ | Paleoclimatic evolution as the main driver of current genomic diversity in the widespread and polymorphic Neotropical songbird <i>Arremon taciturnus</i> . 2020 , 29, 2922-2939 | 1 |
| 7 ⁰⁴ | Spatial predictors of genomic and phenotypic variation differ in a lowland Middle American bird (<i>Icterus gularis</i>). 2020 , 29, 3085-3102 | 2 |
| 7 ⁰³ | The combined role of dispersal and niche evolution in the diversification of Neotropical lizards. 2020 , 10, 2608-2625 | 12 |
| 7 ⁰² | Placing the hybrid origin of the asexual Amazon molly (<i>Poecilia formosa</i>) based on historical climate data. 2020 , 129, 835-843 | 2 |
| 7 ⁰¹ | Minimizing Risk and Maximizing Spatial Transferability: Challenges in Constructing a Useful Model of Potential Suitability for an Invasive Insect. 2020 , 113, 100-113 | 4 |

| | | |
|-----|--|----|
| 700 | Climate-induced changes in the suitable habitat of cold-water corals and commercially important deep-sea fishes in the North Atlantic. 2020 , 26, 2181 | 50 |
| 699 | The Potential Global Distribution of the White Peach Scale <i>Pseudaulacaspis pentagona</i> (Targioni Tozzetti) under Climate Change. 2020 , 11, 192 | 1 |
| 698 | Analysis of potentially suitable habitat within migration connections of an intra-African migrant-the Blue Swallow (<i>Hirundo atrocaerulea</i>). 2020 , 57, 101082 | 6 |
| 697 | Distribution of breeding humpback whale habitats and overlap with cumulative anthropogenic impacts in the Eastern Tropical Atlantic. 2020 , 26, 549-564 | 7 |
| 696 | Environmental Drivers and Distribution Patterns of Carnivoran Assemblages (Mammalia: Carnivora) in the Americas: Past to Present. 2020 , 27, 759-774 | 3 |
| 695 | Potential distribution of a climate sensitive species, the White-winged Snowfinch <i>Montifringilla nivalis</i> in Europe. 2020 , 30, 522-532 | 6 |
| 694 | Genomics of sorghum local adaptation to a parasitic plant. 2020 , 117, 4243-4251 | 31 |
| 693 | Evidence of ecological niche shift in (<i>L.</i>) in Britain: Hybridization as a possible cause of rapid niche expansion. 2020 , 10, 2040-2050 | 15 |
| 692 | Predicting the potential distribution of <i>Amblyomma americanum</i> (Acari: Ixodidae) infestation in New Zealand, using maximum entropy-based ecological niche modelling. 2020 , 80, 227-245 | 4 |
| 691 | Biogeographic barriers, Pleistocene refugia, and climatic gradients in the southeastern Nearctic drive diversification in cornsnakes (<i>Pantherophis guttatus</i> complex). 2020 , 29, 797-811 | 11 |
| 690 | Projected distribution and climate refugia of endangered Kashmir musk deer <i>Moschus cupreus</i> in greater Himalaya, South Asia. 2020 , 10, 1511 | 9 |
| 689 | Pollution control can help mitigate future climate change impact on European grayling in the UK. 2020 , 26, 517-532 | 2 |
| 688 | Mapping tree species vulnerability to multiple threats as a guide to restoration and conservation of tropical dry forests. 2020 , 26, 3552-3568 | 29 |
| 687 | Spatial sampling bias and model complexity in stream-based species distribution models: A case study of Paddlefish (<i>Lepisosteichthys</i>) in the Arkansas River basin, USA. 2020 , 10, 705-717 | 5 |
| 686 | Climatic Change and Habitat Availability for Three Sotol Species in México: A Vision towards Their Sustainable Use. 2020 , 12, 3455 | 1 |
| 685 | Climatic dynamics and topography control genetic variation in Atlantic Forest montane birds. 2020 , 148, 106812 | 6 |
| 684 | Predicting the Distribution of Indicator Taxa of Vulnerable Marine Ecosystems in the Arctic and Sub-arctic Waters of the Nordic Seas. 2020 , 7, | 10 |
| 683 | Ecological Niche Models of Four Hard Tick Genera (Ixodidae) in Mexico. 2020 , 10, | 4 |

| | | |
|-----|---|----|
| 682 | Distribution Pattern of Endangered Plant <i>Semiliquidambar cathayensis</i> (Hamamelidaceae) in Response to Climate Change after the Last Interglacial Period. 2020 , 11, 434 | 12 |
| 681 | Shifting ecosystem connectivity during the Pleistocene drove diversification and gene-flow in a species complex of Neotropical birds (Tityridae: <i>Pachyramphus</i>). 2020 , 47, 1714-1726 | 4 |
| 680 | Genotyping-by-sequencing and ecological niche modeling illuminate phylogeography, admixture, and Pleistocene range dynamics in quaking aspen (). 2020 , 10, 4609-4629 | 3 |
| 679 | Deer density drives habitat use of establishing wolves in the Western European Alps. 2020 , 57, 995-1008 | 4 |
| 678 | Changing Only Slowly: The Role of Phylogenetic Niche Conservatism in Caviidae (Rodentia) Speciation. 2020 , 27, 713-721 | 2 |
| 677 | Co-occurrence or dependence? Using spatial analyses to explore the interaction between palms and <i>Rhodnius</i> triatomines. 2020 , 13, 211 | 0 |
| 676 | Effects of climate change and human influence in the distribution and range overlap between two widely distributed avian scavengers. 2021 , 31, 77-95 | 1 |
| 675 | Including indigenous knowledge in species distribution modeling for increased ecological insights. 2021 , 35, 587-597 | 11 |
| 674 | Oh the places theyll go: improving species distribution modelling for invasive forest pests in an uncertain world. 2021 , 23, 297-349 | 10 |
| 673 | Current landscape attributes and landscape stability in breeding grounds explain genetic differentiation in a long-distance migratory bird. 2021 , 24, 120-134 | 5 |
| 672 | Land use and climate change interaction triggers contrasting trajectories of biological invasion. 2021 , 120, 106936 | 13 |
| 671 | Species distribution models predict the geographic expansion of an enzootic amphibian pathogen. 2021 , 53, 221-231 | 3 |
| 670 | Support for a relationship between demography and modeled habitat suitability is scale dependent for the purple martin <i>Progne subis</i> . 2021 , 90, 356-366 | 1 |
| 669 | Drivers of distributions and niches of North American cold-adapted amphibians: evaluating both climate and land use. 2021 , 31, e2236 | 3 |
| 668 | Idiosyncratic responses to drivers of genetic differentiation in the complex landscapes of Isthmian Central America. 2021 , 126, 251-265 | 0 |
| 667 | Cumulative habitat loss increases conservation threats on endemic species of terrestrial vertebrates in Mexico. 2021 , 253, 108864 | 4 |
| 666 | Genetic similarities versus morphological resemblance: Unraveling a polyploid complex in a Mediterranean biodiversity hotspot. 2021 , 155, 107006 | 3 |
| 665 | Improving biological relevance of model projections in response to climate change by considering dispersal amongst lineages in an amphibian. 2021 , 48, 561-576 | 4 |

| | | |
|-----|--|---|
| 664 | Ecological niche differentiation of two salamanders (Caudata: Hynobiidae) from Hokkaido Island, Japan. 2021 , 36, 281-292 | 2 |
| 663 | Coasting along to a wider range: niche conservatism in the recent range expansion of the Tawny Coster, <i>Acraea terpsicore</i> (Lepidoptera: Nymphalidae). 2021 , 27, 402-415 | 8 |
| 662 | Determining the minimal background area for species distribution models: MinBAR package. 2021 , 439, 109353 | 3 |
| 661 | The story of endurance: Biogeography and the evolutionary history of four Holarctic butterflies with different habitat requirements. 2021 , 48, 590-602 | 0 |
| 660 | Potential regional declines in species richness of tomato pollinators in North America under climate change. 2021 , 31, e02259 | 0 |
| 659 | Geographical pattern of genetic diversity in (Brassicaceae)-A global perspective. 2021 , 11, 199-213 | 4 |
| 658 | Landscape-scale biogeographic distribution analysis of the whitefly, <i>Bemisia tabaci</i> (Gennadius, 1889) in Kenya. 2021 , 41, 1585-1599 | 6 |
| 657 | Potential Distributions of the Invasive Barnacle Scale <i>Ceroplastes cirripediformis</i> (Hemiptera: Coccidae) Under Climate Change and Implications for Its Management. 2021 , 114, 82-89 | 4 |
| 656 | Isolation by environment and recurrent gene flow shaped the evolutionary history of a continentally distributed Neotropical treefrog. 2021 , 48, 760-772 | 7 |
| 655 | Geographic range estimates and environmental requirements for the harpy eagle derived from spatial models of current and past distribution. 2021 , 11, 481-497 | 4 |
| 654 | Genome-wide genetic variation coupled with demographic and ecological niche modeling of the dusky-footed woodrat (<i>Neotoma fuscipes</i>) reveal patterns of deep divergence and widespread Holocene expansion across northern California. 2021 , 126, 521-536 | 0 |
| 653 | White-faced darter distribution is associated with coniferous forests in Great Britain. 2021 , 14, 15-25 | |
| 652 | Influence of spatial extent on habitat suitability models for primate species of Atlantic Forest. 2021 , 61, 101179 | 3 |
| 651 | Balancing transferability and complexity of species distribution models for rare species conservation. 2021 , 27, 95-108 | 7 |
| 650 | Niche-based processes explaining the distributions of closely related subterranean spiders. 2021 , 48, 118-133 | 7 |
| 649 | Intelligent comprehensive evaluation system using artificial intelligence for environmental evaluation. 2021 , 86, 106495 | 2 |
| 648 | OUP accepted manuscript. | 0 |
| 647 | Predicting the current and future distribution of the western black-legged tick, <i>Ixodes pacificus</i> , across the Western US using citizen science collections. 2021 , 16, e0244754 | 6 |

| | | |
|-----|--|---|
| 646 | OUP accepted manuscript. | 1 |
| 645 | Signatures of genetic isolation of the three lineages of the narrow-headed vole <i>Lasiopodomys gregalis</i> (Cricetidae, Rodentia) in a mosaic steppe landscape of South Siberia. 2021 , 101, 275-285 | 1 |
| 644 | Spatial distribution of the pine marten (<i>Martes martes</i>) and stone marten (<i>Martes foina</i>) in the Italian Alps. 2021 , 101, 345-356 | 1 |
| 643 | The geographical origin, refugia, and diversification of honey bees (<i>Apis</i> spp.) based on biogeography and niche modeling. 2021 , 52, 367-377 | 3 |
| 642 | Modelling Beach Litter Accumulation on Mediterranean Coastal Landscapes: An Integrative Framework Using Species Distribution Models. 2021 , 10, 54 | 1 |
| 641 | Physiological and ecological warnings that dodders pose an exigent threat to farmlands in Eastern Africa. 2021 , 185, 1457-1467 | 2 |
| 640 | Predicting the invasion of the acoustic niche: Potential distribution and call transmission efficiency of a newly introduced frog in Cuba. 2021 , 19, 90-97 | 1 |
| 639 | Historical observations for a sparsely recorded woodland butterfly are used to determine factors driving distribution change and to identify conservation strategies. 2021 , 25, 109-121 | |
| 638 | Spatial modelling of <i>Acropora muricata</i> and <i>Porites lutea</i> distribution using environmental descriptors across Lakshadweep-Chagos Archipelago. 2021 , 41, 101619 | |
| 637 | Modeling Niches and Mapping Distributions. 2021 , 315-348 | |
| 636 | Projections for Mexico's Tropical Rainforests Considering Ecological Niche and Climate Change. 2021 , 12, 119 | 7 |
| 635 | Predictively modelling the distribution of the threatened brush-tailed rock-wallaby (<i>Petrogale penicillata</i>) in Oxley Wild Rivers National Park, north-eastern New South Wales, Australia. 2021 , | |
| 634 | Demographic History and Genomic Response to Environmental Changes in a Rapid Radiation of Wild Rats. 2021 , 38, 1905-1923 | 2 |
| 633 | Phenotypic plasticity and the colonization of new habitats: a study of a colonial spider in the Chaco region and the Cerrado. 2021 , 35, 235-251 | 0 |
| 632 | Intraspecific variation and taxonomy of northern pika <i>Ochotona hyperborea</i> (Mammalia, Lagomorpha). 2021 , 102, 28-53 | |
| 631 | Late Pliocene population divergence and persistence despite Pleistocene climatic fluctuations in the Rio Doce snouted Treefrog (<i>Ololygon carnevallii</i>). 2021 , 59, 680-690 | |
| 630 | Investigating niches and distribution of a rare species in a hierarchical framework: Virginia Warbler (<i>Leiothlypis virginiae</i>) at its northeastern range limit. 2021 , 36, 1039-1054 | 2 |
| 629 | Stability in the Balkans: phylogeography of the endemic Greek stream frog, <i>Rana graeca</i> . 2021 , 132, 829-846 | 1 |

| | | |
|-----|--|----|
| 628 | Predictive models of distribution and abundance of a threatened mountain species show that impacts of climate change overrule those of land use change. 2021 , 27, 989-1004 | 2 |
| 627 | Assessing the Potential Distributions of the Invasive Mosquito Vector and Its Natural Infections in Mexico. 2021 , 12, | 5 |
| 626 | Insights into the importance of areas of climatic stability in the evolution and maintenance of avian diversity in the Mesoamerican dry forests. 2021 , 132, 741-758 | 3 |
| 625 | Resistance-Based Connectivity Model to Construct Corridors of the Przewalski's Gazelle (<i>Procapra przewalskii</i>) in Fragmented Landscape. 2021 , 13, 1656 | 1 |
| 624 | Identification, Distribution, and Habitat Suitability Models of Ixodid Tick Species in Cattle in Eastern Bhutan. 2021 , 6, | 1 |
| 623 | A kingdom in decline: Holocene range contraction of the lion (<i>Panthera leo</i>) modelled with global environmental stratification. 2021 , 9, e10504 | 0 |
| 622 | Conservation status assessment of banana crop wild relatives using species distribution modelling. 2021 , 27, 729-746 | 11 |
| 621 | Invasive plants in Brazil: climate change effects and detection of suitable areas within conservation units. 2021 , 23, 1577-1594 | 5 |
| 620 | The Potential Distribution of <i>Neotoma floridana</i> in the Chincoteague National Wildlife Refuge, Virginia. 2021 , 8, 640339 | 2 |
| 619 | Ant-gardens: a specialized ant-epiphyte mutualism capable of facing the effects of climate change. 2021 , 30, 1165-1187 | 3 |
| 618 | Climate change shifts abiotic niche of temperate skates towards deeper zones in the Southwestern Atlantic Ocean. | |
| 617 | Environmental Factors Shape the Nonbreeding Distribution of the Harlan's Red-Tailed Hawk: A Maximum Entropy Approach. 2021 , 55, | 0 |
| 616 | Past, present, and future geographic range of the relict Mediterranean and Macaronesian complex. 2021 , 11, 5075-5095 | 5 |
| 615 | Generalists yet different: distributional responses to climate change may vary in opportunistic bat species sharing similar ecological traits. 2021 , 51, 571-584 | 20 |
| 614 | Niche Contraction of an Endangered Frog Driven by the Amphibian Chytrid Fungus. 2021 , 18, 134-144 | 1 |
| 613 | Predicting the Global Distribution of (Hymenoptera: Formicidae) under Climate Change Using the MaxEnt Model. 2021 , 12, | 6 |
| 612 | Prediction of Potential Geographical Distribution Patterns of <i>Actinidia arguta</i> under Different Climate Scenarios. 2021 , 13, 3526 | 2 |
| 611 | Habitat suitability of cetaceans in the Gulf of Mexico using an ecological niche modeling approach. 2021 , 9, e10834 | 1 |

| | | |
|-----|--|---|
| 610 | Climate Change and Local Host Availability Drive the Northern Range Boundary in the Rapid Expansion of a Specialist Insect Herbivore, <i>Papilio cressphontes</i> . 2021 , 9, | 3 |
| 609 | Combining the responses of habitat suitability and connectivity to climate change for an East Asian endemic frog. 2021 , 18, 14 | 2 |
| 608 | Tolerance niche expansion and potential distribution prediction during Asian openbill bird range expansion. 2021 , 11, 5562-5574 | 2 |
| 607 | Ecological niche modeling as an effective tool to predict the distribution of freshwater organisms: The case of the Sabaleta <i>Brycon henni</i> (Eigenmann, 1913). 2021 , 16, e0247876 | 5 |
| 606 | Vulnerable areas to accidents with scorpions in Brazil. 2021 , 26, 591-601 | 4 |
| 605 | Where is the genus <i>Elysia</i> in the western Atlantic? Potential distribution, species richness and representation in marine protected areas. 2021 , 87, | |
| 604 | Toward a Monte Carlo approach to selecting climate variables in MaxEnt. 2021 , 16, e0237208 | 3 |
| 603 | True blue: Temporal and spatial stability of pelagic wildlife at a submarine canyon. 2021 , 12, e03423 | 2 |
| 602 | How Threatened Is <i>Scincella huanrenensis</i> ? An Update on Threats and Trends. 2021 , 1, 58-72 | 2 |
| 601 | Distribution modelling of the rare stink bug <i>Ceratozygum horridum</i> (Germar, 1839): isolated in small spots across the Neotropics or a continuous population?. 2021 , 55, 649-663 | 0 |
| 600 | Reduction in the potential distribution of bumble bees (Apidae: <i>Bombus</i>) in Mesoamerica under different climate change scenarios: Conservation implications. 2021 , 27, 1772-1787 | 8 |
| 599 | Predicting migration routes for three species of migratory bats using species distribution models. 2021 , 9, e11177 | 1 |
| 598 | <i>Dichrocephala integrifolia</i> (Astereae, Asteraceae), a new exotic genus and species for Mexico and second record for the New World. 2021 , 1, | 0 |
| 597 | Effect of land use, habitat suitability, and hurricanes on the population connectivity of an endemic insular bat. 2021 , 11, 9115 | |
| 596 | Evolutionary history of a desert perennial (Boraginaceae): Intraspecific divergence, regional expansion and asymmetric gene flow.. 2021 , 43, 462-471 | 0 |
| 595 | The four antelope species on the Qinghai-Tibet plateau face habitat loss and redistribution to higher latitudes under climate change. 2021 , 123, 107337 | 2 |
| 594 | Mapping the Habitat Suitability of West Nile Virus Vectors in Southern Quebec and Eastern Ontario, Canada, with Species Distribution Modeling and Satellite Earth Observation Data. 2021 , 13, 1637 | 2 |
| 593 | Anthropogenic pressures increase extinction risk of an isolated Asian elephant (<i>Elephas maximus</i>) population in southwestern China, as revealed by a combination of molecular- and landscape-scale approaches. 2021 , | 1 |

| | | |
|-----|---|----|
| 592 | Potential distribution of aquatic invasive alien plants, <i>Eichhornia crassipes</i> and <i>Salvinia molesta</i> under climate change in Sri Lanka. 2021 , 29, 531-545 | 3 |
| 591 | Structure and robustness of the Neotropical ant-gardens network under climate change. 2021 , 14, 635-646 | 3 |
| 590 | Potential Global Distribution of under Climate Change Based on MaxEnt. 2021 , 12, | 1 |
| 589 | Evolving in islands of mud: old and structured hidden diversity in an endemic freshwater crayfish from the Chilean hotspot. 2021 , 11, 8573 | 3 |
| 588 | Butterflies are weakly protected in a mega-populated country, Bangladesh. 2021 , 26, e01484 | 3 |
| 587 | Where to sleep in the city? How urbanisation impacts roosting habitat availability for an apex predator. 2021 , 26, e01494 | 3 |
| 586 | Ecological niche modelling predicts significant impacts of future climate change on two endemic rodents in eastern Africa. 2021 , 13, 18164-18176 | |
| 585 | Multi-Scale Spatial Prediction of Wild Boar Damage Risk in Hunchun: A Key Tiger Range in China. 2021 , 11, | 0 |
| 584 | Phylogeographic history of Japanese macaques. 2021 , 48, 1420-1431 | 1 |
| 583 | Unusual host selection in larvae (Coleoptera: Chrysomelidae: Bruchinae) and its potential impact on stored almonds (). 2021 , 111, 585-594 | |
| 582 | Migration-tracking integrated phylogeography supports long-distance dispersal-driven divergence for a migratory bird species in the Japanese archipelago. 2021 , 11, 6066-6079 | 2 |
| 581 | Hybridization as a biodiversity driver: The case of <i>Veronica gundisalvi</i> . 42, e67901 | 3 |
| 580 | Species-Distribution Modeling: Advantages and Limitations of Its Application. 2. MaxEnt. 2021 , 11, 265-275 | 10 |
| 579 | Assessment of potential invasion for six phytophagous quarantine pests in Taiwan. 2021 , 11, 10666 | 1 |
| 578 | Modeling breeding habitats of humpback whales <i>Megaptera novaeangliae</i> as a function of group composition. 2021 , 666, 203-215 | 0 |
| 577 | Modeling the Potential Distribution of the Malaria Vector <i>Theobald</i> (Diptera: Culicidae) in Arid Regions of Northern Chile. 2021 , 9, 611152 | 0 |
| 576 | Environmental niche differentiation and paleodistribution of the rare montane woodrats of the genus <i>Nelsonia</i> (Rodentia: Cricetidae). 2021 , 101, 521-530 | 1 |
| 575 | Niche Modeling May Explain the Historical Population Failure of in Taiwan: Implications of Biocontrol Strategies. 2021 , 12, | 0 |

| | | |
|-----|--|--------|
| 574 | Inferring the Potential Distribution of an Emerging Rickettsiosis in America: The Case of. 2021 , 10, | 5 |
| 573 | Static species distribution models in the marine realm: The case of baleen whales in the Southern Ocean. 2021 , 27, 1536-1552 | 4 |
| 572 | Volcanism and palaeoclimate change drive diversification of the world's largest whip spider (Amblypygi). 2021 , 30, 2872-2890 | 3 |
| 571 | The geography of parasite local adaptation to host communities. 2021 , 44, 1205-1217 | 2 |
| 570 | Potential Distribution and Environmental Niche of the Black Corals <i>Antipathes galapagensis</i> and <i>Myriopathes panamensis</i> in the Eastern Tropical Pacific1. 2021 , 75, | 0 |
| 569 | Chikungunya Beyond the Tropics: Where and When Do We Expect Disease Transmission in Europe?. 2021 , 13, | 1 |
| 568 | Evaluating Sierra Nevada Yellow-Legged Frog Distribution Using Environmental DNA. 2021 , 85, 945-952 | 1 |
| 567 | Cautions in weighting individual ecological niche models in ensemble forecasting. 2021 , 448, 109502 | 5 |
| 566 | National assessments of species vulnerability to climate change strongly depend on selected data sources. 2021 , 27, 1367-1382 | 0 |
| 565 | Ensemble Models Predict Invasive Bee Habitat Suitability Will Expand under Future Climate Scenarios in Hawai'i. 2021 , 12, | 5 |
| 564 | Potential distribution models from two highly endemic species of subterranean rodents of Argentina: which environmental variables have better performance in highly specialized species?. 2021 , 101, 503-519 | 1 |
| 563 | In situ adaptation and ecological release facilitate the occupied niche expansion of a non-native Madagascan day gecko in Florida. 2021 , 11, 9410-9422 | 1 |
| 562 | Intraspecific genetic variation matters when predicting seagrass distribution under climate change. 2021 , 30, 3840-3855 | 3 |
| 561 | The impact of climate change on western salamanders' distribution. 2021 , 11, 9370-9384 | 4 |
| 560 | ENMeval 2.0: Redesigned for customizable and reproducible modeling of species niches and distributions. <i>Methods in Ecology and Evolution</i> , 2021 , 12, 1602-1608 | 7.7 24 |
| 559 | Modeling Seasonal Distribution of Irrawaddy Dolphins (<i>Orcaella brevirostris</i>) in a Transnational Important Marine Mammal Area. 2021 , 8, | 0 |
| 558 | Rapid shifts in Arctic tundra species' distributions and inter-specific range overlap under future climate change. 2021 , 27, 1706-1718 | 0 |
| 557 | Temporal matching of occurrence localities and forest cover data helps improve range estimates and predict climate change vulnerabilities. 2021 , 27, e01569 | 0 |

- 556 Phylogeography of *Habia fuscicauda* (Cardinalidae) indicates population isolation, genetic divergence and demographic changes during the Quaternary climate shifts in the Mesoamerican rainforest. **2021**, 162, 961 1
- 555 Species distribution modelling of the Southern Ocean benthos: a review on methods, cautions and solutions. **2021**, 33, 349-372 0
- 554 Open-source Tools in R for Landscape Ecology. **2021**, 6, 97-111 2
- 553 Seasonal spatial dynamics of butterfly migration. **2021**, 24, 1814-1823 1
- 552 Species distribution and conservation assessment of the black-headed night monkey (*Aotus nigriceps*): a species of Least Concern that faces widespread anthropogenic threats. **2021**, 62, 817-825 0
- 551 Climatic suitability of the eastern paralysis tick, *Ixodes holocyclus*, and its likely geographic distribution in the year 2050. **2021**, 11, 15330 2
- 550 Forecasting the Distribution of a Range-Expanding Bat Reveals Future Response to Climate Change and Habitat. **2021**, 23, 0
- 549 Future habitat suitability for species under climate change [Lessons learned from the strawberry tree case study. **2021**, 491, 119150 0
- 548 A genus at risk: Predicted current and future distribution of all three *Lagopus* species reveal sensitivity to climate change and efficacy of protected areas. **2021**, 27, 1759-1774 1
- 547 The role of dispersal limitation and reforestation in shaping the distributional shift of a forest herb under climate change. **2021**, 27, 1775-1791 2
- 546 Austral Yungas under future climate and land-use changes scenarios: the importance of protected areas for long-term amphibian conservation. **2021**, 30, 3335 1
- 545 Estimation of the Recent Expansion Rate of (Orthoptera) on a Regional and Landscape Scale. **2021**, 12, 0
- 544 Pliocene origins, Pleistocene refugia, and postglacial range expansions in southern devil scorpions (*Vaejovidae*: *Vaejovis carolinianus*). **2021**, 21, 575-590
- 543 Genetic structure and temporal environmental niche dynamics of sideoats grama [*Bouteloua curtipendula* (Michx.) Torr.] populations in Mexico. **2021**, 16, e0254566 0
- 542 Diversity, Host Plants and Potential Distribution of Edible Saturniid Caterpillars in Kenya. **2021**, 12, 3
- 541 Distribution patterns, ecological niche and conservation status of endemic *Tillandsia purpurea* along the Peruvian coast. **2021**, 307, 1 0
- 540 The specialist of a specialist: the natural history of the predispersal seed predator weevil *Hemicolpus abdominalis* (Coleoptera: Curculionidae). **2021**, 46, 1006-1018 0
- 539 An Appalachian population of *neochoristoderes* (Diapsida, Choristodera) elucidated using fossil evidence and ecological niche modelling. **2021**, 64, 629-643 1

| | | |
|-----|--|---|
| 538 | Global change on the roof of the world: Vulnerability of Himalayan otter species to land use and climate alterations. | 5 |
| 537 | Origin and Potential Expansion of the Invasive Longan Lanternfly, (Hemiptera: Fulgoridae) in Taiwan. 2021 , 10, | 0 |
| 536 | Historical climate change and megafaunal extinctions linked to genetic diversity declines in shorebirds. | 0 |
| 535 | Climate-based ensemble modelling to evaluate the global distribution of <i>Anoplophora glabripennis</i> (Motschulsky). 2021 , 23, 569 | 1 |
| 534 | Potential distributional shifts in North America of allelopathic invasive plant species under climate change models.. 2022 , 44, 11-19 | 4 |
| 533 | Climatic-niche evolution with key morphological innovations across clades within (Anura: Megophryidae). 2021 , 11, 10353-10368 | 1 |
| 532 | Modelado actual y futuro de la idoneidad de hábitat el ahuehuete (<i>Taxodium mucronatum</i> Ten.): una propuesta para conservación en México. 2021 , 99, 752-770 | 2 |
| 531 | Expected impacts of climate change on tree ferns distribution and diversity patterns in subtropical Atlantic Forest. 2021 , 19, 369-378 | 2 |
| 530 | Climate refugia for two Colombian endemic tamarin primates are critically under-protected. 2021 , 101, 531-543 | 0 |
| 529 | Priority areas for vulture conservation in the Horn of Africa largely fall outside the protected area network. 1-18 | 5 |
| 528 | Disease-driven mass mortality event leads to widespread extirpation and variable recovery potential of a marine predator across the eastern Pacific. 2021 , 288, 20211195 | 4 |
| 527 | Invaders from islands: thermal matching, potential or flexibility?. | |
| 526 | Ecological niche modelling for the conservation of endemic threatened squamates (lizards and snakes) in the Western Ghats. 2021 , 28, e01700 | 2 |
| 525 | Indian Grey Wolf and Striped Hyaena sharing from the same bowl: High niche overlap between top predators in a human-dominated landscape. 2021 , 28, e01682 | 2 |
| 524 | Pleistocene dynamics of the Eurasian steppe as a driving force of evolution: Phylogenetic history of the genus (<i>Brassicaceae</i>). 2021 , 11, 12697-12713 | 1 |
| 523 | Phylogeography of the <i>Poecilimon ampliatus</i> species group (Orthoptera: Tettigoniidae) in the context of the Pleistocene glacial cycles and the origin of the only thelytokous parthenogenetic phaneropterine bush-cricket. 79, 401-418 | 1 |
| 522 | Multi-decadal land use impacts across the vast range of an iconic threatened species. 2021 , 27, 2218 | 0 |
| 521 | Prediction of potentially suitable areas for the introduction of <i>Magnolia wufengensis</i> under climate change. 2021 , 127, 107762 | 6 |

- 520 Comparing the population history of Neotropical annual species: The role of climate change and hybridization between *Tithonia tubaeformis* and *T. rotundifolia* (Asteraceae). **2021**, 23, 962-973
- 519 Seascape genetics of the Atlantic spotted dolphin (*Stenella frontalis*) based on mitochondrial DNA. **2021**,
- 518 Exploring the effects of the quaternary glacial/interglacial cycles on the geographic distributions of tropical Andean rodents: species in the genus *Aepeomys* Thomas, 1898 (Thomasomyini: Sigmodontinae: Cricetidae) as a case study. 1-17 ○
- 517 Delimitation despite discordance: Evaluating the species limits of a confounding species complex in the face of mitonuclear discordance. **2021**, 11, 12739-12753 ○
- 516 United States port interceptions and predicted climate suitability of the red dwarf honeybee, *Apis florea* (Hymenoptera: Apidae), with notes on its identification. **2021**, 52, 915-932
- 515 Habitat use and population genetics of golden jackals in Iran: Insights from a generalist species in a highly heterogeneous landscape. ○
- 514 Global warming drives range shifts in spiny-tailed lizards (Squamata: Agamidae: *Uromastix*) in the African and Arabian deserts. **2021**, 191, 104522 ○
- 513 Mineral lick distribution modeling and NW Amazon conservation planning alternatives. **2021**, 30, 3409 ○
- 512 A Landscape Epidemiological Approach for Predicting Chronic Wasting Disease: A Case Study in Virginia, US. **2021**, 8, 698767 ○
- 511 Characterising the relationship between suitable habitat and gene flow for *Myotis bechsteinii* and *Eptesicus serotinus* in Britain. **2021**, 36, 3419
- 510 Distribution and habitat use of the Madagascar Peregrine Falcon: first estimates for area of habitat and population size.
- 509 Species and Phenotypic Distribution Models Reveal Population Differentiation in Ethiopian Indigenous Chickens. **2021**, 12, 723360 ○
- 508 Modeling the Pacific chub mackerel (*Scomber japonicus*) ecological niche and future scenarios in the northern Peruvian Current System. **2021**, 197, 102672 ○
- 507 Ice matters: Life-history strategies of two Antarctic seals dictate climate change eventualities in the Weddell Sea. **2021**, 27, 6252-6262 ○
- 506 Niche-separation and conservation biogeography of Madagascar fork-marked lemurs (*Cheirogaleidae*: *Phaner*): Evidence of a new cryptic species?. **2021**, 29, e01738 2
- 505 Little hope for the polyploid endemic Pyrenean Larkspur (*Delphinium montanum*): evidences from population genomics and Ecological Niche Modelling.
- 504 Influence of Pleistocene climatic oscillations on the phylogeography and demographic history of endemic vulnerable trees (section) of the Tropical Montane Cloud Forest in Mexico. **2021**, 9, e12181 ○
- 503 Potential distribution of the extremely endangered species *Ostrya rehderiana* (Betulaceae) in China under future climate change. **2021**, 1 ○

| | | |
|-----|---|----|
| 502 | Ecological niche evolution, speciation, and feedback loops: Investigating factors promoting niche evolution in Ordovician brachiopods of eastern Laurentia. 2021 , 578, 110555 | 2 |
| 501 | The future impact of climate and land-use changes on Anatolian ground squirrels under different scenarios. | 1 |
| 500 | Most Mexican hummingbirds lose under climate and land-use change: Long-term conservation implications. 2021 , 19, 487-487 | 1 |
| 499 | Want to model a species niche? A step-by-step guideline on correlative ecological niche modelling. 2021 , 456, 109671 | 9 |
| 498 | Habitat Suitability Modeling to Inform Seascape Connectivity Conservation and Management. 2021 , 13, 465 | 1 |
| 497 | Automatic variable selection in ecological niche modeling: A case study using Cassin's Sparrow (<i>Peucaea cassinii</i>). | |
| 496 | Artificial illumination influences niche segregation in bats. 2021 , 284, 117187 | 9 |
| 495 | Giant Tree Frog diversification in West and Central Africa: Isolation by physical barriers, climate, and reproductive traits. 2021 , | 0 |
| 494 | Bias assessments to expand research harnessing biological collections. 2021 , 36, 1071-1082 | 5 |
| 493 | Is Phylogeographic Congruence Predicted by Historical Habitat Stability, or Ecological Co-associations?. 2021 , 5, | 1 |
| 492 | Broadly Distributed but Genetically Fragmented: Demographic Consequences of Pleistocene Climatic Oscillations in a Common Iberian Grasshopper. 2021 , 5, | |
| 491 | Predicting potential suitable habitats of Chinese fir under current and future climatic scenarios based on Maxent model. 2021 , 64, 101393 | 11 |
| 490 | The role of habitat fragmentation in Pleistocene megafauna extinction in Eurasia. 2021 , 44, 1619 | 2 |
| 489 | White-bellied heron a species on the verge of extinction: Ensemble model reveals loss of habitats and resultant prolonged isolation driving the species to extinction. 2021 , 64, 101383 | 1 |
| 488 | Extremely Endangered Butterflies of Scattered Central European Dry Grasslands Under Current Habitat Alteration. 2021 , 5, | 0 |
| 487 | Driven to the edge: Species distribution modeling of a Clawed Salamander (<i>Hynobiidae</i> :) predicts range shifts and drastic decrease of suitable habitats in response to climate change. 2021 , 11, 14669-14688 | 0 |
| 486 | Local ecological knowledge reveals combined landscape effects of light pollution, habitat loss, and fragmentation on insect populations. 2021 , 262, 109311 | 4 |
| 485 | A multi-scale Maxent approach to model habitat suitability for the giant pandas in the Qionglai mountain, China. 2021 , 30, e01766 | 3 |

| | | |
|-----|--|-------|
| 484 | Where do nivicolous myxomycetes occur? [Modeling the potential worldwide distribution of <i>Physarum albescens</i> . 2021 , 53, 101079 | 0 |
| 483 | Fine-tuning niche models matters in invasion ecology. A lesson from the land planarian <i>Obama nungara</i> . 2021 , 457, 109686 | 2 |
| 482 | Current and potential geographic distribution of red palm mite (<i>Raoiella indica</i> Hirst) in Brazil. 2021 , 65, 101396 | 0 |
| 481 | Methodological analysis of bioclimatic variable selection in species distribution modeling with application to agricultural pests (<i>Metcalfa pruinosa</i> and <i>Spodoptera litura</i>). 2021 , 190, 106430 | 2 |
| 480 | Potential distribution of <i>Amblyomma mixtum</i> (Koch, 1844) in climate change scenarios in the Americas. 2021 , 12, 101812 | 1 |
| 479 | Impact of climate and land cover changes on the potential distribution of four endemic salamanders in Mexico. 2021 , 64, 126066 | 1 |
| 478 | Geographical distribution of As-hyperaccumulator <i>Pteris vittata</i> in China: Environmental factors and climate changes. 2022 , 803, 149864 | 4 |
| 477 | Phylogeography and systematics of <i>Algyroides</i> (Sauria: Lacertidae) of the Balkan Peninsula. 2021 , 50, 282-299 | 0 |
| 476 | Effects of climate change and land cover on the distributions of a critical tree family in the Philippines. 2021 , 11, 276 | 4 |
| 475 | A Maxent Predictive Model for Hunter-Gatherer Sites in the Southern Pampas, Argentina. 2021 , 7, | 2 |
| 474 | Lineage-level distribution models lead to more realistic climate change predictions for a threatened crayfish. 2021 , 27, 684-695 | 6 |
| 473 | Single-locus species delimitation and ecological niche modelling provide insights into the evolution, historical distribution and taxonomy of the Pacific chorus frogs. 2021 , 132, 612-633 | 1 |
| 472 | Climate change impact on the potential geographical distribution of two invading <i>Xylosandrus ambrosia</i> beetles. 2021 , 11, 1339 | 9 |
| 471 | Large-scale, image-based tree species mapping in a tropical forest using artificial perceptual learning. <i>Methods in Ecology and Evolution</i> , 2021 , 12, 608-618 | 7.7 5 |
| 470 | Improving area of occupancy estimates for parapatric species using distribution models and support vector machines. 2021 , 31, e02228 | 3 |
| 469 | The Predictive Power of Ecological Niche Modeling for Global Arbuscular Mycorrhizal Fungal Biogeography. 2017 , 143-158 | 13 |
| 468 | Analysis of the remaining habitat of an endemic species rediscovered. 2020 , 100, 307-314 | 1 |
| 467 | Comparing maximum entropy modelling methods to inform aquaculture site selection for novel seaweed species. 2020 , 429, 109071 | 8 |

| | | |
|-----|---|----|
| 466 | Ecological niche models as hypothesis generators of functional genetic differentiation and potential local adaptation in a Mediterranean alpine ecosystem. | 1 |
| 465 | modleR: a modular workflow to perform ecological niche modeling in R. | 3 |
| 464 | Spatial modeling cannot currently differentiate SARS-CoV-2 coronavirus and human distributions on the basis of climate in the United States. | 4 |
| 463 | A data-driven geospatial workflow to improve mapping species distributions and assessing extinction risk under the IUCN Red List. | 0 |
| 462 | Integrated environmental and genomic analysis reveals the drivers of local adaptation in African indigenous chickens. | 3 |
| 461 | Physiological and ecological warnings that Dodder pose an exigent threat to farmlands in Eastern Africa. | 0 |
| 460 | First genome-wide analysis of an endangered lichen reveals isolation by distance and strong population structure. | 2 |
| 459 | Using insects to detect, monitor and predict the distribution of <i>Xylella fastidiosa</i> : a case study in Corsica. | 4 |
| 458 | blockCV: an R package for generating spatially or environmentally separated folds for k-fold cross-validation of species distribution models. | 6 |
| 457 | Detecting spatial dynamics of range expansions with geo-referenced genomewide SNP data and the geographic spectrum of shared alleles. | 2 |
| 456 | Spatial distribution and spread potential of sixteen <i>Leptospira</i> serovars in a subtropical region of Brazil. | 1 |
| 455 | Assessment and representation of variability in ecological niche model predictions. | 3 |
| 454 | Evaluating species distribution models with discrimination accuracy is uninformative for many applications. | 2 |
| 453 | Phylogenomic approaches reveal how a climatic inversion and glacial refugia shape patterns of diversity in an African rain forest tree species. | 3 |
| 452 | Climate change and local host availability drive the northern range boundary in the rapid northward expansion of the eastern giant swallowtail butterfly. | 1 |
| 451 | Predictor complexity and feature selection affect Maxent model transferability: Evidence from global freshwater invasive species. 2021 , 27, 497-511 | 9 |
| 450 | Potential distribution of <i>Pythium insidiosum</i> in Rio Grande do Sul, Brazil, and projections to neighbour countries. 2018 , 65, 1671-1679 | 9 |
| 449 | <i>Schistosoma japonicum</i> transmission risk maps at present and under climate change in mainland China. 2017 , 11, e0006021 | 19 |

- 448 The Taxonomic Status of *Mazama bricenii* and the Significance of the Tñhira Depression for Mammalian Endemism in the Cordillera de Mérida, Venezuela. **2015**, 10, e0129113 21
- 447 Adding Biotic Interactions into Paleodistribution Models: A Host-Cleptoparasite Complex of Neotropical Orchid Bees. **2015**, 10, e0129890 8
- 446 Andean Condor (*Vultur gryphus*) in Ecuador: Geographic Distribution, Population Size and Extinction Risk. **2016**, 11, e0151827 20
- 445 Comparing Distribution of Harbour Porpoises (*Phocoena phocoena*) Derived from Satellite Telemetry and Passive Acoustic Monitoring. **2016**, 11, e0158788 10
- 444 Genetic and ecological insights into glacial refugia of walnut (*Juglans regia* L.). **2017**, 12, e0185974 20
- 443 Using distribution models to estimate blooms of phytosanitary cyanobacteria in Brazil. **2020**, 20, 2
- 442 Clearing up the Crystal Ball: Understanding Uncertainty in Future Climate Suitability Projections for Amphibians. **2020**, 76, 108 5
- 441 Extinctions of Threatened Frogs may Impact Ecosystems in a Global Hotspot of Anuran Diversity. **2020**, 76, 121 4
- 440 sdmbench: R package for benchmarking species distribution models. **2018**, 3, 847 1
- 439 Getting to the bottom of bycatch: a GIS-based toolbox to assess the risk of marine mammal bycatch. **2020**, 42, 37-57 8
- 438 Effects of spatial resolution on predicting the distribution of aquatic invasive species in nearshore marine environments. **2016**, 556, 17-30 15
- 437 Distributional changes in a guild of non-indigenous tunicates in the NW Atlantic under high-resolution climate projections. **2017**, 570, 173-186 6
- 436 Methods for improving species distribution models in data-poor areas: example of sub-Antarctic benthic species on the Kerguelen Plateau. **2018**, 594, 149-164 6
- 435 Long-term trends in habitat use and site fidelity by Australian humpback dolphins *Sousa sahulensis* in a near-urban embayment. **2018**, 603, 227-242 4
- 434 Human Habitat Selection: Using Tools from Wildlife Ecology to Predict Recreation in Natural Landscapes. **2019**, 39, 142 5
- 433 Predicting the Potential Distribution of the Sierra Nevada Red Fox in the Oregon Cascades. **2018**, 9, 351-366 2
- 432 Spatial modeling could not differentiate early SARS-CoV-2 cases from the distribution of humans on the basis of climate in the United States. **2020**, 8, e10140 4
- 431 Climatic niche comparison between closely related trans-Palearctic species of the genus (Insecta: Heteroptera: Miridae: Orthotylinae). **2020**, 8, e10517 5

| | | |
|-----|--|----|
| 430 | Climate-change-induced range shifts of three allergenic ragweeds (L.) in Europe and their potential impact on human health. 2017 , 5, e3104 | 30 |
| 429 | Maxent-directed field surveys identify new populations of narrowly endemic habitat specialists. 2017 , 5, e3632 | 37 |
| 428 | Predicting the potential distributions of the invasive cycad scale (Hemiptera: Diaspididae) under different climate change scenarios and the implications for management. 2018 , 6, e4832 | 21 |
| 427 | Historical connections among river basins and climatic changes explain the biogeographic history of a water rat. 2018 , 6, e5333 | 3 |
| 426 | First steps towards assessing the evolutionary history and phylogeography of a widely distributed Neotropical grassland bird (Motacillidae:). 2018 , 6, e5886 | 8 |
| 425 | Habitat suitability and connectivity inform a co-management policy of protected area network for Asian elephants in China. 2019 , 7, e6791 | 11 |
| 424 | Species delimitation of the ticks based on phylogenetic clustering and niche modeling. 2019 , 7, e6911 | 13 |
| 423 | One-class land-cover classification using MaxEnt: the effect of modelling parameterization on classification accuracy. 2019 , 7, e7016 | 7 |
| 422 | Climatic niche comparison across a cryptic species complex. 2019 , 7, e7042 | 4 |
| 421 | Spread of the non-native anemone Hüssermann & Fßterra, 2001 along the Humboldt-current large marine ecosystem: an ecological niche model approach. 2019 , 7, e7156 | 2 |
| 420 | Exploring snake occurrence records: Spatial biases and marginal gains from accessible social media. 2019 , 7, e8059 | 4 |
| 419 | Predicting hedgehog mortality risks on British roads using habitat suitability modelling. 2020 , 7, e8154 | 20 |
| 418 | (Hymenoptera, Mymaridae), a potential biocontrol agent to reduce populations of (Hemiptera, Aphrophoridae) the main vector of in Europe. 2020 , 8, e8591 | 7 |
| 417 | Genetic variation of the Chilean endemic long-haired mouse (Rodentia, Supramyomorph, Cricetidae) in a geographical and environmental context. 2020 , 8, e9517 | 6 |
| 416 | Including Host Availability and Climate Change Impacts on the Global Risk Area of <i>Carpomya pardalina</i> (Diptera: Tephritidae). 2021 , 9, | 0 |
| 415 | Genetic diversity and functional connectivity of a critically endangered salamander. | 2 |
| 414 | Life-history traits and habitat availability shape genomic diversity in birds: implications for conservation. 2021 , 288, 20211441 | 4 |
| 413 | Biogeography of <i>D. Don</i> (Bignoniaceae): Diversification, Andean Uplift and Niche Conservatism. 2021 , 12, 724057 | 1 |

- 412 Updated distribution maps of predominant *Culex* mosquitoes across the Americas. **2021**, 14, 547 2
- 411 Population Genetics and Biogeography of the Lungwort Lichen in North America Support Distinct Eastern and Western Gene Pools. **2021**, 2
- 410 ClineHelpR: an R package for genomic cline outlier detection and visualization. **2021**, 22, 501 0
- 409 Niche evolution and historical biogeography of lady slipper orchids in North America and Eurasia. **2021**, 48, 2727-2741 2
- 408 Species Distribution Models of the Loisel in Its Origin and Invasive Country Reveal an Ecological Niche Shift. **2021**, 12, 738769 2
- 407 Predictive performance of presence-only species distribution models: a benchmark study with reproducible code. e01486 17
- 406 Combining Remote Sensing and Species Distribution Modelling to Assess *Pinus hartwegii* Response to Climate Change and Land Use from Izta-Popo National Park, Mexico. **2021**, 10, 1037 0
- 405 Evidence of cryptic diversity in *Podarcis peloponnesiacus* and re-evaluation of its current taxonomy; insights from genetic, morphological, and ecological data. 0
- 404 SDSim: A generalized user friendly web ABM system to simulate spatiotemporal distribution of species under environmental scenarios. **2021**, 105234 2
- 403 Geographical isolation versus dispersal: Relictual alpine grasshoppers support a model of interglacial diversification with limited hybridization. **2021**, 0
- 402 Projected dynamics of breeding habitat suitability for a steppe-land bird warrant anticipatory conservation actions. 0
- 401 Identifying prioritized planting areas for medicinal plant *Thesium chinense* Turcz. under climate change in China. **2021**, 66, 101459 0
- 400 Analysis of the distribution pattern of Chinese *Ziziphus jujuba* under climate change based on optimized biomod2 and MaxEnt models. **2021**, 132, 108256 11
- 399 ENVIREM: An expanded set of bioclimatic and topographic variables increases flexibility and improves performance of ecological niche modeling. 1
- 398 Akaike information criteria and predictive geographical accuracy are not related in ecological niche modeling. 0
- 397 BioModelos: a collaborative online system to map species distributions. 0
- 396 Islands in the desert: Environmental distribution modelling of endemic flora reveals the extent of Pleistocene tropical relict vegetation in southern Arabia. 0
- 395 Mapping changes in the spatiotemporal distribution of lumpy skin disease virus. 0

- 394 Differentiation between wild and domesticated Ungulates based on ecological niches. 2
- 393 Co-occurrence or dependence? Using spatial analyses to explore the interaction between palms and triatomines (Chagas disease insect vectors).
- 392 Genomics of sorghum local adaptation to a parasitic plant. 0
- 391 Distribuci3n potencial bajo escenarios de cambio clim3tico de corales del g3nero Pocillopora (Anthozoa: Scleractinia) en el Pac3fico oriental tropical. **2019**, 90, 2
- 390 Climate change and forest plagues: assessing current and future impacts of diprionid sawflies on the pine forests of north-western Mexico. **2019**, 7, e7220 2
- 389 Effects of prey turnover on poison frog toxins: a landscape ecology approach to assess how biotic interactions affect species phenotypes. 1
- 388 Evaluating the boundaries of marine biogeographic regions of the Southwestern Atlantic using halacarid mites (Halacaridae), meiobenthic organisms with a low dispersal potential.
- 387 Barriers to coexistence limit the poleward range of a globally-distributed plant.
- 386 Assessing the Relative Role of Environmental Factors That Limit the Distribution of the Yucatan Rattlesnake (*Crotalus tzabcan*). **2020**, 54, 216 1
- 385 Distribution of *Malpighia mexicana* in Mexico and its implications for Barranca del R3o Santiago. **2021**, 32, 1095-1103 0
- 384 Comparative phylogeography reveals how a barrier filters and structures taxa in North American warm deserts.
- 383 Fine-scale landscape genetics unveiling contemporary asymmetric movement of red panda (*Ailurus fulgens*) in Kangchenjunga landscape, India.
- 382 Changes in the potential distribution of the guava fruit fly *Anastrepha striata* (Diptera, Tephritidae) during the current and possible future climate scenarios in Colombia.
- 381 Potential habitats of an alien species (*Asterias rubens* Linnaeus, 1758) in the Black Sea: its current and future distribution patterns. **2021**, 1 1
- 380 Environmental matching reveals non-uniform range-shift patterns in benthic marine Crustacea. **2021**, 168, 1 2
- 379 Species distribution modeling as an approach to studying the processes of landscape domestication in central southern Mexico. 1
- 378 Ecological niche complexity of invasive and native cryptic species of the *Bemisia tabaci* species complex in China. 1 2
- 377 Expected impacts of climate change on tree ferns distribution and diversity patterns in subtropical Atlantic Forest.

- 376 Ecological Niche Modeling and Other Tools for the Study of Avian Malaria Distribution in the Neotropics: A Short Literature Review. **2020**, 251-280 1
- 375 Predictive Habitat Model Reveals Specificity in a Broadly Distributed Forest Raptor, The Harpy Eagle. **2020**, 54, 0
- 374 Roles of hostplant availability and quality for the distribution and climate change response of a dietary specialist herbivore.
- 373 Influence of montane altitudinal ranges on species distribution models; evidence in Andean blow flies. **2020**, 8, e10370 0
- 372 Does the Mediterranean water shrew *Neomys anomalus* (Soricidae, Eulipotyphla) expand the eastern part of the distribution range?. **2020**, 19, 112-130 0
- 371 The role of littoral cliffs in the niche delimitation on a microendemic plant facing climate change. **2021**, 16, e0258976 0
- 370 Risks of giant hogweed (*Heracleum mantegazzianum*) range increase in North America. **2022**, 24, 299 2
- 369 Too little, too late? Conservation exigencies for Borneo inferred from biogeographic considerations of its endemic plant genera against intense landscape modifications. 1
- 368 Effects of climate change on the ecological niche of common hornbeam (*Carpinus betulus* L.). **2021**, 66, 101478 1
- 367 Potential geographic distribution of , the vector of Powassan virus.. **2021**, 46, 155-162 0
- 366 Potential areas of spread of *Trioza erytreae* over mainland Portugal and Spain. 1
- 365 Toward a Monte Carlo approach to selecting climate variables in MaxEnt: A case study using Cassin's Sparrow (*Peucaea cassinii*).
- 364 Distribution, richness and conservation of the genus (Lamiaceae) in the State of Michoacán, Mexico. **2020**, 8, e56827
- 363 Spatial modeling of the ecological niche of *Pinus greggii* Engelm. (Pinaceae): a species conservation proposal in Mexico under climatic change scenarios. **2020**, 13, 426-434 4
- 362 ClinePlotR: Visualizing genomic clines and detecting outliers in R. 0
- 361 Spatial Suitability Modeling of Zoonosis: Implicated Risk Areas of *B. anthracis* and Trends under climate change Scenarios in Ethiopia. 0
- 360 OUP accepted manuscript.
- 359 Not going with the flow: Ecological niche of a migratory seabird, the South American Tern *Sterna hirundinacea*. **2022**, 463, 109804 0

- 358 Predicting future distribution patterns of *Jatropha gossypifolia* L. in South Africa in response to climate change. **2022**, 146, 417-425 1
- 357 Assessing Suitable Habitats for Treefrog Species after Previous Declines in Costa Rica. **2021**, 13, 577 0
- 356 Effects of geographic variation in host plant resources for a specialist herbivore's contemporary and future distribution. **2021**, 12, e03822 1
- 355 Spatial patterns of phylogenetic diversity and endemism in the Western Ghats, India: A case study using ancient predatory arthropods.. **2021**, 11, 16499-16513 1
- 354 Potential range shifts and climatic refugia of rupicolous reptiles in a biodiversity hotspot of South Africa. **2021**, 48, 264-273 0
- 353 Changes in the potential distribution of the guava fruit fly (Diptera, Tephritidae) under current and possible future climate scenarios in Colombia. **2021**, 1-12
- 352 Modeling the past, present, and future distributions of endangered white abalone (*Haliotis sorenseni*) to inform recovery efforts in California. **2021**, 16, e0259716 0
- 351 Prediction of Potentially Suitable Distributions of *Codonopsis pilosula* in China Based on an Optimized MaxEnt Model. **2021**, 9, 2
- 350 River network rearrangements promote speciation in lowland Amazonian birds. 0
- 349 Updated range metrics and a global population estimate for the Critically Endangered Philippine Eagle using a spatial ensemble habitat model. 0
- 348 Potential distribution of the primary malaria vector *Anopheles gambiae* Giles [Diptera: Culicidae] in Southwest Nigeria under current and future climatic conditions. **2021**, 82, 0
- 347 Reconstruction and variability of tropical pollination networks in the Brazilian Atlantic Forest.
- 346 Ecological assessment and environmental niche modelling of Himalayan rhubarb (*Rheum webbianum* Royle) in northwest Himalaya. **2021**, 16, e0259345 2
- 345 Genome-scale data reveal deep lineage divergence and a complex demographic history in the Texas horned lizard (*Phrynosoma cornutum*) throughout the southwestern and central US. **2021**, 1
- 344 Niche dynamics of in Sri Lanka: Distribution patterns, climate change effects, and conservation priorities.. **2021**, 11, 18196-18215
- 343 Reconsideration of the native range of the Chinese Swamp Cypress (*Glyptostrobus pensilis*) based on new insights from historic, remnant and planted populations. **2021**, 32, e01927 0
- 342 Temporal changes in the potential geographic distribution of (Chiroptera, Vespertilionidae), the "decade effect".. **2021**, 11, 16972-16980
- 341 Modelling habitat suitability of the invasive tick *Rhipicephalus microplus* in West Africa.. **2022**,

- 340 Predicting the in-between: Present and future habitat suitability of an intertidal euryhaline fish. **2022**, 68, 101523
- 339 Climatic Suitability for *Haplodrassus rufipes* in a Mediterranean Area: Linking a Predaceous Species to the Olive Grove. **2020**, 4,
- 338 MODELING THE DISTRIBUTION OF THE SOUTHERN YELLOW-CHEEKED GIBBON (*NOMASCUS GABRIELLAE*) USING MAXENT. **2021**, 59,
- 337 Estimating real snakebite incidence in Colombia by using mathematical modelling and statistical inference.
- 336 Molecular phylogeography and evolutionary history of the pink rice borer (Lepidoptera: Noctuidae): Implications for refugia identification and pest management. o
- 335 Reduced range size and Important Bird and Biodiversity Area coverage for the Harpy Eagle predicted from future climate change scenarios. 1
- 334 Predicting the potential distribution of the fall armyworm *Spodoptera frugiperda* (J.E. Smith) under climate change in China. **2022**, 33, e01994 o
- 333 Effects of climate change in the seas of China: Predicted changes in the distribution of fish species and diversity. **2022**, 134, 108489 1
- 332 Fine-Tuned Ecological Niche Models Unveil Climatic Suitability and Association with Vegetation Groups for Selected *Chaetocnema* Species in South Africa (Coleoptera: Chrysomelidae). **2022**, 14, 100 o
- 331 Presence and potential distribution of malaria-infected New World primates of Costa Rica.. **2022**, 21, 17 o
- 330 Systematic study and niche differentiation of the genus *Aporocactus* (Hylocereae, Cactoideae, Cactaceae). **2021**, 1, o
- 329 Mapping the habitat suitability of *Ottelia* species in Africa. **2022**, 1
- 328 Automatic variable selection in ecological niche modeling: A case study using Cassin's Sparrow (*Peucaea cassinii*).. **2022**, 17, e0257502 o
- 327 Bioclimatic habitat limitations for argan trees (*Argania spinosa* (L.) Skeels) in Northern Africa and Spain. **2022**, 22, 1 o
- 326 Phylogeography of the Neotropical sciurid *Guerlinguetus brasiliensis* (Rodentia: Sciuridae).
- 325 Connectivity conservation at the crossroads: protected areas versus payments for ecosystem services in conserving connectivity for Colombian carnivores.. **2022**, 9, 201154 1
- 324 Biotic factors limit the invasion of the plague pathogen (*Yersinia pestis*) in novel geographical settings. o
- 323 Endemism, projected climate change, and identifying species of critical concern in the Scrub Mint clade (Lamiaceae). o

| | | |
|-----|---|---|
| 322 | Modeling the Climate Suitability of Northernmost Mangroves in China under Climate Change Scenarios. 2022 , 13, 64 | 2 |
| 321 | Groundwater Potential Mapping Using Maximum Entropy. 2022 , 239-256 | 0 |
| 320 | Variable vulnerability to climate change in New Zealand lizards. | 3 |
| 319 | Predicting potential suitable habitat for <i>Ensete glaucum</i> (Roxb.) Cheesman using MaxEnt modelling. 2022 , 287, 152007 | 0 |
| 318 | Ancient introgression underlying the unusual mito-nuclear discordance and coat phenotypic variation in the Moupin pika. | 1 |
| 317 | Distribution limits, natural history and conservation status of the poorly known Peruvian gracile mouse opossum (<i>Didelphimorphia</i> : <i>Didelphidae</i>). 1-17 | 0 |
| 316 | Including climate change to predict the global suitable area of an invasive pest: <i>Bactrocera correcta</i> (Diptera: Tephritidae). 2022 , 34, e02021 | 1 |
| 315 | Ecological divergence and synchronous Pleistocene diversification in the widespread South American butter frog complex.. 2022 , 107398 | |
| 314 | Distribution and conservation status of the mountain tapir (<i>Tapirus pinchaque</i>) in Peru. 2022 , 66, 126130 | |
| 313 | Distribution modelling of deep-sea elasmobranchs on the Mid-Atlantic Ridge for spatial management. 2022 , 182, 103707 | 0 |
| 312 | Shedding light on a threatened midwife toad: fine-scale niche modelling of the Moroccan endemic <i>Alytes maurus</i> (Pasteur and Bons 1962). 1-11 | 0 |
| 311 | Goitered Gazelle's (<i>Gazella Subgutturosa</i>) Habitat Desirability Modeling by Using Maximum Entropy (Maxent) Method. | |
| 310 | EU-Trees4F, a dataset on the future distribution of European tree species.. 2022 , 9, 37 | 1 |
| 309 | Threatened skates exhibit abiotic niche stability despite climate change in the Southwestern Atlantic Ocean. | |
| 308 | Prediction of Suitable Distribution of a Critically Endangered Plant <i>Glyptostrobus pensilis</i> . 2022 , 13, 257 | 0 |
| 307 | Functional divergence from ecological baselines on Caribbean coral reefs. | 1 |
| 306 | Tropical and subtropical Asia's valued tree species under threat. 2021 , | 1 |
| 305 | One step ahead of sea anemone invasions with ecological niche modeling: potential distributions and niche dynamics of three successful invasive species. | 1 |

- 304 Conservation assessment of a range-edge population of *Trichophorum planifolium* (Cyperaceae) reveals range-wide inbreeding and locally divergent environmental conditions. 0
- 303 Modeling the risk of illegal forest activity and its distribution in the southeastern region of the Sierra Madre Mountain Range, Philippines. **2022**, 15, 63-70
- 302 Species distribution models and a 60-year-old transplant experiment reveal inhibited forest plant range shifts under climate change. **2022**, 49, 537-550 2
- 301 Prediction of Potentially Suitable Distribution Areas for *Prunus tomentosa* in China Based on an Optimized MaxEnt Model. **2022**, 13, 381 1
- 300 Strengthening Angel Shark Conservation in the Northeastern Mediterranean Sea. **2022**, 10, 269 0
- 299 European Ground Squirrels at the Edge: Current Distribution Status and Anticipated Impact of Climate on Europe's Southernmost Population. **2022**, 11, 301 0
- 298 Computer vision for assessing species color pattern variation from web-based community science images.
- 297 Past climate cooling promoted global dispersal of amphipods from Tian Shan montane lakes to circumboreal lakes.. **2022**, 1
- 296 Synergistic effects of climate and landscape change on the conservation of Amazonian lizards.. **2022**, 10, e13028
- 295 The effect of past defaunation on ranges, niches, and future biodiversity forecasts.. **2022**, 1
- 294 Little hope for the polyploid endemic Pyrenean Larkspur (*Primula*): Evidences from population genomics and Ecological Niche Modeling.. **2022**, 12, e8711 2
- 293 Native range estimates for red-listed vascular plants.. **2022**, 9, 117 0
- 292 Climate biogeography of *Arabidopsis thaliana*: linking distribution models, individual performance, and life history. 0
- 291 Spatio-temporal dynamics of rabies and habitat suitability of the common marmoset *Callithrix jacchus* in Brazil.. **2022**, 16, e0010254
- 290 Linking ecological niche models and common garden experiments to predict phenotypic differentiation in stressful environments: Assessing the adaptive value of marginal populations in an alpine plant.. **2022**, 0
- 289 'Fly to a Safer North': Distributional Shifts of the Orchid *L. Due* to Climate Change.. **2022**, 11, 1
- 288 An isolated population of the secretive, endemic Aztec Rail (*Rallus tenuirostris*) in Chihuahua, Mexico. **2022**, 133,
- 287 Predicted changes in the functional structure of earthworm assemblages in France driven by climate change. 0

- 286 Modelado Espacial Actual y Futuro de la Idoneidad de HBitat de Triatoma nitida Usinger1 en Latinoamfca. **2022**, 47,
- 285 Climate change effects on the global distribution and range shifts of citrus longhorned beetle Anoplophora chinensis.
- 284 Landscape Genomics Provides Evidence of Ecotypic Adaptation and a Barrier to Gene Flow at Treeline for the Arctic Foundation Species .. **2022**, 13, 860439
- 283 Climate-induced Range Shifts of Invasive Species (Diaphorina citri Kuwayama).. **2022**, 3
- 282 The genomic landscapes of desert birds are structured by contemporary features. 0
- 281 Historical connections between Atlantic Forest and Amazonia drove genetic and ecological diversity in Lithobates palmipes (Anura, Ranidae). **2022**, 20, 19-19
- 280 Potential geographic distribution of relict plant Pteroceltis tatarinowii in China under climate change scenarios.. **2022**, 17, e0266133 2
- 279 Prey resources are more important than climatic conditions for predicting the distribution of a broad-ranged apex predator.
- 278 Does size matter? An analysis of the niche width and vulnerability to climate change of fourteen species of the genus from North America.. **2022**, 10, e13154
- 277 The tangled evolutionary history of a long-debated Mesoamerican taxon: The Velazquez Woodpecker (Melanerpes santacruzi, Aves: Picidae).. **2022**, 170, 107445 0
- 276 Estimates of breeding season location for 4 mesic prairie bird species wintering along the Gulf Coast. **2021**, 133, 1
- 275 A spatial ecological analysis of the reintroduction of the Eurasian beaver. **2021**, 937, 022003
- 274 Global invasion risk of Apocephalus borealis, a honey bee parasitoid. **2021**, 52, 1128-1140
- 273 Are Climates in Canada and the United States Suitable for the European Spruce Bark Beetle, Ips typographus, and Its Fungal Associate, Endoconidiophora polonica?. **2021**, 12, 1725 1
- 272 Urban Environments Aid Invasion of Brown Widows (Theridiidae: Latrodectus geometricus) in North America, Constraining Regions of Overlap and Mitigating Potential Impact on Native Widows. **2021**, 9, 1
- 271 Exploring the Potential Distribution of Relic Trochodendron aralioides: An Approach Using Open-Access Resources and Free Software. **2021**, 12, 1749
- 270 Estimating the Potential Impacts of Climate Change on the Spatial Distribution of Garuga forrestii, an Endemic Species in China. **2021**, 12, 1708 1
- 269 The Cenozoic history of palms: Global diversification, biogeography and the decline of megathermal forests. **2022**, 31, 425-439 4

- 268 Centres of endemism in Borneo and their environmental correlates revealed by endemic plant genera. **2021**, 285, 151966 0
- 267 Seed Menus: An integrated decision-support framework for native plant restoration in the Mojave Desert.. **2022**, 12, e8805 1
- 266 Widespread occurrence of in Ontario, Canada, and predicted habitat suitability for the emerging .. **2022**, 12, e8798 0
- 265 Climatic niche comparisons of eastern North American and eastern Asian disjunct plant genera. 1
- 264 Biodiversity and conservation of Solar-powered sea slugs from the Western Atlantic under climate change scenarios.
- 263 Ecological niche models reveal the potential zones of invasion of the cobia (*Rachycentron canadum*) in the Eastern Pacific Ocean. **2022**, 849, 2413
- 262 Identifying climate refugia for high-elevation Alpine birds under current climate warming predictions.. **2022**, 3
- 261 Modelling potential distribution of an endemic plant (*Ferula Microcolea*) based on bioclimatic variables in Iran. **2022**, 64, 157-172
- 260 Post-Pleistocene dispersal explains the Rapoport effect in North American salamanders.
- 259 Identification of risk areas for and in China, based on the major host plant and CMIP6 climate scenarios.. **2022**, 12, e8824 2
- 258 Global disparity of research allocation and the Aichi biodiversity conservation targets.
- 257 The Colombian Caribbean Sea: a tropical habitat for the Vulnerable sperm whale *Physeter macrocephalus*?. 1-11
- 256 Changing patterns of conflict between humans, carnivores and crop-raiding prey as large carnivores recolonize human-dominated landscapes. **2022**, 269, 109553 0
- 255 Data_Sheet_1.PDF. **2020**,
- 254 Data_Sheet_1.docx. **2019**,
- 253 Table_1.xlsx. **2019**,
- 252 Data_Sheet_1.docx. **2019**,
- 251 Data_Sheet_1.docx. **2019**,

250 Data_Sheet_1.docx. **2020**,

249 Image_1.PDF. **2018**,

248 Image_2.pdf. **2018**,

247 DataSheet_1.zip. **2019**,

246 Data_Sheet_1.pdf. **2018**,

245 River network rearrangements promote speciation in lowland Amazonian birds.. **2022**, 8, eabn1099

o

244 OUP accepted manuscript.

243 Differentiated historical demography and ecological niche forming present distribution and genetic structure in coexisting two salamanders (Amphibia, Urodela, Hynobiidae) in a small island, Japan.. **2022**, 10, e13202

242 Extensive protected area coverage and an updated global population estimate for the Endangered Madagascar Serpent-eagle identified from species-habitat associations using remote sensing data.

241 Climate warming since the Holocene accelerates west-east communication for a Eurasian temperate species *Aquarius paludum*.. **2022**,

1

240 Sustainable human population density in Western Europe between 560.000 and 360.000 years ago.. **2022**, 12, 6907

o

239 Effects of future climate change on the forests of Madagascar. **2022**, 13,

3

238 Predicting the distribution of Australian frogs and their overlap with *Batrachochytrium dendrobatidis* under climate change.

o

237 Niche and geographical expansions of North American trees and tall shrubs in Europe.

o

236 MaxEnt Modelling and Impact of Climate Change on Habitat Suitability Variations of Economically Important Chilgoza Pine (*Pinus gerardiana* Wall.) in South Asia. **2022**, 13, 715

9

235 Feature tuning improves MAXENT predictions of the potential distribution of *Rudolph* and its variant.. **2022**, 10, e13337

o

234 Competition mediates understorey species range shifts under climate change.

o

233 A gap analysis of reconnaissance surveys assessing the impact of the 2019-20 wildfires on vertebrates in Australia. **2022**, 270, 109573

- 232 Extinction risk assessment of the endemic terrestrial vertebrates in Mexico. **2022**, 270, 109562 0
- 231 Site-specific scaling of remote sensing-based estimates of woody cover and aboveground biomass for mapping long-term tropical dry forest degradation status. **2022**, 276, 113040 0
- 230 POTENTIAL DISTRIBUTION OF DOMESTICATED *Sechium edule* (CUCURBITACEAE) IN MEXICO. **2022**, 27, 0
- 229 Predicting the Risk of Exotic Plant Invasions in the Orinoco Region: Importance of Distribution Models, Climatic Niche and Functional Richness. **2022**, 10, 0
- 228 VenomMaps: Updated species distribution maps and models for New World pitvipers (Viperidae: Crotalinae). **2022**, 9, 0
- 227 Habitat distribution change of commercial species in the Adriatic Sea during the COVID-19 pandemic. **2022**, 101675 0
- 226 Climatic niche evolution of infectious diseases driving amphibian declines. 0
- 225 Karuk ecological fire management practices promote elk habitat in northern California. 0
- 224 Assessing the distribution of invasive Asian mosquitoes in Northern Italy and modelling the potential spread of *Aedes koreicus* in Europe. **2022**, 232, 106536 0
- 223 The future impact of climate and land-use changes on Anatolian ground squirrels under different scenarios. **2022**, 101693 2
- 222 Two Sides of the Same Desert: Floristic Connectivity and Isolation Along the Hyperarid Coast and Precordillera in Peru and Chile. 10, 0
- 221 Predicting habitat suitability and range shifts under projected climate change for two octocorals in the north-east Atlantic. 10, e13509 0
- 220 Intertidal beach habitat suitability model for Pacific sand lance, *Ammodytes personatus*, in the Salish Sea, Canada. 0
- 219 Fin whale (*Balaenoptera physalus*) distribution modeling on their Nordic and Barents Seas feeding grounds. 0
- 218 Predicted Pleistocene-Holocene range and connectivity declines of the vulnerable fishing cat and insights for current conservation. 0
- 217 A MaxEnt modelling approach to understand the climate change effects on the distributional range of White-bellied Sholokili *Sholicola albiventris* (Blanford, 1868) in the Western Ghats, India. **2022**, 101702 1
- 216 Cabruca agroforestry systems reduce vulnerability of cacao plantations to climate change in southern Bahia. **2022**, 42, 0
- 215 Potential global distribution of a temperate marine coastal predator: The role of barriers and dispersal corridors on subpopulation connectivity. 1

- 214 Distribution of the greater naked-tailed armadillo *Cabassous tatouay* (Desmarest, 1804) in South America, with new records and species distribution modeling. 1-9
- 213 Potential distributions of the parasite *Trypanosoma cruzi* and its vector *Dipetalogaster maxima* highlight areas at risk of Chagas disease transmission in Baja California Sur, Mexico, under climate change. ○
- 212 Nuclear and Mitochondrial Phylogenomics of the Sifakas Reveal Cryptic Variation in the Diademed Sifaka. **2022**, 13, 1026
- 211 Isolation by distance and past climate resistance shaped the distribution of genealogical lineages of a neotropical lizard. **2022**, 20,
- 210 Distribution and habitat use of the Madagascar Peregrine Falcon: first estimates for area of habitat and population size. 1-17 ○
- 209 Distribution models calibrated with independent field data predict two million ancient and veteran trees in England. ○
- 208 Climate change on potential suitable areas for typical Pinaceae species on the Western Sichuan Plateau. 1-19
- 207 Finding what you don't know: Testing SDM methods for poorly known species. 1
- 206 Comprehensively evaluating the performance of species distribution models across clades and resolutions: choosing the right tool for the job. ○
- 205 Extreme environments filter functionally rich communities of Atlantic Forest treefrogs along altitudinal and latitudinal gradients.
- 204 Current and future geographic patterns of bird diversity dimensions of the Yucatan Peninsula and their representativeness in natural protected areas. **2022**, 8, 242-252 1
- 203 Ecological correlates of extinction risk and persistence of direct-developing stream-dwelling frogs in Mesoamerica. **2022**, 38, e02197 1
- 202 Local abundance of neotropical orchid bees in Amazon forests not related to large-scale climate suitability.
- 201 Predicting the potential distribution of a near-extinct avian predator on the Arabian Peninsula: implications for its conservation management. **2022**, 194, ○
- 200 A framework for near-real time monitoring of diversity patterns based on indirect remote sensing, with an application in the Brazilian Atlantic rainforest. 10, e13534
- 199 High Wind Speed Prevents the Establishment of the Disease Vector Mosquito *Aedes albopictus* in Its Climatic Niche in Europe. 10, ○
- 198 Constructing ecological indices for urban environments using species distribution models.
- 197 Introduction of high-value *Crocus sativus* (saffron) cultivation in non-traditional regions of India through ecological modelling. **2022**, 12, ○

- 196 Predicted impacts of climate change on wild and commercial berry habitats will have food security, conservation and agricultural implications. **2022**, 157341 o
- 195 Phylogenomics of arboreal alligator lizards shed light on the geographical diversification of cloud forest-adapted biotas. o
- 194 Changes in Climate Extremes and Their Effect on Maize (*Zea mays* L.) Suitability Over Southern Africa. 4,
- 193 Simulation of the Potential Suitable Distribution of the Endangered *Cremastra appendiculata* in China Under Global Climate Change. 10,
- 192 Niche diversification of Mediterranean and southwestern Asian tortoises. 10, e13702 o
- 191 Forecasting shifts in habitat suitability of three marine predators suggests a rapid decline in inter-specific overlap under future climate change. **2022**, 12,
- 190 Long-term funding of community projects has contributed to mitigation of illegal activities within a premier African protected area, Bwindi impenetrable National Park, Uganda. o
- 189 A novel multivariate ecological approach to modeling freshwater mussel habitats verified by ground truthing. o
- 188 Identification of degradation factors in mountain semiarid rangelands using spatial distribution modelling and ecological niche theory. 1-17
- 187 Ecological specialization and reproductive isolation among closely related sympatric ant-eating spiders.
- 186 Modelling the current and future potential distribution of the bean bug *Riptortus pedestris* with increasingly serious damage to soybean. o
- 185 Prediction of the potential distribution of the predatory mite *Neoseiulus californicus* (McGregor) in China under current and future climate scenarios. **2022**, 12, 1
- 184 Computer vision for assessing species color pattern variation from web-based community science images. **2022**, 104784 2
- 183 Biogeography of cereal stemborers and their natural enemies: forecasting pest management efficacy under changing climate. o
- 182 Determination of priority areas for amphibian conservation in Guerrero (Mexico), through systematic conservation planning tools. **2022**, 68, 126235 o
- 181 Habitat Suitability Assessment of Black-Necked Crane (*Grus nigricollis*) in the Zoige Grassland Wetland Ecological Function Zone on the Eastern Tibetan Plateau. **2022**, 14, 579 1
- 180 Climate change and cocoyam (*Colocasia esculenta* (L.) Schott) production: assessing impacts and potential adaptation strategies in Zimbabwe. **2022**, 27,
- 179 Distribution Prediction of *Hyalomma asiaticum* (Acari: Ixodidae) in a Localized Region in Northwestern China. **2022**, 108,

- 178 Habitat Suitability and Population Size Estimates for the Mongoose Lemurs (*Eulemur mongoz*) of Madagascar and the Comoro Islands, and Implications for Their Conservation.
- 177 Species distribution models applied to mosquitoes: Use, quality assessment, and recommendations for best practice. **2022**, 472, 110073 ○
- 176 Stay home, stay safe? High habitat suitability and environmental connectivity increases road mortality in a colonizing mesocarnivore. **2022**, 37, 2343-2361
- 175 Ecological Niche Modeling of Water Lily (*Nymphaea* L.) Species in Australia under Climate Change to Ascertain Habitat Suitability for Conservation Measures. **2022**, 11, 1874 ○
- 174 Climate Change Impact on the Habitat Suitability of *Pseudotsuga menziesii* Mirb. Franco in Mexico: An Approach for Its Conservation. **2022**, 14, 8888 ○
- 173 The combination of genomic offset and niche modelling provides insights into climate change-driven vulnerability. **2022**, 13, 1
- 172 Identifying the natural reserve area of *Cistanche salsa* under the effects of multiple host plants and climate change conditions using a maximum entropy model in Xinjiang, China. 13, ○
- 171 Assessing genomic and ecological differentiation among subspecies of the Rough-footed Mud Turtle, *Kinosternon hirtipes*.
- 170 Estimation of the potential geographical distribution of a new potato pest (*Schrankia costae* strigalis) in China under climate change. **2022**,
- 169 Long term (1985–2018) changes of the habitat suitability of European spruce assessed by Maxent modelling based on Landsat satellite imagery: A case study from a mountain landscape of Central Bulgaria. **2022**, 68, 277-292
- 168 Trees planted under a global restoration pledge have mixed futures under climate change. 1
- 167 Linking life history to landscape for threatened species conservation in a multiuse region.
- 166 Ecological niche modelling for delineating livestock ecotypes and exploring environmental genomic adaptation: The example of Ethiopian village chicken. 10, ○
- 165 Predicting the potential distribution of four endangered holoparasites and their primary hosts in China under climate change. 13, ○
- 164 Optimising the benefit–cost ratio of fishing grounds for a multi-species fishery in the waters of northern Taiwan.
- 163 Assessing the Effectiveness of Correlative Ecological Niche Model Temporal Projection through Floristic Data. **2022**, 11, 1219 ○
- 162 Staying close to home: Marine habitat selection by foraging yellow-eyed penguins using spatial distribution models. 9,
- 161 Evaluation of Spatial Distribution of Three Major *Leptocorisa* (Hemiptera: Alydidae) Pests Using MaxEnt Model. **2022**, 13, 750 ○

- 160 Geography vs. past climate: the drivers of population genetic structure of the Himalayan langur. **2022**, 22, 0
- 159 Modeling the distribution of the endangered Jemez Mountains salamander (*Plethodon neomexicanus*) in relation to geology, topography, and climate. **2022**, 12, 0
- 158 Global invasibility potential of the shrub *Baccharis dracunculifolia*. 0
- 157 Incorporating interspecific relationships into species distribution models can better assess the response of species to climate change, a case study of two Chinese primates. **2022**, 142, 109255 0
- 156 Prediction of the dynamic distribution for *Eucheuma denticulatum* (Rhodophyta, Solieriaceae) under climate change in the Indo-Pacific Ocean. **2022**, 180, 105730 0
- 155 Going west: Range expansion for loggerhead sea turtles in the Mediterranean Sea under climate change. **2022**, 38, e02264 1
- 154 Analysis of the impact of climate change on the distribution and active compound content of the plateau medicinal plant *Nardostachys jatamansi* (D. Don) DC. **2022**, 187, 115438 0
- 153 Reconstructing the distribution of Chacoan biota from current and past evidence: the case of the southern three-banded armadillo *Tolypeutes matacus* (Desmarest, 1804). 0
- 152 Mapping habitats sensitive to overgrazing in the Swiss Northern Alps using habitat suitability modeling. **2022**, 274, 109742 0
- 151 A machine learning algorithm-based approach (MaxEnt) for predicting invasive potential of *Trioza erytreae* on a global scale. **2022**, 71, 101792 0
- 150 Beyond topo-climatic predictors: Does habitats distribution and remote sensing information improve predictions of species distribution models?. **2022**, 39, e02286 0
- 149 Mapping the changing distribution of two important pollinating giant honeybees across 21000 years. **2022**, 39, e02282 1
- 148 Impacts of climate change on the distributions and diversity of the giant panda with its sympatric mammalian species. **2022**, 144, 109452 1
- 147 Future changes in the seasonal habitat suitability for anchovy (*Engraulis japonicus*) in Korean waters projected by a maximum entropy model. 9, 0
- 146 A spatially explicit risk assessment of salamander populations to *Batrachochytrium* salamandrivorans in the United States. 0
- 145 Evaluation of the Likelihood of Establishing False Codling Moth (*Thaumatotibia leucotreta*) in Australia via the International Cut Flower Market. **2022**, 13, 883 0
- 144 Seasonal models reveal niche changes during invasion in *Capsella bursa-pastoris*. 0
- 143 Jack of all trades: genome assembly of Wild Jack and comparative genomics of *Artocarpus*. 0

- 142 The genomic landscapes of desert birds form over multiple time scales. o
- 141 Predicting invasion risk of rugose spiralling whitefly, *Aleurodicus rugioperculatus* in India based on CMIP6 projections by MaxEnt. o
- 140 Spotting what's important: Priority areas, connectivity, and conservation of the Northern Tiger Cat (*Leopardus tigrinus*) in Colombia. **2022**, 17, e0273750 o
- 139 Widespread exposure to Crimean-Congo haemorrhagic fever in Uganda might be driven by transmission from *Rhipicephalus* ticks: Evidence from cross-sectional and modelling studies. **2022**, o
- 138 An exhaustive evaluation of modeling ecological niches above species level to predict marine biological invasions. o
- 137 Predicting the Potential Distribution of Endangered *Parrotia subaequalis* in China. **2022**, 13, 1595 o
- 136 Influence of Pleistocene climate fluctuations on the demographic history and distribution of the critically endangered Chinese pangolin (*Manis pentadactyla*). **2022**, 7, o
- 135 Species distribution models and climatic niche comparisons provide clues on the geographic origin of a spider invasion in the Americas. o
- 134 Amphibian Diversity of the Yucatan Peninsula: Representation in Protected Areas and Climate Change Impacts. **2022**, 14, 813 o
- 133 Prediction of Suitable Distribution Area of Plateau pika (*Ochotona curzoniae*) in the Qinghai-Tibet Plateau under Shared Socioeconomic Pathways (SSPs). **2022**, 14, 12114 3
- 132 Ecological niche modelling as a tool to identify candidate indigenous chicken ecotypes of Tigray (Ethiopia). 13, o
- 131 Predicting the distribution of suitable habitat of the poisonous weed *Astragalus variabilis* in China under current and future climate conditions. 13, o
- 130 Comparing climatic suitability and niche distances to explain populations responses to extreme climatic events. o
- 129 The influence of road networks on brown bear spatial distribution and habitat suitability in a human-modified landscape. o
- 128 Reconstruction and variability of tropical pollination networks in the Brazilian Atlantic Forest. **2022**, 23, 315-325 o
- 127 Sumatra-wide assessment of spatiotemporal niche partitioning among small carnivore species. o
- 126 Spatial ensemble modeling for predicting the potential distribution of *Lymantria dispar asiatica* (Lepidoptera: Erebidae: Lymantriinae) in South Korea. **2022**, 194, o
- 125 Potential Global Distribution of Invasive Alien Species, *Anthonomus grandis* Boheman, under Current and Future Climate Using Optimal MaxEnt Model. **2022**, 12, 1759 1

- 124 Habitat Suitability of Fig (*Ficus carica* L.) in Mexico under Current and Future Climates. **2022**, 12, 1816 ○
- 123 Predicting suitable habitat for the endangered plant *Cephalotaxus oliveri* Mast. in China. 1-8 ○
- 122 Molecular, morphometric, and spatial data analyses provide new insights into the evolutionary history of the *Peromyscus boylii* species complex (Rodentia: Cricetidae) in the mountains of Mexico. **2022**, 20, 1-19 ○
- 121 Spatial Evaluation of Machine Learning-Based Species Distribution Models for Prediction of Invasive Ant Species Distribution. **2022**, 12, 10260 ○
- 120 Incorporating evolutionary and threat processes into crop wild relatives conservation. **2022**, 13, 1
- 119 Preferable and limiting conditions of trans-Palearctic Orthocephalus species (Heteroptera: Miridae). **2022**, 67, ○
- 118 Assessment of Climate Change and Land Use Effects on Water Lily (*Nymphaea* L.) Habitat Suitability in South America. **2022**, 14, 830 ○
- 117 Range expansion and establishment of a non-indigenous tunicate (*Diplosoma listerianum*) in thermal refugia is mediated by environmental variability in changing coastal environments.. ○
- 116 The possibilities of explicit *Striga* (*Striga asiatica*) risk monitoring using phenometric, edaphic, and climatic variables, demonstrated for Malawi and Zambia. **2022**, 194, ○
- 115 Testing the assumption of environmental equilibrium in an invasive plant species over a 130 year history. ○
- 114 Snow limits polecat *Mustela putorius* distribution in Sweden. ○
- 113 Realized niche shift of an invasive widow spider: drivers and impacts of human activities. **2022**, 19, ○
- 112 Linking environmental stability with genetic diversity and population structure in two Atlantic Forest palm trees. ○
- 111 Developing species distribution models for critically endangered species using participatory data: The European sturgeon marine habitat suitability. **2022**, 108136 ○
- 110 Using MaxEnt to Predict the Potential Distribution of the Little Fire Ant (*Wasmannia auropunctata*) in China. **2022**, 13, 1008 1
- 109 New genetic information and geographic distribution of charismatic carnivores: the olingos (Procyonidae: *Bassaricyon*) in Colombia. ○
- 108 Threatened species could be more vulnerable to climate change in tropical countries. **2023**, 858, 159989 3
- 107 Updated geographical distribution of the Sunbittern (<i>Eurypyga helias</i>; Eurypigidae). **2022**, 44, 612-618 ○

| | | |
|-----|--|---|
| 106 | Simulation the potential distribution of <i>Dendrolimus houi</i> and its hosts, <i>Pinus yunnanensis</i> and <i>Cryptomeria fortunei</i> , under climate change in China. 13, | o |
| 105 | Potential global geographical distribution of <i>Lolium temulentum</i> L. under climate change. 13, | o |
| 104 | Landscape-scale drivers of liana load across a Southeast Asian forest canopy differ to the Neotropics. | o |
| 103 | Together forever? Hummingbird-plant relationships in the face of climate warming. 2022 , 175, | o |
| 102 | Risk analysis of the spread of the quarantine pest mite <i>Schizotetranychus hindustanicus</i> in Brazil. | o |
| 101 | Identifying the Past, Present, and Future Distribution Patterns of the Balkan Wall Lizard (<i>Sauria: Lacertidae: Podarcis tauricus</i>) by Ecological Niche Modelling. 146-159 | o |
| 100 | Protected areas fail to cover the full annual cycle of migratory butterflies. | o |
| 99 | Minimal climate change impacts on the geographic distribution of <i>Nepeta glomerulosa</i> , medicinal species endemic to southwestern and central Asia. 2022 , 12, | o |
| 98 | Mapping the potential northern limits and promotion extent of ratoon rice in China. 2023 , 150, 102822 | 1 |
| 97 | A curated list of R packages for ecological niche modelling. 2023 , 476, 110242 | o |
| 96 | Rarity and conservation status of the Colombian Speckled Tree Rat, <i>Pattonomys semivillosus</i> (l. Geoffroy, 1838). 2022 , | o |
| 95 | Priorities for expanding the protected area system in Bangladesh. | o |
| 94 | <i>Cordyceps cicadae</i> and <i>Cordyceps gunnii</i> have closer species correlation with <i>Cordyceps sinensis</i> : from the perspective of metabonomic and MaxEnt models. 2022 , 12, | o |
| 93 | Modeling the geographical distributions of <i>Chordodes formosanus</i> and its mantis hosts in Taiwan, with considerations for their niche overlaps. 2022 , 12, | o |
| 92 | Back to the future: Climate change effects on habitat suitability of <i>Parnassius apollo</i> throughout the Quaternary glacial cycles. | o |
| 91 | Predicting Climate Change Effects on the Potential Distribution of Two Invasive Cryptic Species of the <i>Bemisia tabaci</i> Species Complex in China. 2022 , 13, 1081 | o |
| 90 | Predicting potential transmission risk of Everglades virus in Florida using mosquito blood meal identifications. 2, | o |
| 89 | Predicting the Potential Distribution of Pine Wilt Disease in China under Climate Change. 2022 , 13, 1147 | o |

- 88 Predicting the Impact of Climate Change on the Distribution of a Neglected Arboviruses Vector (*Armigeres subalbatus*) in China. **2022**, 7, 431 ○
- 87 Ecological determinants and risk areas of *Striga hermonthica* infestation in western Kenya under changing climate. ○
- 86 Effects of Climate Change and Environmental Factors on Bamboo (*Ferocalamus strictus*), a PSESP Unique to China. **2022**, 13, 2108 ○
- 85 Surveillance and invasive risk of the red imported fire ant, *Solenopsis invicta* Buren in China. ○
- 84 Range restricted old and young lineages show the southern Western Ghats to be both museum and cradle of diversity for woody plants. ○
- 83 Modeling of historical and current distributions of lone star tick, *Amblyomma americanum* (Acari: Ixodidae), is consistent with ancestral range recovery. 1
- 82 Habitat suitability and area of occupancy defined for rare New World sea snake. ○
- 81 Jack of all trades: Genome assembly of Wild Jack and comparative genomics of *Artocarpus*. 13, ○
- 80 Population development and landscape preference of reintroduced wild ungulates: successful rewilding in Southern Italy. 10, e14492 ○
- 79 Importance of data selection and filtering in species distribution models: A case study on the Cantabrian brown bear. **2022**, 13, 1
- 78 Estructura filogeográfica de *Crescentia alata* (Bignoniaceae): los huertos como reservorios de diversidad local. **2022**, 101, 164-185 ○
- 77 Gene flow assessment helps to distinguish strong genomic structure from speciation in an Iberian ant-eating spider. **2022**, 107682 ○
- 76 Genomic Consequences of and Demographic Response to Pervasive Hybridization over Time in Climate-sensitive Pikas. ○
- 75 Modeling the potential distribution of two immortality flora in the Philippines: Applying MaxEnt and GARP algorithms under different climate change scenarios. ○
- 74 Primates facing climate crisis in a tropical forest hotspot will lose climatic suitable geographical range. **2023**, 13, ○
- 73 The current and future seasonal geographic distribution of largehead hairtail *Trichiurus japonicus* in the Beibu Gulf, South China Sea. 9, ○
- 72 Using Species Distribution Models (SDMs) to Estimate the Suitability of European Mediterranean Non-Native Area for the Establishment of *Toumeyella Parvicornis* (Hemiptera: Coccidae). **2023**, 14, 46 ○
- 71 The Different Fate of the Pyrenean Desman (*Galemys pyrenaicus*) and the Eurasian Otter (*Lutra lutra*) under Climate and Land Use Changes. **2023**, 13, 274 ○

- 70 A review on the status and modeling of suitable habitats of the southern white-cheeked gibbon. ○
- 69 Mapping the phosphorus sorption capacity of Danish soils in four depths with quantile regression forests and uncertainty propagation. **2023**, 430, 116316 ○
- 68 Modeling climate change impacts on potential global distribution of *Tamarixia radiata* Waterston (Hymenoptera: Eulophidae). **2023**, 864, 160962 ○
- 67 Phylogeography within the *Peromyscus maniculatus* species group: Understanding past distribution of genetic diversity and areas of refugia in western North America. **2023**, 180, 107701 ○
- 66 Climate influences the genetic structure and niche differentiation among populations of the olive field mouse *Abrothrix olivacea* (Cricetidae: Abrotrichini). **2022**, 12, 1
- 65 Transmission Risk Predicting for Schistosomiasis in Mainland China by Exploring Ensemble Ecological Niche Modeling. **2023**, 8, 24 ○
- 64 Predicting the geographical distribution of *Acropora muricata* in two lesser-known reef systems of the Andaman Sea. **2022**, 26, ○
- 63 Rarely naturalized, but widespread and even invasive: the paradox of a popular pet terrapin expansion in Eurasia. 81, 91-127 ○
- 62 Ecological-niche modeling reveals current opportunities for Agave dryland farming in Sonora, Mexico and Arizona, USA. **2023**, 18, e0279877 ○
- 61 The benefits of being smaller: Consistent pattern for climate-induced range shift and morphological difference of three falconiforme species. **2023**, 14, 100079 ○
- 60 Modelling distribution and fate of coralligenous habitat in the Northern Adriatic Sea under a severe climate change scenario. 10, ○
- 59 Maximum Entropy Modeling of Giant Pangolin *Smutsia gigantea* (Illiger, 1815) habitat suitability in a protected forest-savannah transition area of Central Cameroon. **2023**, e02395 ○
- 58 The impact of climate change and human activities over the past 2000 years has increased the spatial-temporal extinction rate of gibbons. **2023**, 281, 109998 ○
- 57 Where to go? Habitat preferences and connectivity at a crossroad of European brown bear metapopulations. **2023**, 43, e02460 ○
- 56 Climate change impacts on the availability of anti-malarial plants in Kenya. **2023**, 5, 100070 ○
- 55 A new species of small-eared shrew (Soricidae, *Cryptotis*) from El Triunfo Biosphere Reserve, Chiapas, Mexico. ○
- 54 Microhabitat modeling of the invasive Asian longhorned tick (*Haemaphysalis longicornis*) in New Jersey, USA. **2023**, 14, 102126 ○
- 53 The identification and conservation of climate refugia for two Colombian endemic titi (*Plecturocebus*) monkeys. **2023**, 72, 126345 ○

- 52 Flexible species distribution modelling methods perform well on spatially separated testing data. **2023**, 32, 369-383 ○
- 51 The TOP-100 most dangerous invasive alien species in Northern Eurasia: invasion trends and species distribution modelling. **2023**, 82, 23-56 ○
- 50 Assessing risk from invasive alien plants in China: Reconstructing invasion history and estimating distribution patterns of *Lolium temulentum* and *Aegilops tauschii*. **2023**, 14, ○
- 49 Species distribution modelling of *Monothea buxifolia* (Falc.) A. DC.: Present distribution and impacts of potential climate change. **2023**, 9, e13417 1
- 48 Habitat and ecological niche characteristics of the elusive Hairy-eared Dwarf Lemur (*Allocebus trichotis*) with updated occurrence and geographic range data. **2023**, 85, ○
- 47 Effects of climate change and anthropogenic activity on ranges of vertebrate species endemic to the Qinghai-Tibet Plateau over 40 years. ○
- 46 Contrasting ecological information content in whaling archives with modern cetacean surveys for conservation planning and identification of historical distribution changes. ○
- 45 Evaluating niche changes during invasion with seasonal models in *Capsella bursa-pastoris*. **2023**, 110, ○
- 44 Predicting current and future potential distribution of *Cynaesus angustus* (Coleoptera: Tenebrionidae) in global scale using the MaxEnt model. **2023**, 101, 102089 ○
- 43 Prey resources are equally important as climatic conditions for predicting the distribution of a broad-ranged apex predator. **2023**, 29, 613-628 ○
- 42 Integration matters: Combining socio-cultural and biophysical methods for mapping ecosystem service bundles. ○
- 41 Potentially Suitable Geographical Area for *Monochamus alternatus* under Current and Future Climatic Scenarios Based on Optimized MaxEnt Model. **2023**, 14, 182 ○
- 40 MaxEnt Modeling for Predicting Suitable Habitat for Endangered Tree *Keteleeria davidiana* (Pinaceae) in China. **2023**, 14, 394 ○
- 39 Areography, environmental heterogeneity and spatial models explain patterns of past and present diversity in *Edraianthus* (Campanulaceae). ○
- 38 Projecting environmental suitability areas for the seaweed *Gracilaria birdiae* (Rhodophyta) in Brazil: Implications for the aquaculture pertaining to five environmentally crucial parameters. **2023**, 35, 773-784 ○
- 37 MaxEnt brings comparable results when the input data are being completed; Model parameterization of four species distribution models. **2023**, 13, ○
- 36 Habitat Suitability Evaluation of Different Forest Species in Lvliang Mountain by Combining Prior Knowledge and MaxEnt Model. **2023**, 14, 438 ○
- 35 An exhaustive evaluation of modeling ecological niches above species level to predict marine biological invasions. **2023**, 186, 105926 ○

- 34 Current and future distribution of a parasite with complex life cycle under global change scenarios: *Echinococcus multilocularis* in Europe. **2023**, 29, 2436-2449 ○
- 33 Multi-temporal ecological niche modeling for bird conservation in the face of climate change scenarios in Caatinga, Brazil. 11, e14882 ○
- 32 You're stressing me out! Effect of interspecific competition from red deer on roe deer physiological stress response. ○
- 31 Extensive protected area coverage and an updated global population estimate for the Endangered Madagascar Serpent-eagle *Eutriorchis astur*. **2023**, 33, ○
- 30 Dual domestications and origin of traits in grapevine evolution. **2023**, 379, 892-901 1
- 29 Climatic niche shift and distribution of *Melanagromyza sojae* under current and future climate scenarios: does this species pose a risk to soybean production?. ○
- 28 Prediction of the potentially suitable areas of *Litsea cubeba* in China based on future climate change using the optimized MaxEnt model. **2023**, 148, 110093 ○
- 27 Projected Shifts in Bird Distribution in India under Climate Change. **2023**, 15, 404 ○
- 26 Current and Potential Future Global Distribution of the Raisin Moth *Cadra figulilella* (Lepidoptera: Pyralidae) under Two Different Climate Change Scenarios. **2023**, 12, 435 ○
- 25 Geometric morphometrics and ecological niche modelling for delimitation of *Triatoma pallidipennis* (Hemiptera: Reduviidae: Triatominae) haplogroups. **2023**, 3, 100119 ○
- 24 An Invisible Boundary between Geographic Ranges of Cryptic Species of Narrow-Headed Voles (*Stenocranius*, *Lasiopodomys*, Cricetidae) in Transbaikalia. **2023**, 15, 439 ○
- 23 Assessing environmental suitability of *Ligusticum chuanxiong* based on ecological analyses with chemical and molecular verification. **2023**, 9, e14629 ○
- 22 Molecular data, ecological niche, and dispersal models reveal a trans-Atlantic shallow-water octopus species. **2023**, 213, 103019 ○
- 21 The evolution of white-tailed jackrabbit camouflage in response to past and future seasonal climates. **2023**, 379, 1238-1242 ○
- 20 Prediction of global potential suitable habitats of *Nicotiana glauca* Link et Otto based on MaxEnt model. **2023**, 13, ○
- 19 Climatic envelopes of the genus *Lacerta* Linnaeus, 1758 in Türkiye: an application of ecological niche modeling. **2023**, 30, 56382-56397 ○
- 18 Habitat connectivity supports the local abundance of fire salamanders (*Salamandra atra*) but also the spread of *Batrachochytrium salamandrivorans*. ○
- 17 Ecological Niche Modelling Approaches: Challenges and Applications in Vector-Borne Diseases. **2023**, 8, 187 ○

- 16 Desert Locust (*Schistocerca gregaria*) Invasion Risk and Vegetation Damage in a Key Upsurge Area. **2023**, 4, 187-208 ○
- 15 Effects of Environmental Variation in Structuring Population Genetic Variation in the False-Water Cobras (Xenodontinae: Hydrodynastes). ○
- 14 The Fate of *Guzmania monostachia* in Florida Rests with Humans. **2023**, 15, 525 ○
- 13 N-SDM : a high-performance computing pipeline for Nested Species Distribution Modelling. ○
- 12 Priorities for conserving the world's terrestrial mammals based on over-the-horizon extinction risk. **2023**, 33, 1381-1388.e6 ○
- 11 Neglected tropical diseases risk correlates with poverty and early ecosystem destruction. **2023**, 12, ○
- 10 The habitat-suitability models of the European mole cricket (*Gryllotalpa gryllotalpa*) as information tool for conservation and pest management. **2023**, 9, e14826 ○
- 9 Climate change and the potential distribution of the invasive shrub, *Leucaena leucocephala* (Lam.) De Wit in Africa. ○
- 8 Spatial phylogenetics of Fagales: Investigating the history of temperate forests. ○
- 7 The ecological response of commercial fishes and shrimps to climate change: predicting global distributional shifts under future scenarios. **2023**, 23, ○
- 6 Effects of Climate Change on the Habitat Suitability and Distribution of Endemic Freshwater Fish Species in Semi-Arid Central Anatolian Ecoregion in Türkiye. **2023**, 15, 1619 ○
- 5 Range restricted old and young lineages show the southern Western Ghats to be both a museum and a cradle of diversity for woody plants. **2023**, 290, ○
- 4 Transmission risk of Oropouche fever across the Americas. **2023**, 12, ○
- 3 Contribution of the public to the modelling of the distributions of species: Occurrence and current and potential distribution of the ant *Manica rubida* (Hymenoptera: Formicidae). 120, 137-148 ○
- 2 Predicting Climate Change Impacts on Candelilla (*Euphorbia antisiphilitica* Zucc.) for Mexico: An Approach for Mexico's Primary Harvest Area. **2023**, 15, 7737 ○
- 1 Multitemporal relative landslide exposure and risk analysis for the sustainable development of rapidly growing cities. ○