
## CITATION REPORT List of articles citing



DOI: 10.1002/14651858.cd009124.pub2 Cochrane Database of Systematic Reviews, 2015, , CD009124.

Source: https://exaly.com/paper-pdf/60572221/citation-report.pdf

Version: 2024-04-25

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
25	Effect of the type of maternal pushing during the second stage of labour on obstetric and neonatal outcome: a multicentre randomised trial-the EOLE study protocol. <i>BMJ Open</i> , <b>2016</b> , 6, e012290	3	4
24	Methods of pushing during vaginal delivery and pelvic floor and perineal outcomes: a review. <i>Current Opinion in Obstetrics and Gynecology</i> , <b>2016</b> , 28, 470-476	2.4	9
23	Prevention of pelvic floor disorders: international urogynecological association research and development committee opinion. <i>International Urogynecology Journal</i> , <b>2016</b> , 27, 1785-1795	2	23
22	Committee Opinion No. 687: Approaches to Limit Intervention During Labor and Birth. <i>Obstetrics and Gynecology</i> , <b>2017</b> , 129, e20-e28	4.9	76
21	Perineal techniques during the second stage of labour for reducing perineal trauma. <i>The Cochrane Library</i> , <b>2017</b> , 6, CD006672	5.2	56
20	Pushing/bearing down methods for the second stage of labour. <i>The Cochrane Library</i> , <b>2017</b> , 3, CD00912	245.2	13
19	Durations of second stage of labor and pushing, and adverse neonatal outcomes: a population-based cohort study. <i>Journal of Perinatology</i> , <b>2017</b> , 37, 236-242	3.1	41
18	Current Resources for Evidence-Based Practice, May/June 2017. <i>Journal of Midwifery and Womens Health</i> , <b>2017</b> , 62, 373-379	2.2	
17	Labour and normal birth. <b>2017</b> , 1-48		
16	MidwivesWManagement during the Second Stage of Labor in Relation to Second-Degree Tears-An Experimental Study. <i>Birth</i> , <b>2017</b> , 44, 86-94	3.1	31
15	Effectiveness of breathing exercises during the second stage of labor on labor pain and duration: a randomized controlled trial. <i>Journal of Integrative Medicine</i> , <b>2017</b> , 15, 456-461	4	19
14	Practices used by midwives during the second stage of labor to facilitate birth - Are they related to perineal trauma?. <i>Sexual and Reproductive Healthcare</i> , <b>2018</b> , 15, 18-22	2.4	1
13	Factors associated with obstetric anal sphincter injuries in midwife-led birth: A cross sectional study. <i>Midwifery</i> , <b>2018</b> , 62, 264-272	2.8	11
12	Pilates program design and health benefits for pregnant women: Alpractitioners Waurvey. <i>Journal of Bodywork and Movement Therapies</i> , <b>2018</b> , 22, 411-417	1.6	7
11	Methods for spontaneous delivery. <b>2018</b> , 569-581		
10	Implementation of evidence-based practices in normal delivery care. <i>Revista Latino-Americana De Enfermagem</i> , <b>2018</b> , 26, e2988	1.5	7
9	Associations between improved care during the second stage of labour and maternal and neonatal health outcomes in a rural hospital in Bangladesh. <i>Midwifery</i> , <b>2018</b> , 66, 30-35	2.8	2

## CITATION REPORT

8	Intact Perineum: What are the Predictive Factors in Spontaneous Vaginal Birth?. <i>Materia Socio-medica</i> , <b>2019</b> , 31, 25-30	0.9	2
7	Factors related to episiotomy practice: an evidence-based medicine systematic review. <i>Journal of Obstetrics and Gynaecology</i> , <b>2019</b> , 39, 737-747	1.3	7
6	A protocol for developing, disseminating, and implementing a core outcome set (COS) for childbirth pelvic floor trauma research. <i>BMC Pregnancy and Childbirth</i> , <b>2020</b> , 20, 376	3.2	10
5	The Role of Intrapartum Sonography in Prolonged Labor. <b>2021</b> , 155-167		
4	Effect of Breathing Technique of Blowing on the Extent of Damage to the Perineum at the Moment of Delivery: A Randomized Clinical Trial. <i>Iranian Journal of Nursing and Midwifery Research</i> , <b>2017</b> , 22, 62-6	163	11
3	The effects of ropivacaine 0.0625% and levobupivacaine 0.0625% on uterine and abdominal muscle electromyographic activity during the second stage of labor. <i>Minerva Anestesiologica</i> , <b>2019</b> , 85, 854-861	1.9	1
2	Clinical evaluation of labor: an evidence- and experience-based approach. <i>Journal of Perinatal Medicine</i> , <b>2021</b> , 49, 241-253	2.7	0
1	Management of the active second stage of labor in waterbirths compared with conventional births - a prospective cohort study <i>Midwifery</i> , <b>2022</b> , 107, 103283	2.8	