## QTL mapping in multiple populations and development trait loci for fruit size in cucumbers of different market

Theoretical and Applied Genetics 128, 1747-1763 DOI: 10.1007/s00122-015-2544-7

**Citation Report** 

| #  | ARTICLE  | IF              | CITATIONS    |
|----|--|-----------------|--------------|
| 1  | A High-Density Genetic Linkage Map for Cucumber (Cucumis sativus L.): Based on Specific Length<br>Amplified Fragment (SLAF) Sequencing and QTL Analysis of Fruit Traits in Cucumber. Frontiers in Plant<br>Science, 2016, 7, 437.    | 3.6             | 120          |
| 2  | Genetic Mapping of Complex Traits in Cucurbits. Plant Genetics and Genomics: Crops and Models, 2016, , 269-290.  | 0.3             | 3            |
| 3  | Genomic Analysis of Cucurbit Fruit Growth. Plant Genetics and Genomics: Crops and Models, 2016, , 321-344.   | 0.3             | 7            |
| 4  | SHORT HYPOCOTYL 1 Encodes a SMARCA3-like Chromatin Remodeling Factor Regulating Elongation.<br>Plant Physiology, 2016, 172, pp.00501.2016.   | 4.8             | 22           |
| 5  | QTL mapping of parthenocarpic fruit set in North American processing cucumber. Theoretical and Applied Genetics, 2016, 129, 2387-2401.   | 3.6             | 42           |
| 6  | Rapid identification of fruit length loci in cucumber (Cucumis sativus L.) using next-generation sequencing (NGS)-based QTL analysis. Scientific Reports, 2016, 6, 27496.  | 3.3             | 63           |
| 7  | Integrated analysis in bi-parental and natural populations reveals CsCLAVATA3 (CsCLV3) underlying carpel number variations in cucumber. Theoretical and Applied Genetics, 2016, 129, 1007-1022.                                      | 3.6             | 43           |
| 8  | Variation in cucumber (Cucumis sativus L.) fruit size and shape results from multiple components acting pre-anthesis and post-pollination. Planta, 2017, 246, 641-658.   | 3.2             | 53           |
| 9  | Transcriptomic analysis of short-fruit 1 (sf1) reveals new insights into the variation of fruit-related traits in Cucumis sativus. Scientific Reports, 2017, 7, 2950.  | 3.3             | 26           |
| 10 | QTL mapping of domestication and diversifying selection related traits in round-fruited semi-wild<br>Xishuangbanna cucumber (Cucumis sativus L. var. xishuangbannanesis). Theoretical and Applied<br>Genetics, 2017, 130, 1531-1548. | 3.6             | 60           |
| 11 | Quantitative trait locus analysis of cucumber fruit morphological traits based on image analysis.<br>Euphytica, 2017, 213, 1.  | 1.2             | 15           |
| 12 | Round fruit shape in WI7239 cucumber is controlled by two interacting quantitative trait loci with one putatively encoding a tomato SUN homolog. Theoretical and Applied Genetics, 2017, 130, 573-586.                               | 3.6             | 80           |
| 13 | Genetics and Genomics of Cucurbitaceae. Plant Genetics and Genomics: Crops and Models, 2017, , .   | 0.3             | 22           |
| 14 | A high-density linkage map and QTL mapping of fruit-related traits in pumpkin (Cucurbita moschata) Tj ETQq1 1  | 0.784314<br>3.3 | rgBT /Overic |
| 15 | STAYGREEN (CsSGR) is a candidate for the anthracnose (Colletotrichum orbiculare) resistance locus cla in Gy14 cucumber. Theoretical and Applied Genetics, 2018, 131, 1577-1587.  | 3.6             | 21           |
| 16 | Genetic mapping reveals a candidate gene (CIFS1) for fruit shape in watermelon (Citrullus lanatus L.).<br>Theoretical and Applied Genetics, 2018, 131, 947-958.  | 3.6             | 94           |
| 17 | Dynamic and epistatic QTL mapping reveals the complex genetic architecture of waterlogging tolerance in chrysanthemum. Planta, 2018, 247, 899-924.   | 3.2             | 21           |
| 18 | QTL Analysis for Downy Mildew Resistance in Cucumber Inbred Line PI 197088. Plant Disease, 2018, 102, 1240-1245.   | 1.4             | 33           |

| #  | Article   | IF  | CITATIONS |
|----|---|-----|-----------|
| 19 | Genetic variation of dynamic fiber elongation and developmental quantitative trait locus mapping of fiber length in upland cotton (Gossypium hirsutum L.). BMC Genomics, 2018, 19, 882.   | 2.8 | 33        |
| 20 | Identification of Conserved and Diverse Metabolic Shift of the Stylar, Intermediate and Peduncular<br>Segments of Cucumber Fruit during Development. International Journal of Molecular Sciences, 2018,<br>19, 135.                     | 4.1 | 14        |
| 21 | <i>LITTLELEAF</i> ( <i>LL</i> ) encodes a WD40 repeat domain ontaining protein associated with organ size variation in cucumber. Plant Journal, 2018, 95, 834-847.  | 5.7 | 58        |
| 22 | Identification of <scp>QTL</scp> s controlling lowâ€ŧemperature tolerance during the germination stage in cucumber ( <i>Cucumis sativus</i> L.). Plant Breeding, 2018, 137, 629-637.  | 1.9 | 16        |
| 23 | STAYGREEN, STAY HEALTHY: a lossâ€ofâ€susceptibility mutation in the <i>STAYGREEN</i> gene provides<br>durable, broadâ€spectrum disease resistances for over 50Âyears of US cucumber production. New<br>Phytologist, 2019, 221, 415-430. | 7.3 | 72        |
| 24 | Quantitative trait loci analysis of hormone levels in Arabidopsis roots. PLoS ONE, 2019, 14, e0219008.  | 2.5 | 3         |
| 25 | Mapping quantitative trait loci for yield-related traits and predicting candidate genes for grain weight in maize. Scientific Reports, 2019, 9, 16112.  | 3.3 | 15        |
| 26 | QTL mapping and genome-wide association study reveal two novel loci associated with green flesh<br>color in cucumber. BMC Plant Biology, 2019, 19, 243.   | 3.6 | 35        |
| 27 | Target SSR-Seq: A Novel SSR Genotyping Technology Associate With Perfect SSRs in Genetic Analysis of<br>Cucumber Varieties. Frontiers in Plant Science, 2019, 10, 531.  | 3.6 | 37        |
| 28 | Genetic Regulation of Ethylene Dosage for Cucumber Fruit Elongation. Plant Cell, 2019, 31, 1063-1076.   | 6.6 | 85        |
| 29 | A Functional Allele of <i>CsFUL1</i> Regulates Fruit Length through Repressing <i>CsSUP</i> and Inhibiting Auxin Transport in Cucumber. Plant Cell, 2019, 31, 1289-1307.  | 6.6 | 84        |
| 30 | QTL mapping of low temperature germination ability in cucumber. Euphytica, 2019, 215, 1.  | 1.2 | 17        |
| 31 | Genetic analysis and QTL mapping of fruit length and diameter in a cucumber (Cucumber sativus L.) recombinant inbred line (RIL) population. Scientia Horticulturae, 2019, 250, 214-222.   | 3.6 | 13        |
| 32 | QTL mapping for 11 agronomic traits based on a genome-wide Bin-map in a large F2 population of foxtail<br>millet (Setaria italica (L.) P. Beauv). Molecular Breeding, 2019, 39, 1.  | 2.1 | 24        |
| 33 | Quantitative Trait Loci Mapping and Candidate Gene Analysis of Low Temperature Tolerance in<br>Cucumber Seedlings. Frontiers in Plant Science, 2019, 10, 1620.  | 3.6 | 19        |
| 34 | QTL and Transcriptomic Analyses Implicate Cuticle Transcription Factor SHINE as a Source of Natural<br>Variation for Epidermal Traits in Cucumber Fruit. Frontiers in Plant Science, 2019, 10, 1536.                                    | 3.6 | 12        |
| 35 | Novel loci fsd6.1 and Csgl3 regulate ultra-high fruit spine density in cucumber. Theoretical and Applied Genetics, 2019, 132, 27-40.  | 3.6 | 23        |
| 36 | Molecular basis of cucumber fruit domestication. Current Opinion in Plant Biology, 2019, 47, 38-46.   | 7.1 | 96        |

CITATION REPORT

\_

| #  | Article  | IF  | CITATIONS |
|----|--|-----|-----------|
| 37 | Origin and domestication of Cucurbitaceae crops: insights from phylogenies, genomics and archaeology. New Phytologist, 2020, 226, 1240-1255.   | 7.3 | 134       |
| 38 | Quantitative trait loci for fruit size and flowering timeâ€related traits under domestication and diversifying selection in cucumber ( <i>Cucumis sativus</i> ). Plant Breeding, 2020, 139, 176-191.             | 1.9 | 26        |
| 39 | Genome-wide Target Mapping Shows Histone Deacetylase Complex1 Regulates Cell Proliferation in<br>Cucumber Fruit. Plant Physiology, 2020, 182, 167-184.   | 4.8 | 47        |
| 40 | Map-based cloning of qBWT-c12 discovered brassinosteroid-mediated control of organ size in cotton.<br>Plant Science, 2020, 291, 110315.  | 3.6 | 4         |
| 41 | The majorâ€effect quantitative trait locus Fnl7.1 encodes a late embryogenesis abundant protein associated with fruit neck length in cucumber. Plant Biotechnology Journal, 2020, 18, 1598-1609.                 | 8.3 | 14        |
| 42 | Genetic architecture of fruit size and shape variation in cucurbits: a comparative perspective.<br>Theoretical and Applied Genetics, 2020, 133, 1-21.  | 3.6 | 111       |
| 43 | Identification of fruit size associated quantitative trait loci featuring SLAF based high-density linkage<br>map of goji berry (Lycium spp.). BMC Plant Biology, 2020, 20, 474.                                  | 3.6 | 18        |
| 44 | A durum wheat adult plant stripe rust resistance QTL and its relationship with the bread wheat Yr80<br>locus. Theoretical and Applied Genetics, 2020, 133, 3049-3066.  | 3.6 | 10        |
| 45 | Dynamic Changes of Fruit Shape Traits and Its Correlation to the Morphology of Other Organs in<br>Cucumber. Agronomy, 2020, 10, 1111.  | 3.0 | 2         |
| 46 | Identification of QTLs for grain yield and other traits in tropical maize under Striga infestation. PLoS<br>ONE, 2020, 15, e0239205.   | 2.5 | 14        |
| 47 | QTL Mapping for Cucumber Fruit Size and Shape with Populations from Long and Round Fruited Inbred<br>Lines. Horticultural Plant Journal, 2020, 6, 132-144.   | 5.0 | 20        |
| 48 | Detection of putative QTL regions associated with ovary traits in melon using SNP-CAPS markers.<br>Scientia Horticulturae, 2020, 270, 109445.  | 3.6 | 14        |
| 50 | Gene regulatory network of carpel number variation in cucumber. Development (Cambridge), 2020, 147,  | 2.5 | 16        |
| 51 | The MADS-Box Gene CsSHP Participates in Fruit Maturation and Floral Organ Development in Cucumber. Frontiers in Plant Science, 2019, 10, 1781.   | 3.6 | 11        |
| 52 | Cucumber Fruit Size and Shape Variations Explored from the Aspects of Morphology, Histology, and<br>Endogenous Hormones. Plants, 2020, 9, 772.   | 3.5 | 30        |
| 53 | Localization of quantitative trait loci for cucumber fruit shape by a population of chromosome segment substitution lines. Scientific Reports, 2020, 10, 11030.  | 3.3 | 11        |
| 54 | Markers for selection of three alleles of ClSUN25-26-27a (Cla011257) associated with fruit shape in watermelon. Molecular Breeding, 2020, 40, 1.   | 2.1 | 12        |
| 55 | QTL for horticulturally important traits associated with pleiotropic andromonoecy and carpel<br>number loci, and a paracentric inversion in cucumber. Theoretical and Applied Genetics, 2020, 133,<br>2271-2290. | 3.6 | 8         |

| #  | Article   | IF               | CITATIONS           |
|----|---|------------------|---------------------|
| 56 | Quantitative trait loci for horticulturally important traits defining the Sikkim cucumber, Cucumis sativus var. sikkimensis. Theoretical and Applied Genetics, 2021, 134, 229-247.  | 3.6              | 19                  |
| 58 | Genetic linkage mapping and QTLs identification for morphology and fruit quality related traits of melon by SNP based CAPS markers. Scientia Horticulturae, 2021, 278, 109849.  | 3.6              | 18                  |
| 59 | Identification of quantitative trait loci for powdery mildew resistance in highly resistant cucumber<br>( <i>Cucumis sativus</i> L.) using ddRAD-seq analysis. Breeding Science, 2021, 71, 326-333.                                 | 1.9              | 2                   |
| 60 | Quantitative trait locus analysis of cucumber fruit texture using doubleâ€digest<br>restrictionâ€siteâ€associated DNA sequencing. Euphytica, 2021, 217, 1.  | 1.2              | 2                   |
| 61 | CsKTN1 for a katanin p60 subunit is associated with the regulation of fruit elongation in cucumber (Cucumis sativus L.). Theoretical and Applied Genetics, 2021, 134, 2429-2441.  | 3.6              | 15                  |
| 62 | Molecular research progress and improvement approach of fruit quality traits in cucumber.<br>Theoretical and Applied Genetics, 2021, 134, 3535-3552.  | 3.6              | 18                  |
| 63 | Recent advances in cucumber ( <i>Cucumis sativus</i> L.). Journal of Horticultural Science and<br>Biotechnology, 2022, 97, 3-23.  | 1.9              | 6                   |
| 64 | Longâ€read genome assembly and genetic architecture of fruit shape in the bottle gourd. Plant Journal, 2021, 107, 956-968.  | 5.7              | 23                  |
| 65 | Genomic Prediction and the Practical Breeding of 12 Quantitative-Inherited Traits in Cucumber (Cucumis sativus L.). Frontiers in Plant Science, 2021, 12, 729328.   | 3.6              | 2                   |
| 66 | Fine mapping and identification of the candidate gene BFS for fruit shape in wax gourd (Benincasa) Tj ETQq1 1 0   | .784314 r<br>3.6 | gBT /Overloci<br>16 |
| 67 | A Genotyping-by-sequencing Single Nucleotide Polymorphism–based Map and Genetic Analysis of Root<br>Traits in an Interspecific Tomato Population. Journal of the American Society for Horticultural<br>Science, 2019, 144, 394-404. | 1.0              | 4                   |
| 68 | Classical Genetics and Traditional Breeding in Cucumber (Cucumis sativus L.). , 0, , .  |                  | 1                   |
| 69 | Genome-wide analysis of expansins and their role in fruit spine development in cucumber (Cucumis) Tj ETQq0 0 (  | ) rgBT /Ov       | erlock 10 Tf 5      |
| 70 | <i>CmFSI8</i> / <i>CmOFP13</i> encoding an OVATE family protein controls fruit shape in melon.<br>Journal of Experimental Botany, 2022, 73, 1370-1384.  | 4.8              | 17                  |
| 71 | Promoter variation in a homeobox gene, CpDll, is associated with deeply lobed leaf in Cucurbita pepo<br>L Theoretical and Applied Genetics, 2022, 135, 1223-1234.   | 3.6              | 5                   |
| 72 | Genetic and Molecular Regulation Mechanisms in the Formation and Development of Vegetable Fruit<br>Shape. Applied Sciences (Switzerland), 2022, 12, 1514.   | 2.5              | 4                   |
| 73 | Mapping of genetic loci controlling fruit linked morphological traits of melon using developed CAPS markers. Molecular Biology Reports, 2022, 49, 5459-5472.  | 2.3              | 9                   |
| 74 | Genome-wide analysis of OVATE family proteins in cucumber (Cucumissativus L.). Journal of Integrative   | 3.5              | 5                   |

CITATION REPORT

## # ARTICLE

76 Mapping of fruit apex shape related QTLs across multi-genetic backgrounds in cucumber (Cucumis) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 5

| 100 | Functional copy number variation of CsSHINE1 is associated with fruit skin netting intensity in cucumber, Cucumis sativus. Theoretical and Applied Genetics, 2022, 135, 2101-2119.      | 3.6 | 5 |
|-----|---|-----|---|
| 101 | Identification of QTLs linked with watermelon fruit and seed traits using GBS-based high-resolution genetic mapping. Scientia Horticulturae, 2022, 303, 111237.                         | 3.6 | 8 |
| 102 | Serial-Omics and Molecular Function Study Provide Novel Insight into Cucumber Variety Improvement.<br>Plants, 2022, 11, 1609.   | 3.5 | 2 |
| 103 | QTL mapping for pumpkin fruit traits using a GBS-based high-density genetic map. Euphytica, 2022, 218, .  | 1.2 | 2 |
| 104 | Quantitative trait locus mapping of fruit aroma compounds in cucumber ( <i>Cucumber sativus</i> L.)<br>based on a recombinant inbred line population. Horticulture Research, 2022, 9, . | 6.3 | 2 |

Mapping and Validation of a New Quantitative Trait Loci (Qtl) for Fruit Size in Watermelon (Citrullus) Tj ETQq0 0 0 rgBT /Overlock 10 Tf

| 106 | Research Progress on the Leaf Morphology, Fruit Development and Plant Architecture of the Cucumber. Plants, 2022, 11, 2128.  | 3.5 | 3  |
|-----|--|-----|----|
| 107 | Identification of QTLs controlling cold tolerance in cucumber (Cucumis sativus L.) seedlings. Scientia<br>Horticulturae, 2022, 305, 111383.  | 3.6 | 2  |
| 108 | QTL analysis of early flowering of female flowers in zucchini (Cucurbita pepo L.). Journal of<br>Integrative Agriculture, 2023, 22, 3321-3330.   | 3.5 | 3  |
| 109 | Genomic Designing for Abiotic Stress Tolerance in Cucurbits. , 2022, , 187-252.  |     | 0  |
| 110 | Sigma factor binding protein 1 (CsSIB1) is a putative candidate of the major-effect QTL dm5.3 for downy mildew resistance in cucumber (Cucumis sativus). Theoretical and Applied Genetics, 2022, 135, 4197-4215.                                   | 3.6 | 4  |
| 111 | The CsHEC1-CsOVATE module contributes to fruit neck length variation via modulating auxin<br>biosynthesis in cucumber. Proceedings of the National Academy of Sciences of the United States of<br>America, 2022, 119, .                            | 7.1 | 10 |
| 112 | Ectopic Expression of CsSUN in Tomato Results in Elongated Fruit Shape via Regulation of Longitudinal Cell Division. International Journal of Molecular Sciences, 2022, 23, 9973.  | 4.1 | 1  |
| 113 | Phenotypic Characterization and Fine Mapping of a Major-Effect Fruit Shape QTL FS5.2 in Cucumber,<br>Cucumis sativus L., with Near-Isogenic Line-Derived Segregating Populations. International Journal of<br>Molecular Sciences, 2022, 23, 13384. | 4.1 | 2  |
| 114 | Relationship between Key Environmental Factors and the Architecture of Fruit Shape and Size in<br>Near-Isogenic Lines of Cucumber (Cucumis sativus L.). International Journal of Molecular Sciences,<br>2022, 23, 14033.                           | 4.1 | 7  |
| 115 | Natural variation in CRABS CLAW contributes to fruit length divergence in cucumber. Plant Cell, 2023, 35, 738-755.   | 6.6 | 11 |
| 116 | Morphological and Genetic Diversity of Cucumber (Cucumis sativus L.) Fruit Development. Plants, 2023, 12, 23.  | 3.5 | 3  |

| #   | Article  | IF              | CITATIONS  |
|-----|--|-----------------|------------|
| 117 | Primary mapping of quantitative trait loci regulating multivariate horticultural phenotypes of watermelon (Citrullus lanatus L.). Frontiers in Plant Science, 0, 13, .   | 3.6             | 3          |
| 118 | <i>CsTRM5</i> regulates fruit shape via mediating cell division direction and cell expansion in cucumber. Horticulture Research, 2023, 10, .   | 6.3             | 9          |
| 119 | Genetic Mapping and QTL Analysis of Fruit Traits in Melon (Cucumis melo L.). Current Issues in<br>Molecular Biology, 2023, 45, 3419-3433.  | 2.4             | 3          |
| 120 | Mapping and validation of a new quantitative trait locus (QTL) for fruit size in watermelon (Citrullus) Tj ETQq1 1 (   | ).784314<br>3.6 | rgBT /Over |
| 121 | Molecular and genetic regulations of fleshy fruit shape and lessons from Arabidopsis and rice.<br>Horticulture Research, 0, , .  | 6.3             | 2          |
| 122 | Transcriptional and phytohormone regulatory network involved in LITTLELEAF-mediated organ size development in cucumber (Cucumis sativus). Scientia Horticulturae, 2023, 321, 112294.                                       | 3.6             | 0          |
| 123 | QTL Mapping for Ovary- and Fruit-Related Traits in Cucumis sativus-C. hystrix Introgression Line IL52.<br>Genes, 2023, 14, 1133.   | 2.4             | 1          |
| 124 | Fine genetic mapping of the Mottled Rind Color (Morc) locus reveals a 4895-bp presence-absence variation contributing to the mottled or unmottled fruit rind color in cucumber. Scientia Horticulturae, 2023, 321, 112303. | 3.6             | 0          |
| 125 | QTL mapping and stability analysis of trichome density in zucchini (Cucurbita pepo L.). Frontiers in<br>Plant Science, 0, 14, .  | 3.6             | 1          |
| 126 | QTL mapping and transcriptomic analysis of fruit length in cucumber. Frontiers in Plant Science, 0, 14,  | 3.6             | 0          |
| 127 | Identification of major QTLs for drought tolerance in soybean, together with a novel candidate gene,<br><i>GmUAA6</i> . Journal of Experimental Botany, 2024, 75, 1852-1871.   | 4.8             | 0          |
| 128 | Morphological Analyses and QTL Mapping of Mottled Leaf in Zucchini (Cucurbita pepo L.).<br>International Journal of Molecular Sciences, 2024, 25, 2491.  | 4.1             | 0          |

CITATION REPORT