

The aryl hydrocarbon receptor-dependent disruption of WB-F344 epithelial cells is linked with induction of survival apoptosis

Toxicology

333, 37-44

DOI: [10.1016/j.tox.2015.04.001](https://doi.org/10.1016/j.tox.2015.04.001)

Citation Report

#	ARTICLE	IF	CITATIONS
1	Environmental Ligands of the Aryl Hydrocarbon Receptor and Their Effects in Models of Adult Liver Progenitor Cells. <i>Stem Cells International</i> , 2016, 2016, 1-14.	1.2	19
2	Effect of TCDD on the fate of epithelial cells isolated from human fetal palatal shelves (hFPECs). <i>Toxicology and Applied Pharmacology</i> , 2016, 305, 186-193.	1.3	14
3	2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD)-mediated deregulation of myeloid and sebaceous gland stem/progenitor cell homeostasis. <i>Archives of Toxicology</i> , 2017, 91, 2295-2301.	1.9	13
4	2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD) Disrupts Control of Cell Proliferation and Apoptosis in a Human Model of Adult Liver Progenitors. <i>Toxicological Sciences</i> , 2019, 172, 368-384.	1.4	9
5	Diesel exhaust particles inhibit lung branching morphogenesis via the YAP/TAZ pathway. <i>Science of the Total Environment</i> , 2023, 861, 160682.	3.9	0