The microtubule-associated protein PRC1 promotes early recurrence of hepatocellular carcinoma in association with the Wnt/?-catenin signalling pathway

| # | Paper | IF | Citations |
|---|-------|-----|-----------|
| 194 | MAPping the Wnt pathway to hepatocellular carcinoma recurrence. *Gut*, **2016**, 65, 1397-400 | 19.2 | 1 |
| 193 | TMED3 promotes hepatocellular carcinoma progression via IL-11/STAT3 signaling. *Scientific Reports*, **2016**, 6, 37070 | 4.9 | 37 |
| 192 | Genetic and functional analyses do not explain the association of high PRC1 expression with poor survival of breast carcinoma patients. *Biomedicine and Pharmacotherapy*, **2016**, 83, 857-864 | 7.5 | 12 |
| 191 | MELK is an oncogenic kinase essential for early hepatocellular carcinoma recurrence. **2016**, 383, 85-93 | | 51 |
| 190 | Elevated PRC1 in gastric carcinoma exerts oncogenic function and is targeted by piperlongumine in a p53-dependent manner. **2017**, 21, 1329-1341 | | 28 |
| 189 | Expression and functional assessment of candidate type 2 diabetes susceptibility genes identify four new genes contributing to human insulin secretion. **2017**, 6, 459-470 | | 32 |
| 188 | Atypical regulators of Wnt/Ωcatenin signaling as potential therapeutic targets in Hepatocellular Carcinoma. **2017**, 242, 1142-1149 | | 33 |
| 187 | Transcriptomic analysis of gene expression profiles of stomach carcinoma reveal abnormal expression of mitotic components. **2017**, 170, 41-49 | | 13 |
| 186 | STC2 as a novel mediator for Mus81-dependent proliferation and survival in hepatocellular carcinoma. **2017**, 388, 177-186 | | 27 |
| 185 | Screening and identification of key biomarkers in hepatocellular carcinoma: Evidence from bioinformatic analysis. *Oncology Reports*, **2017**, 38, 2607-2618 | 3.5 | 109 |
| 184 | Identification of prognostic genes through expression differentiation during metastatic process in lung adenocarcinoma. *Scientific Reports*, **2017**, 7, 11119 | 4.9 | 1 |
| 183 | The Transcriptomic Signature Of Disease Development And Progression Of Nonalcoholic Fatty Liver Disease. *Scientific Reports*, **2017**, 7, 17193 | 4.9 | 31 |
| 182 | PRC1 contributes to tumorigenesis of lung adenocarcinoma in association with the Wnt/Ωcatenin signaling pathway. *Molecular Cancer*, **2017**, 16, 108 | 42.1 | 52 |
| 181 | Emergence of aspirin as a promising chemopreventive and chemotherapeutic agent for liver cancer. **2017**, 8, e3112 | | 5 |
| 180 | Protein Regulator of Cytokinesis PRC1 Confers Chemoresistance and Predicts an Unfavorable Postoperative Survival of Hepatocellular Carcinoma Patients. **2017**, 8, 801-808 | | 19 |
| 179 | Protein regulator of cytokinesis-1 expression: prognostic value in lung squamous cell carcinoma patients. **2017**, 9, 2054-2060 | | 12 |
| 178 | SPC25 upregulation increases cancer stem cell properties in non-small cell lung adenocarcinoma cells and independently predicts poor survival. *Biomedicine and Pharmacotherapy*, **2018**, 100, 233-239 | 7.5 | 21 |

177    Transmembrane protein 170B is a novel breast tumorigenesis suppressor gene that inhibits the Wnt/ß-catenin pathway. **2018**, 9, 91    17

176    PRC1: Linking Cytokinesis, Chromosomal Instability, and Cancer Evolution. **2018**, 4, 59-73    30

175    Expression Differentiation Is Not Helpful in Identifying Prognostic Genes Based on TCGA Datasets. **2018**, 11, 292-299    7

174    Spindle pole body component 25 homolog expressed by ECM stiffening is required for lung cancer cell proliferation. **2018**, 500, 937-943    18

173    Chitosan-coated doxorubicin nano-particles drug delivery system inhibits cell growth of liver cancer via p53/PRC1 pathway. **2018**, 495, 414-420    29

172    Upregulation of thyroid hormone receptor interactor 13 is associated with human hepatocellular carcinoma. *Oncology Reports*, **2018**, 40, 3794-3802    3.5    7

171    CDK1-mediated BCL9 phosphorylation inhibits clathrin to promote mitotic Wnt signalling. **2018**, 37,    15

170    The microtubule-associated protein PRC1 is a potential therapeutic target for lung cancer. *Oncotarget*, **2018**, 9, 4985-4997    3.3    8

169    CDK7 activated beta-catenin/TCF signaling in hepatocellular carcinoma. **2018**, 370, 461-467    5

168    Reducing protein regulator of cytokinesis 1 as a prospective therapy for hepatocellular carcinoma. **2018**, 9, 534    24

167    MiR-137 suppresses migration and invasion by targeting EZH2-STAT3 signaling in human hepatocellular carcinoma. **2018**, 214, 1980-1986    23

166    Androgen receptor drives hepatocellular carcinogenesis by activating enhancer of zeste homolog 2-mediated Wnt/ß-catenin signaling. **2018**, 35, 155-166    19

165    p27 regulates the microtubule bundling activity of PRC1. **2018**, 1865, 1630-1639    8

164    ß-catenin regulates effects of miR-24 on the viability and autophagy of glioma cells. *Experimental and Therapeutic Medicine*, **2019**, 18, 1285-1290    2.1    4

163    Mechanisms of the Ase1/PRC1/MAP65 family in central spindle assembly. **2019**, 94, 2033-2048    10

162    EZH2 cooperates with E2F1 to stimulate expression of genes involved in adrenocortical carcinoma aggressiveness. **2019**, 121, 384-394    12

161    KIFC1 is activated by TCF-4 and promotes hepatocellular carcinoma pathogenesis by regulating HMGA1 transcriptional activity. *Journal of Experimental and Clinical Cancer Research*, **2019**, 38, 329    12.8    14

160    HEG1 indicates poor prognosis and promotes hepatocellular carcinoma invasion, metastasis, and EMT by activating Wnt/ß-catenin signaling. **2019**, 133, 1645-1662    32

| | | | |
|---|---|---|---|
| 159 | CCL14 serves as a novel prognostic factor and tumor suppressor of HCC by modulating cell cycle and promoting apoptosis. **2019**, 10, 796 | | 26 |
| 158 | Development and Validation of a 9-Gene Prognostic Signature in Patients With Multiple Myeloma. *Frontiers in Oncology*, **2018**, 8, 615 | 5.3 | 13 |
| 157 | HBX-induced miR-5188 impairs FOXO1 to stimulate ßcatenin nuclear translocation and promotes tumor stemness in hepatocellular carcinoma. **2019**, 9, 7583-7598 | | 38 |
| 156 | Weighted gene coexpression network analysis identifies a new biomarker of CENPF for prediction disease prognosis and progression in nonmuscle invasive bladder cancer. **2019**, 7, e982 | | 17 |
| 155 | Upregulation of long non-coding RNA FOXP4-AS1 and its regulatory network in hepatocellular carcinoma. **2019**, 12, 7025-7038 | | 10 |
| 154 | Screening and Identification of Potential Prognostic Biomarkers in Adrenocortical Carcinoma. *Frontiers in Genetics*, **2019**, 10, 821 | 4.5 | 13 |
| 153 | PRC1 gene silencing inhibits proliferation, invasion, and angiogenesis of retinoblastoma cells through the inhibition of the Wnt/ßcatenin signaling pathway. **2019**, 120, 16840-16852 | | 9 |
| 152 | Pseudogene RACGAP1P activates RACGAP1/Rho/ERK signalling axis as a competing endogenous RNA to promote hepatocellular carcinoma early recurrence. **2019**, 10, 426 | | 17 |
| 151 | Identification and validation of a potent multi-mRNA signature for the prediction of early relapse in hepatocellular carcinoma. *Carcinogenesis*, **2019**, 40, 840-852 | 4.6 | 9 |
| 150 | Phosphoregulation of the oncogenic protein regulator of cytokinesis 1 (PRC1) by the atypical CDK16/CCNY complex. **2019**, 51, 1-17 | | 11 |
| 149 | Analysis of potential key genes in very early hepatocellular carcinoma. **2019**, 17, 77 | | 19 |
| 148 | MiR-1-3p Inhibits Lung Adenocarcinoma Cell Tumorigenesis via Targeting Protein Regulator of Cytokinesis 1. *Frontiers in Oncology*, **2019**, 9, 120 | 5.3 | 13 |
| 147 | Transcriptome Analysis Revealed a Highly Connected Gene Module Associated With Cirrhosis to Hepatocellular Carcinoma Development. *Frontiers in Genetics*, **2019**, 10, 305 | 4.5 | 16 |
| 146 | AKIP1 promotes early recurrence of hepatocellular carcinoma through activating the Wnt/ßcatenin/CBP signaling pathway. **2019**, 38, 5516-5529 | | 25 |
| 145 | MicroRNA-194 inhibits cell invasion and migration in hepatocellular carcinoma through PRC1-mediated inhibition of Wnt/ßcatenin signaling pathway. **2019**, 51, 1314-1322 | | 15 |
| 144 | Identification of key genes and pathways in hepatocellular carcinoma: A preliminary bioinformatics analysis. **2019**, 98, e14287 | | 18 |
| 143 | Knockdown of the long noncoding RNA HOTTIP inhibits cell proliferation and enhances cell sensitivity to cisplatin by suppressing the Wnt/ßcatenin pathway in prostate cancer. **2019**, 120, 8965-8974 | | 21 |
| 142 | Prognostic genes of hepatocellular carcinoma based on gene coexpression network analysis. **2019**, 120, 11616 | | 22 |

| 141 | Identifying novel biomarkers in hepatocellular carcinoma by weighted gene co-expression network analysis. **2019**, 120, 11418 | | 26 |
|---|---|---|---|
| 140 | Identification of potential hub genes related to the progression and prognosis of hepatocellular carcinoma through integrated bioinformatics analysis. *Oncology Reports*, **2020**, 43, 133-146 | 3.5 | 29 |
| 139 | Identification of prognostic markers of lung cancer through bioinformatics analysis and in vitro experiments. *International Journal of Oncology*, **2020**, 56, 193-205 | 4.4 | 9 |
| 138 | Comprehensive investigation of key biomarkers and pathways in hepatitis B virus-related hepatocellular carcinoma. **2019**, 10, 5689-5704 | | 8 |
| 137 | PRC1 promotes GLI1-dependent osteopontin expression in association with the Wnt/£atenin signaling pathway and aggravates liver fibrosis. **2019**, 9, 100 | | 5 |
| 136 | Screening Hub Genes as Prognostic Biomarkers of Hepatocellular Carcinoma by Bioinformatics Analysis. **2019**, 28, 76S-86S | | 39 |
| 135 | Astragaloside IV (AS-IV) alleviates the malignant biological behavior of hepatocellular carcinoma Wnt/£atenin signaling pathway.. **2019**, 9, 35473-35482 | | 1 |
| 134 | miR-338-5p inhibits cell proliferation, colony formation, migration and cisplatin resistance in esophageal squamous cancer cells by targeting FERMT2. *Carcinogenesis*, **2019**, 40, 883-892 | 4.6 | 22 |
| 133 | Exploring and comparing of the gene expression and methylation differences between lung adenocarcinoma and squamous cell carcinoma. **2019**, 234, 4454-4459 | | 12 |
| 132 | Identification of core genes and outcomes in hepatocellular carcinoma by bioinformatics analysis. **2019**, 120, 10069-10081 | | 51 |
| 131 | Cytokinesis defects and cancer. **2019**, 19, 32-45 | | 99 |
| 130 | Erianin inhibits the oncogenic properties of hepatocellular carcinoma via inducing DNA damage and aberrant mitosis. **2020**, 182, 114266 | | 6 |
| 129 | ZFP36 Binds With PRC1 to Inhibit Tumor Growth and Increase 5-Fu Chemosensitivity of Hepatocellular Carcinoma. *Frontiers in Molecular Biosciences*, **2020**, 7, 126 | 5.6 | 5 |
| 128 | Aberrant DNA methylation results in altered gene expression in non-alcoholic steatohepatitis-related hepatocellular carcinomas. **2020**, 146, 2461-2477 | | 8 |
| 127 | Oncogenic Gene-Expression Programs in Leiomyosarcoma and Characterization of Conventional, Inflammatory, and Uterogenic Subtypes. **2020**, 18, 1302-1314 | | 9 |
| 126 | Genomic Landscape of HCC. **2020**, 19, 448-461 | | 6 |
| 125 | A genomic-clinicopathologic nomogram for predicting overall survival of hepatocellular carcinoma. *BMC Cancer*, **2020**, 20, 1176 | 4.8 | 2 |
| 124 | KIF2C: a novel link between Wnt/£atenin and mTORC1 signaling in the pathogenesis of hepatocellular carcinoma. **2021**, 12, 788-809 | | 20 |

| | | | |
|---|---|---|---|
| 123 | Using mRNAsi to identify prognostic-related genes in endometrial carcinoma based on WGCNA. **2020**, 258, 118231 | | 9 |
| 122 | Potential mechanism of RRM2 for promoting Cervical Cancer based on weighted gene co-expression network analysis. **2020**, 17, 2362-2372 | | 6 |
| 121 | Overexpression of microRNA-203 Suppresses Proliferation, Invasion, and Migration while Accelerating Apoptosis of CSCC Cell Line SCL-1. **2020**, 21, 428-440 | | 1 |
| 120 | The RNA-binding protein RBM47 inhibits non-small cell lung carcinoma metastasis through modulation of AXIN1 mRNA stability and Wnt/βcatentin signaling. **2020**, 34, 31-39 | | 12 |
| 119 | Random-forest algorithm based biomarkers in predicting prognosis in the patients with hepatocellular carcinoma. *Cancer Cell International*, **2020**, 20, 251 | 6.4 | 11 |
| 118 | Protein regulator of cytokinesis 1 regulates chromosome dynamics and cytoplasmic division during mouse oocyte meiotic maturation and early embryonic development. **2020**, 287, 5130-5147 | | 0 |
| 117 | Role of lncRNAHCP5/microRNA-525-5p/PRC1 crosstalk in the malignant behaviors of ovarian cancer cells. **2020**, 394, 112129 | | 11 |
| 116 | Heat-Shock protein A12A is a novel PCNA-binding protein and promotes hepatocellular carcinoma growth. **2020**, 287, 5464-5477 | | 5 |
| 115 | Tumor Microenvironment. *Advances in Experimental Medicine and Biology*, **2020**, | 3.6 | 1 |
| 114 | Silencing MYH9 blocks HBx-induced GSK3βubiquitination and degradation to inhibit tumor stemness in hepatocellular carcinoma. **2020**, 5, 13 | | 43 |
| 113 | URH49 exports mRNA by remodeling complex formation and mediating the NXF1-dependent pathway. **2020**, 1863, 194480 | | 3 |
| 112 | Establishment of a Genomic-Clinicopathologic Nomogram for Predicting Early Recurrence of Hepatocellular Carcinoma After R0 Resection. **2021**, 25, 112-124 | | 5 |
| 111 | KIF11 is upregulated in colorectal cancer and silencing of it impairs tumor growth and sensitizes colorectal cancer cells to oxaliplatin via p53/GSK3βsignaling. **2021**, 12, 3741-3753 | | 4 |
| 110 | A Deep Look into the Program of Rapid Tumor Growth of Hepatocellular Carcinoma. **2021**, 9, 22-31 | | 0 |
| 109 | Analysis of transcriptome in the relationship between expression of PRC1 protein and prognosis of patients with cholangiocarcinoma. **2021**, 49, 300060521989200 | | 0 |
| 108 | Tektin4 loss promotes triple-negative breast cancer metastasis through HDAC6-mediated tubulin deacetylation and increases sensitivity to HDAC6 inhibitor. **2021**, 40, 2323-2334 | | 1 |
| 107 | Shc3 promotes hepatocellular carcinoma stemness and drug resistance by interacting with βcatenin to inhibit its ubiquitin degradation pathway. **2021**, 12, 278 | | 8 |
| 106 | Hsa-circRNA-G004213 promotes cisplatin sensitivity by regulating miR-513b-5p/PRPF39 in liver cancer. **2021**, 23, | | 4 |

| | | | |
|---|---|---|---|
| 105 | Sec62 promotes stemness and chemoresistance of human colorectal cancer through activating Wnt/β-catenin pathway. *Journal of Experimental and Clinical Cancer Research*, **2021**, 40, 132 | 12.8 | 26 |
| 104 | EB2 promotes hepatocellular carcinoma proliferation and metastasis via MAPK/ERK pathway by modulating microtubule dynamics. **2021**, 135, 847-864 | | 3 |
| 103 | Dysregulation of Wnt/β-catenin signaling by protein kinases in hepatocellular carcinoma and its therapeutic application. **2021**, 112, 1695-1706 | | 12 |
| 102 | Cytokinesis regulators as potential diagnostic and therapeutic biomarkers for human hepatocellular carcinoma. **2021**, 246, 1343-1354 | | 0 |
| 101 | Integrative analysis identifies key mRNA biomarkers for diagnosis, prognosis, and therapeutic targets of HCV-associated hepatocellular carcinoma. *Aging*, **2021**, 13, 12865-12895 | 5.6 | 2 |
| 100 | Recipient C7 rs9292795 genotype and the risk of hepatocellular carcinoma recurrence after orthotopic liver transplantation in a Han Chinese population. *BMC Cancer*, **2021**, 21, 521 | 4.8 | 0 |
| 99 | Identification and validation of a hub gene prognostic index for hepatocellular carcinoma. **2021**, 17, 2193-2208 | | 1 |
| 98 | TSHZ2 is an EGF-regulated tumor suppressor that binds to the cytokinesis regulator PRC1 and inhibits metastasis. **2021**, 14, | | 0 |
| 97 | Gene expression of cytokinesis regulators and has no prognostic role in colorectal and pancreatic cancer. *Oncology Letters*, **2021**, 22, 598 | 2.6 | 2 |
| 96 | High SGO2 Expression Predicts Poor Overall Survival: A Potential Therapeutic Target for Hepatocellular Carcinoma. **2021**, 12, | | 0 |
| 95 | Paeoniflorin Affects Hepatocellular Carcinoma Progression by Inhibiting Wnt/β-Catenin Pathway through Downregulation of 5-HT1D. **2021**, 22, 1246-1253 | | 2 |
| 94 | DDX56 modulates post-transcriptional Wnt signaling through miRNAs and is associated with early recurrence in squamous cell lung carcinoma. *Molecular Cancer*, **2021**, 20, 108 | 42.1 | 4 |
| 93 | BIOINFORMATICS APPROACH ON ACTION AND MECHANISM OF BROMELAIN IN HEPATOCELLULAR CARCINOMA. *Asian Journal of Pharmaceutical and Clinical Research*, 122-129 | 0.4 | |
| 92 | Viral Hepatitides, Inflammation and Tumour Microenvironment. *Advances in Experimental Medicine and Biology*, **2020**, 1263, 25-43 | 3.6 | 4 |
| 91 | MicroRNA-766-3p Inhibits Tumour Progression by Targeting Wnt3a in Hepatocellular Carcinoma. *Molecules and Cells*, **2018**, 41, 830-841 | 3.5 | 25 |
| 90 | Identification of hub genes in hepatocellular carcinoma using integrated bioinformatic analysis. *Aging*, **2020**, 12, 5439-5468 | 5.6 | 18 |
| 89 | KIF23 activated Wnt/β-catenin signaling pathway through direct interaction with Amer1 in gastric cancer. *Aging*, **2020**, 12, 8372-8396 | 5.6 | 13 |
| 88 | Central spindle proteins and mitotic kinesins are direct transcriptional targets of MuvB, B-MYB and FOXM1 in breast cancer cell lines and are potential targets for therapy. *Oncotarget*, **2017**, 8, 11160-11172 | 3.3 | 28 |

| 87 | A novel PHD-finger protein 14/KIF4A complex overexpressed in lung cancer is involved in cell mitosis regulation and tumorigenesis. *Oncotarget*, **2017**, 8, 19684-19698 | 3.3 | 17 |
|---|---|---|---|
| 86 | Identification of Platform-Independent Diagnostic Biomarker Panel for Hepatocellular Carcinoma Using Large-Scale Transcriptomics Data. *Frontiers in Genetics*, **2019**, 10, 1306 | 4.5 | 18 |
| 85 | Aberrant regulation of Wnt signaling in hepatocellular carcinoma. *World Journal of Gastroenterology*, **2016**, 22, 7486-99 | 5.6 | 51 |
| 84 | CCNB2, NUSAP1 and TK1 are associated with the prognosis and progression of hepatocellular carcinoma, as revealed by co-expression analysis. *Experimental and Therapeutic Medicine*, **2020**, 19, 2679-2689 | 2.1 | 1 |
| 83 | Overexpression of PRC1 indicates a poor prognosis in ovarian cancer. *International Journal of Oncology*, **2020**, 56, 685-696 | 4.4 | 10 |
| 82 | Identification of the association between HMMR expression and progression of hepatocellular carcinoma via construction of a co-expression network. *Oncology Letters*, **2020**, 20, 2645-2654 | 2.6 | 6 |
| 81 | MicroRNAs target the Wnt/ß-catenin signaling pathway to regulate epithelial-mesenchymal transition in cancer (Review). *Oncology Reports*, **2020**, 44, 1299-1313 | 3.5 | 12 |
| 80 | Bioinformatics analysis of LINC01554 and its co-expressed genes in hepatocellular carcinoma. *Oncology Reports*, **2020**, 44, 2185-2197 | 3.5 | 3 |
| 79 | Immunobiology of hepatocarcinogenesis: Ways to go or almost there?. *World Journal of Gastrointestinal Pathophysiology*, **2016**, 7, 242-55 | 3.2 | 10 |
| 78 | A genomic-clinical nomogram predicting recurrence-free survival for patients diagnosed with hepatocellular carcinoma. *PeerJ*, **2019**, 7, e7942 | 3.1 | 5 |
| 77 | Transcriptome analysis revealed key prognostic genes and microRNAs in hepatocellular carcinoma. *PeerJ*, **2020**, 8, e8930 | 3.1 | 18 |
| 76 | The phosphorylated retinoid X receptor-ß promotes diethylnitrosamine-induced hepatocarcinogenesis in mice through the activation of ß-catenin signaling pathway. *Carcinogenesis*, **2021**, | 4.6 | 0 |
| 75 | Circular RNA circIPO11 drives self-renewal of liver cancer initiating cells via Hedgehog signaling. *Molecular Cancer*, **2021**, 20, 132 | 42.1 | 11 |
| 74 | Screening and predicted value of potential biomarkers for breast cancer using bioinformatics analysis. *Scientific Reports*, **2021**, 11, 20799 | 4.9 | 3 |
| 73 | Whole-exome sequencing of rectal cancer identifies locally recurrent mutations in the Wnt pathway. *Aging*, **2021**, 13, 23262-23283 | 5.6 | 0 |
| 72 | Wnt Signaling Pathway Is among the Drivers of Liver Metastasis. *Livers*, **2021**, 1, 180-200 | | 0 |
| 71 | Identification of Platform-Independent Diagnostic Biomarker Panel for Hepatocellular Carcinoma using Large-scale Transcriptomics Data. | | |
| 70 | promotes cell proliferation and cell cycle progression by regulating family molecules and pathway in non-small cell lung cancer.. *Translational Cancer Research*, **2019**, 8, 2059-2072 | 0.3 | 2 |

| 69 | Nuclear receptor FXR impairs SK-Hep-1 cell migration and invasion by inhibiting the Wnt/Ecatenin signaling pathway. *Oncology Letters*, **2020**, 20, 161 | 2.6 | 2 |
|---|---|---|---|
| 68 | Transcriptome and interactome analyses identify theTP53interacting geneRCCD1as a candidate susceptibility gene at the 15p26.1 breast and ovarian cancer risk locus. | | |
| 67 | Nuclear division cycle 80 complex is associated with malignancy and predicts poor survival of hepatocellular carcinoma. *International Journal of Clinical and Experimental Pathology*, **2019**, 12, 1233-1247 | 1.4 | 3 |
| 66 | Identifying drug candidates for hepatocellular carcinoma based on differentially expressed genes. *American Journal of Translational Research (discontinued)*, **2020**, 12, 2664-2674 | 3 | |
| 65 | Silybum marianum total extract, silymarin and silibinin abate hepatocarcinogenesis and hepatocellular carcinoma growth via modulation of the HGF/c-Met, Wnt/Ecatenin, and PI3K/Akt/mTOR signaling pathways. *Biomedicine and Pharmacotherapy*, **2021**, 145, 112409 | 7.5 | 2 |
| 64 | Pleiotropic roles of FXR in liver and colorectal cancers.. *Molecular and Cellular Endocrinology*, **2022**, 111543 | 4.1 | 0 |
| 63 | Construction and Comprehensive Analysis of a circRNA-miRNA-mRNA Regulatory Network to Reveal the Pathogenesis of Hepatocellular Carcinoma.. *Frontiers in Molecular Biosciences*, **2022**, 9, 801478 | 5.6 | |
| 62 | Combinatorial Cetuximab targeted polymeric nanocomplexes reduce PRC1 level and abrogate growth of metastatic hepatocellular carcinoma in vivo with efficient radionuclide uptake.. *Nanomedicine: Nanotechnology, Biology, and Medicine*, **2022**, 102529 | 6 | 0 |
| 61 | Study on the expression of TOP2A in hepatocellular carcinoma and its relationship with patient prognosis.. *Cancer Cell International*, **2022**, 22, 29 | 6.4 | 1 |
| 60 | Identification of the critical genes and miRNAs in hepatocellular carcinoma by integrated bioinformatics analysis.. *Medical Oncology*, **2022**, 39, 21 | 3.7 | 0 |
| 59 | Novel Gene Signatures as Prognostic Biomarkers for Predicting the Recurrence of Hepatocellular Carcinoma.. *Cancers*, **2022**, 14, | 6.6 | 0 |
| 58 | RNA Binding Protein in Hepatocellular Carcinoma: Recent Research Progress. *Advances in Clinical Medicine*, **2022**, 12, 2103-2109 | 0 | |
| 57 | Dysregulated cell-signaling pathways in hepatocellular carcinoma: causes and therapeutic options. **2022**, 337-355 | | |
| 56 | An Immune-Related Gene Signature Predicts Prognosis and Immunotherapy Response in Hepatocellular Carcinoma.. *Combinatorial Chemistry and High Throughput Screening*, **2022**, | 1.3 | |
| 55 | Identification of Immune-Related Breast Cancer Chemotherapy Resistance Genes Bioinformatics Approaches.. *Frontiers in Oncology*, **2022**, 12, 772723 | 5.3 | |
| 54 | NAP1L1 promotes tumor proliferation through HDGF/C-JUN signaling in ovarian cancer.. *BMC Cancer*, **2022**, 22, 339 | 4.8 | 1 |
| 53 | The aberrant upregulation of exon 10-inclusive SREK1 through SRSF10 acts as an oncogenic driver in human hepatocellular carcinoma.. *Nature Communications*, **2022**, 13, 1363 | 17.4 | 0 |
| 52 | Encouraging specific biomarkers-based therapeutic strategies for hepatocellular carcinoma. *World Journal of Clinical Cases*, **2022**, 10, 3321-3333 | 1.6 | 0 |

| 51 | CCNB2, CDC20, AURKA, TOP2A, MELK, NCAPG, KIF20A, UBE2C, PRC1, and ASPM May Be Potential Therapeutic Targets for Hepatocellular Carcinoma Using Integrated Bioinformatic Analysis.. *International Journal of General Medicine*, **2021**, 14, 10185-10194 | 2.3 | 2 |
|---|---|---|---|
| 50 | Liquid-Liquid Phase Separation-Related Genes Associated with Tumor Grade and Prognosis in Hepatocellular Carcinoma: A Bioinformatic Study.. *International Journal of General Medicine*, **2021**, 14, 9671-9679 | 2.3 | 1 |
| 49 | Image_1.TIF. **2019**, | | |
| 48 | Image_2.TIF. **2019**, | | |
| 47 | Image_3.TIF. **2019**, | | |
| 46 | Image_4.TIF. **2019**, | | |
| 45 | Image_5.TIF. **2019**, | | |
| 44 | Image_6.TIF. **2019**, | | |
| 43 | Table_1.XLSX. **2019**, | | |
| 42 | Table_2.XLSX. **2019**, | | |
| 41 | Table_3.XLSX. **2019**, | | |
| 40 | Table_4.XLSX. **2019**, | | |
| 39 | DataSheet_1.xlsx. **2020**, | | |
| 38 | Image_1.jpeg. **2020**, | | |
| 37 | Image_2.jpeg. **2020**, | | |
| 36 | Image_3.jpeg. **2020**, | | |
| 35 | Image_4.png. **2020**, | | |
| 34 | Image_5.png. **2020**, | | |

33    Image_6.png. **2020**,

32    Image_1.tif. **2019**,

31    Image_2.tif. **2019**,

30    Image_3.tif. **2019**,

29    Image_4.tif. **2019**,

28    Image_5.tif. **2019**,

27    Image_1.TIF. **2020**,

26    Data_Sheet_1.docx. **2019**,

25    Data_Sheet_2.docx. **2019**,

24    Image_1.TIF. **2019**,

23    Image_2.TIF. **2019**,

22    Table_1.docx. **2019**,

21    Table_2.docx. **2019**,

20    Table_3.docx. **2019**,

19    CDK16 promotes the progression and metastasis of triple-negative breast cancer by phosphorylating PRC1.. *Journal of Experimental and Clinical Cancer Research*, **2022**, 41, 149        12.8    2

18    Identification of a circRNA/miRNA/mRNA ceRNA Network as a Cell Cycle-Related Regulator for Chronic Sinusitis with Nasal Polyps.. *Journal of Inflammation Research*, **2022**, 15, 2601-2615        4.8

17    Kinesin family member 23, regulated by FOXM1, promotes triple negative breast cancer progression via activating Wnt/Ε-catenin pathway.. *Journal of Experimental and Clinical Cancer Research*, **2022**, 41, 168        12.8    1

16    Protein Regulator of Cytokinesis 1 (PRC1) Upregulation Promotes Immune Suppression in Liver Hepatocellular Carcinoma. **2022**, 2022, 1-27

| 15 | TRIM36 inhibits tumorigenesis through the Wnt/ßcatenin pathway and promotes caspase-dependent apoptosis in hepatocellular carcinoma. **2022**, 22, | 0 |
| 14 | MicroRNA-mediated regulation of key signaling pathways in hepatocellular carcinoma: A mechanistic insight. 13, | 2 |
| 13 | Expression of the cytokinesis regulator PRC1 results in p53-pathway activation in A549 cells but does not directly regulate gene expression in the nucleus. 1-14 | 0 |
| 12 | Bioinformatic and Experimental Analyses Reveal That KIF4A Is a Biomarker of Therapeutic Sensitivity and Predicts Prognosis in Cervical Cancer Patients. | 0 |
| 11 | System Biology Approaches Identified Novel Biomarkers and their Signaling Pathways Involved in Renal Cell Carcinoma (RCC) with Different Human Diseases. | 1 |
| 10 | ESPL1 is Elevated in Hepatocellular Carcinoma and Predicts Prognosis. Volume 15, 8381-8398 | 0 |
| 9 | Prognosis-Related Molecular Subtypes and Immune Features Associated with Hepatocellular Carcinoma. **2022**, 14, 5721 | 0 |
| 8 | Comprehensive analysis of prognostic value, relationship to cell cycle, immune infiltration and m6A modification of ZSCAN20 in hepatocellular carcinoma. | 0 |
| 7 | Multi-omics on truncatingASXL1mutations in Bohring Opitz syndrome identify dysregulation of canonical and non-canonical Wnt signaling. | 0 |
| 6 | CircRNA-mTOR Promotes Hepatocellular Carcinoma Progression and Lenvatinib Resistance via PSIP1/c-Myc Axis Nuclear Translocation and Partially through Increasing EGFR Expression. | 0 |
| 5 | Feiyiliu Mixture sensitizes EGFRDel19/T790M/C797S mutant non-small cell lung cancer to osimertinib by attenuating the PRC1/Wnt/EGFR pathway. 14, | 0 |
| 4 | Combined transcriptomics and in-silico approach uncovers the role of prognostic biomarkers in hepatocellular carcinoma. **2023**, 35, 201154 | 0 |
| 3 | KIF23, under regulation by androgen receptor, can promote the deterioration of nasopharyngeal carcinoma by activating the Wnt/ßcatenin signaling pathway. | 0 |
| 2 | Clinical value of PRC1 and DLGAP5 and immunosuppressive T cells overexpressing them in HCC based on transcriptome data. | 0 |
| 1 | KIF23, under regulation by androgen receptor, contributes to nasopharyngeal carcinoma deterioration by activating the Wnt/ßcatenin signaling pathway. **2023**, 23, | 0 |