

# MMP-2 gene polymorphisms are associated with type A diameters in patients

Medicine (United States)

95, e5175

DOI: [10.1097/md.0000000000005175](https://doi.org/10.1097/md.0000000000005175)

Citation Report

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 1  | Polymorphisms and pharmacogenomics for the toxicity of methotrexate monotherapy in patients with rheumatoid arthritis. <i>Medicine (United States)</i> , 2017, 96, e6337.  | 0.4 | 33        |
| 2  | MRI-measured myocardial iron load in patients with severe diabetic heart failure. <i>Clinical Radiology</i> , 2018, 73, 324.e1-324.e7.   | 0.5 | 3         |
| 3  | Matrix metalloproteinase family polymorphisms and the risk of aortic aneurysmal diseases: A systematic review and meta-analysis. <i>Clinical Genetics</i> , 2018, 93, 15-32.   | 1.0 | 26        |
| 4  | Matrix metalloproteinases are regulated by MicroRNA 320 in macrophages and are associated with aortic dissection. <i>Experimental Cell Research</i> , 2018, 370, 98-102.   | 1.2 | 18        |
| 5  | Cerebral small vessel disease and systemic arteriopathy in intracranial arterial dolichoectasia patients. <i>Acta Neurologica Scandinavica</i> , 2019, 139, 150-157.   | 1.0 | 5         |
| 6  | Insulin-like growth factor-binding protein 3 inhibits angiotensin II-induced aortic smooth muscle cell phenotypic switch and matrix metalloproteinase expression. <i>Experimental Physiology</i> , 2020, 105, 1827-1839.           | 0.9 | 4         |
| 7  | Increased Circulating Angiopoietin-Like Protein 8 Levels Are Associated with Thoracic Aortic Dissection and Higher Inflammatory Conditions. <i>Cardiovascular Drugs and Therapy</i> , 2020, 34, 65-77.                             | 1.3 | 25        |
| 8  | Efficient detection of differentially methylated regions in the genome of patients with thoracic aortic dissection and association with MMP2 hypermethylation. <i>Experimental and Therapeutic Medicine</i> , 2020, 20, 1073-1081. | 0.8 | 7         |
| 9  | Analysis of patients with thoracic aortic aneurysm taking into account risk factors, procedure and prognosis in 5-year clinical observation. <i>Annales Academiae Medicae Silesiensis</i> , 2018, 72, 193-202.                     | 0.1 | 0         |
| 10 | Remodeling of abdominal aortic aneurysm sac following endovascular aortic repair: association with clinical, surgical, and genetic factors. <i>Cardiovascular Pathology</i> , 2022, 58, 107405.                                    | 0.7 | 6         |
| 11 | The Predictive Role of Plasma Biomarkers in the Evolution of Aortopathies Associated with Congenital Heart Malformations. <i>International Journal of Molecular Sciences</i> , 2022, 23, 4993.                                     | 1.8 | 3         |
| 12 | ADAMTS-7 deficiency attenuates thoracic aortic aneurysm and dissection in mice. <i>Journal of Molecular Medicine</i> , 2023, 101, 237-248.   | 1.7 | 4         |