

CITATION REPORT

List of articles citing

Sitting and Activity Time in People With Stroke

DOI: 10.2522/ptj.20140522

Physical Therapy, 2016, 96, 193-201.

Source: <https://exaly.com/paper-pdf/65496854/citation-report.pdf>

Version: 2024-04-25

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
123	Recovery of ambulation activity across the first six months post-stroke. 2016 , 49, 271-276		25
122	Sitting time and physical activity after stroke: physical ability is only part of the story. <i>Topics in Stroke Rehabilitation</i> , 2016 , 23, 36-42	2.6	40
121	Considerations when using the activPAL monitor in field-based research with adult populations. 2017 , 6, 162-178		209
120	How Physically Active Are People Following Stroke? Systematic Review and Quantitative Synthesis. <i>Physical Therapy</i> , 2017 , 97, 707-717	3.3	126
119	Sit to stand activity during stroke rehabilitation. <i>Topics in Stroke Rehabilitation</i> , 2017 , 24, 562-566	2.6	10
118	Cardiorespiratory fitness and walking endurance improvements after 12 months of an individualised home and community-based exercise programme for people after stroke. 2017 , 31, 1617-1624		10
117	From Disease to Health: Physical Therapy Health Promotion Practices for Secondary Prevention in Adult and Pediatric Neurologic Populations. 2017 , 41 Suppl 3, S46-S54		9
116	Sleep Duration, Sedentary Behavior, Physical Activity, and Quality of Life after Inpatient Stroke Rehabilitation. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2017 , 26, 2004-2012	2.8	29
115	Breaking up sitting time after stroke (BUST-Stroke). 2017 , 12, 425-429		11
114	Measuring Activity Performance of Older Adults Using the activPAL: A Rapid Review. 2017 , 5,		9
113	Perceptions of physical activity and walking in an early stage after stroke or acquired brain injury. 2017 , 12, e0173463		22
112	Inactive lifestyles and sedentary behavior in persons with chronic aneurysmal subarachnoid hemorrhage: evidence from accelerometer-based activity monitoring. <i>Journal of NeuroEngineering and Rehabilitation</i> , 2017 , 14, 120	5.3	5
111	Effect of accelerometer-based feedback on physical activity in hospitalized patients with ischemic stroke: a randomized controlled trial. 2018 , 32, 1047-1056		15
110	Reliability of intensity-based physical activity measurement using an activity monitor in people with subacute stroke in the hospital setting: a cross-sectional study. <i>Topics in Stroke Rehabilitation</i> , 2018 , 25, 288-294	2.6	10
109	Objectively assessed physical activity and associated factors of sedentary behavior among survivors of stroke living in Cape Town, South Africa. 2018 , 40, 2509-2515		14
108	Best practice guidelines for the measurement of physical activity levels in stroke survivors: a secondary analysis of an observational study. 2018 , 41, 14-19		21
107	Rating of perceived exertion with Borg scale in stroke over two common activities of the daily living. <i>Topics in Stroke Rehabilitation</i> , 2018 , 25, 145-149	2.6	8

106	Human Postural Balance Based on the Surface Gravity Measuring System. 2018,	
105	Effectiveness of a multimodal low-to-moderate intensity exercise rehabilitation program for stroke survivors. 2018, 53, 147-153	2
104	Using Intervention Mapping to develop and implement a home-based sedentary behavior change intervention after stroke. 2020, 10, 87-95	3
103	Quality of life after stroke: impact of clinical and sociodemographic factors. 2018, 73, e418	57
102	The Feasibility and Longitudinal Effects of a Home-Based Sedentary Behavior Change Intervention After Stroke. 2018, 99, 2540-2547	22
101	Muscle Thickness and Echo Intensity of the Abdominal and Lower Extremity Muscles in Stroke Survivors. 2018, 14, 549-554	13
100	Frequent, short bouts of light-intensity exercises while standing decreases systolic blood pressure: Breaking Up Sitting Time after Stroke (BUST-Stroke) trial. 2018, 13, 932-940	25
99	Breaking up sitting time after stroke (BUST-stroke). 2018, 13, 921-931	11
98	Difference in independent mobility improvement from admission to discharge between subacute stroke patients using knee-ankle-foot and those using ankle-foot orthoses. 2018, 30, 1003-1008	3
97	Go Home, Sit Less: The Impact of Home Versus Hospital Rehabilitation Environment on Activity Levels of Stroke Survivors. 2018, 99, 2216-2221.e1	16
96	Protocol and pilot study of a short message service-guided training after acute stroke/transient ischemic attack to increase walking capacity and physical activity. 2018, 11, 109-114	6
95	Use of Mobile Devices to Measure Outcomes in Clinical Research, 2010-2016: A Systematic Literature Review. 2018, 2, 11-30	52
94	Interventions for reducing sedentary behaviour in people with stroke. 2018,	1
93	Activity monitors for increasing physical activity in adult stroke survivors. 2018, 7, CD012543	15
92	The known-groups validity of intensity-based physical activity measurement using an accelerometer in people with subacute stroke. 2018, 30, 507-513	9
91	Home-based physical activity incentive and education program in subacute phase of stroke recovery (Ticaa'dom): study protocol for a randomized controlled trial. 2018, 19, 68	9
90	Preferred Placement and Usability of a Smart Textile System vs. Inertial Measurement Units for Activity Monitoring. 2018, 18,	30
89	Sedentary behavior after stroke: A new target for therapeutic intervention. 2019, 14, 9-11	6

88	Characteristics of intensity-based physical activity according to gait ability in people hospitalized with subacute stroke: a cross-sectional study. 2019 , 22, 17-25		7
87	Patterns of Sedentary Behavior in the First Month After Acute Coronary Syndrome. 2019 , 8, e011585		2
86	Association of Perceived Built Environment Attributes with Objectively Measured Physical Activity in Community-Dwelling Ambulatory Patients with Stroke. 2019 , 16,		6
85	Three distinct physical behavior types in fatigued patients with multiple sclerosis. <i>Journal of NeuroEngineering and Rehabilitation</i> , 2019 , 16, 105	5-3	5
84	Physical Activity and Sedentary Behaviour Patterns in 326 Persons with COPD before Starting a Pulmonary Rehabilitation: A Cluster Analysis. 2019 , 8,		16
83	What is the effect of interrupting prolonged sitting with frequent bouts of physical activity or standing on first or recurrent stroke risk factors? A scoping review. 2019 , 14, e0217981		8
82	Poststroke Fatigue. 2019 , 50, 1927-1933		14
81	Identifying factors associated with sedentary time after stroke. Secondary analysis of pooled data from nine primary studies. <i>Topics in Stroke Rehabilitation</i> , 2019 , 26, 327-334	2.6	13
80	Supervised exercise delivered via telehealth in real time to manage chronic conditions in adults: a protocol for a scoping review to inform future research in stroke survivors. 2019 , 9, e027416		7
79	Development and Validation of a Questionnaire to Assess Barriers to Physical Activity After Stroke: The Barriers to Physical Activity After Stroke Scale. 2019 , 100, 1672-1679		3
78	Participation, Fear of Falling, and Upper Limb Impairment are Associated with High Sitting Time in People with Stroke. 2019 , 33, 181-196		5
77	Measurement properties of self-report physical activity assessment tools for patients with stroke: a systematic review. 2019 , 23, 476-490		9
76	Early effects of a knee-ankle-foot orthosis on static standing balance in people with subacute stroke. 2019 , 31, 127-131		5
75	Prevalence of 3 Healthy Lifestyle Behaviors Among US Adults With and Without History of Stroke. 2019 , 16, E23		5
74	Functional measurement post-stroke via mobile application and body-worn sensor technology. 2019 , 5, 47		5
73	Wireless Physical Activity Monitor Use Among Adults Living With HIV: A Scoping Review. 2019 , 37, 17-28		3
72	Measuring Community Mobility in Survivors of Stroke Using Global Positioning System Technology: A Prospective Observational Study. 2019 , 43, 175-185		3
71	Exploring the Association Between Physical Activity, Sedentary Behavior, and High-Sensitivity C-Reactive Protein Among Stroke Survivors. 2019 , 27, 360-366		2

70	Two Days of Measurement Provides Reliable Estimates of Physical Activity Poststroke: An Observational Study. 2019 , 100, 883-890		13
69	Breaking up sitting time after stroke - How much less sitting is needed to improve blood pressure after stroke (BUST-BP-Dose): Protocol for a dose-finding study. 2019 , 13, 100310		2
68	Long-Term Effect of Promoting In-Hospital Physical Activity on Postdischarge Patients with Mild Ischemic Stroke. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2019 , 28, 1048-1055	2.8	4
67	Level of physical activity in men and women with chronic stroke. <i>Physiotherapy Theory and Practice</i> , 2019 , 35, 947-955	1.5	7
66	Efficacy of interventions aimed at improving physical activity in individuals with stroke: a systematic review. 2020 , 42, 902-917		9
65	Examining daily physical activity in community-dwelling adults with stroke using social cognitive theory: an exploratory, qualitative study. 2020 , 42, 2631-2639		10
64	Daytime physical activity at admission is associated with improvement of gait independence 1 month later in people with subacute stroke: a longitudinal study. <i>Topics in Stroke Rehabilitation</i> , 2020 , 27, 25-32	2.6	7
63	Influence of Interventions on Daily Physical Activity and Sedentary Behavior after Stroke: A Systematic Review. 2020 , 12, 186-201		6
62	Exercise Programs Delivered According to Guidelines Improve Mobility in People With Stroke: A Systematic Review and Meta-analysis. 2020 , 101, 154-165		5
61	Impact of mHealth technology on adherence to healthy PA after stroke: a randomized study. <i>Topics in Stroke Rehabilitation</i> , 2020 , 27, 354-368	2.6	18
60	Is long-bout sedentary behaviour associated with long-term glucose levels 3 months after acute ischaemic stroke? A prospective observational cohort study. 2020 , 10, e037475		1
59	A co-production approach guided by the behaviour change wheel to develop an intervention for reducing sedentary behaviour after stroke. 2020 , 6, 115		7
58	Factors influencing sedentary behaviours after stroke: findings from qualitative observations and interviews with stroke survivors and their caregivers. 2020 , 20, 967		9
57	A comparison of self-reported and device measured sedentary behaviour in adults: a systematic review and meta-analysis. 2020 , 17, 31		81
56	Barriers and Facilitators to Aerobic Exercise Implementation in Stroke Rehabilitation: A Scoping Review. 2020 , 44, 179-187		6
55	Active and sedentary bouts in people after stroke and healthy controls: An observational study. 2020 , 25, e1845		1
54	Effects on walking performance and lower body strength by short message service guided training after stroke or transient ischemic attack (The STROKEWALK Study): a randomized controlled trial. 2021 , 35, 276-287		1
53	Levels of physical activity and quality of life among community-dwelling adults with stroke in a developing country. 2021 , 23, 165-170		5

52	Heart Disease and Stroke Statistics-2021 Update: A Report From the American Heart Association. <i>Circulation</i> , 2021 , 143, e254-e743	16.7	1087
51	Association of Barriers, Fear of Falling and Fatigue with Objectively Measured Physical Activity and Sedentary Behavior in Chronic Stroke. 2021 , 10,		2
50	Levels of physical activity before and after stroke in relation to early cognitive function. 2021 , 11, 9078		4
49	Validity of the International Physical Activity Questionnaire Long Form for Assessing Physical Activity and Sedentary Behavior in Subjects with Chronic Stroke. 2021 , 18,		0
48	An Exploration of Sedentary Behavior Patterns in Community-Dwelling People With Stroke: A Cluster-Based Analysis. 2021 , 45, 221-227		0
47	How active are stroke patients in physiotherapy sessions and is this associated with stroke severity?. 2021 , 1-7		1
46	Decision-Making on Referral to Primary Care Physiotherapy After Inpatient Stroke Rehabilitation. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2021 , 30, 105667	2.8	
45	Acute Effects of Frequent Light-Intensity Standing-Based Exercises That Interrupt 8 Hours of Prolonged Sitting on Postprandial Glucose in Stroke Survivors: A Dose-Escalation Trial. 2021 , 18, 644-652		0
44	Interventions for reducing sedentary behaviour in people with stroke. 2021 , 6, CD012996		6
43	Comparing the physical activity of stroke survivors in high-income countries and low to middle-income countries. 2021 , 26, e1918		
42	Behavioral Medicine for Sedentary Behavior, Daily Physical Activity, and Exercise to Prevent Cardiovascular Disease: A Review. 2021 , 23, 48		0
41	Longitudinal Impact of Community-Based Rehabilitation Programs on Functional Recovery After Stroke: A Scoping Review. 108482232110355		
40	A qualitative study of sedentary behaviours in stroke survivors: non-participant observations and interviews with stroke service staff in stroke units and community services. 2021 , 1-10		1
39	2021 Guideline for the Prevention of Stroke in Patients With Stroke and Transient Ischemic Attack: A Guideline From the American Heart Association/American Stroke Association. 2021 , 52, e364-e467		201
38	The Effects of Interrupting Prolonged Sitting With Frequent Bouts of Light-Intensity Standing Exercises on Blood Pressure in Stroke Survivors: A Dose Escalation Trial. 2021 , 18, 988-997		
37	Association Between Physical Activity and Mortality Among Community-Dwelling Stroke Survivors. 2021 ,		1
36	Objectively Measured Physical Activity and Sedentary Time Among Adults With and Without Stroke: A National Cohort Study. 2021 , 52, e729-e732		1
35	The Effects of a Movement-to-Music (M2M) Intervention on Physical and Psychosocial Outcomes in People Poststroke: A Randomized Controlled Trial.. 2021 , 3, 100160		0

34	A Comparison of the Physiology of Sedentary Behavior and Light Physical Activity in Adults With and Without a Physical Disability. 2019 , 16, 894-901		2
33	Effectiveness of a multimodal exercise rehabilitation program on walking capacity and functionality after a stroke. 2017 , 13, 666-675		7
32	Differences in muscle thickness and echo intensity between stroke survivors and age- and sex-matched healthy older adults. 2020 , 23, 188-194		4
31	Sedentary Behavior in People with and without a Chronic Health Condition: How Much, What and When?. 2016 , 3, 503-519		8
30	Promoting Physical Activity and Nutrition in People With Stroke. 2017 , 71, 7105360010p1-7105360010p5		9
29	Actividad física, barreras y beneficios en personas con ictus crónico: estudio transversal de encuesta. 2019 , 41, 275-284		0
28	Feasibility of Aerobic Interval Training in Nonambulant Persons after Stroke. 2019 , 8, 97-101		0
27	Factors influencing sedentary behaviours after stroke: findings from qualitative observations and interviews with stroke survivors and their caregivers.		
26	What is the relationship between physical activity and cardiovascular risk factors in stroke survivors post completion of rehabilitation? Protocol for a longitudinal study. 2017 , 7, e019193		2
25	mHealth impact on secondary stroke prevention: a scoping review of randomized controlled trials among stroke survivors between 2010-2020.. 2022 , 8, 19		0
24	The Impact of Wearable Technologies in Health Research: Scoping Review (Preprint).		
23	Heart Disease and Stroke Statistics-2022 Update: A Report From the American Heart Association.. <i>Circulation</i> , 2022 , CIR0000000000001052	16.7	196
22	Fit for purpose. Co-production of complex behavioural interventions. A practical guide and exemplar of co-producing a telehealth-delivered exercise intervention for people with stroke.. <i>Health Research Policy and Systems</i> , 2022 , 20, 2	3.7	0
21	Can telerehabilitation services combined with caregiver-mediated exercises improve early supported discharge services poststroke? A study protocol for a multicentre, observer-blinded, randomized controlled trial.. <i>BMC Neurology</i> , 2022 , 22, 29	3.1	1
20	The Impact of Wearable Technologies in Health Research: Scoping Review.. <i>JMIR MHealth and UHealth</i> , 2022 , 10, e34384	5.5	4
19	Early Wearing of Knee-Ankle-Foot Orthosis Improves Functional Prognosis in Patients after Stroke.. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2022 , 31, 106261	2.8	0
18	People with Stroke Are Most Sedentary in the Afternoon and Evening.. <i>Cerebrovascular Diseases</i> , 2022 , 1-6	3.2	1
17	Appetite and energy intake following a bout of circuit resistance training in chronic hemiparetic stroke patients: a preliminary randomized controlled trial.. <i>Topics in Stroke Rehabilitation</i> , 2022 , 1-14	2.6	

16	Physical activity level of post-stroke individuals that use the Brazilian public health system. <i>Fisioterapia Em Movimento</i> , 35,	0.8	
15	Differences in paretic lower limb loading and fluidity in sit-to-walk according to selection of the leading limb in individuals with stroke.. <i>Clinical Biomechanics</i> , 2022, 94, 105639	2.2	
14	Physical activity dimensions after stroke: patterns and relation with lower limb motor function.. <i>Journal of NeuroEngineering and Rehabilitation</i> , 2021, 18, 171	5.3	1
13	Physical activity according to sit-to-stand, standing, and stand-to-sit abilities in subacute stroke with walking difficulty: a cross-sectional study.. <i>Physiotherapy Theory and Practice</i> , 2022, 1-9	1.5	
12	Between commitment and avoidance - working age stroke survivors' perceptions of physical activity and sedentary behaviour: a qualitative study.. <i>BMC Neurology</i> , 2022, 22, 183	3.1	0
11	Predictors of non-stepping time in people with chronic stroke. 1-9		
10	Study Protocol: Sleep Effects on Post-Stroke Rehabilitation Study. Publish Ahead of Print,		0
9	Relationships Among Environmental Variables, Physical Capacity, Balance Self-Efficacy, and Real-World Walking Activity Post-Stroke. 2022, 36, 535-544		0
8	The longitudinal association between movement behavior patterns and the course of participation up to one year after stroke. 1-9		
7	Accuracy of gait and posture classification using movement sensors in individuals with mobility impairment after stroke. 13,		0
6	Beyond steps per day: other measures of real-world walking after stroke related to cardiovascular risk. 2022, 19,		0
5	Exploring the perspectives of people with stroke, caregivers and healthcare professionals on the design and delivery of a mHealth adaptive physical activity intervention: a qualitative study protocol. 5, 66		0
4	Life and Leisure Activities following Stroke or Transient Ischaemic Attack (TIA): An Observational, Multi-Centre, 6-Month Follow-Up Study. 2022, 19, 13848		0
3	Multidimensional analysis of sedentary behaviour and participation in Spanish stroke survivors (Part&Sed-Stroke): a protocol for a longitudinal multicentre study. 2023, 13, e065628		0
2	Mobile health to promote physical activity in people post stroke or transient ischemic attack □ study protocol for a feasibility randomised controlled trial. 2023, 23,		0
1	Circadian Factors in Stroke: A Clinician's Perspective.		0