

Exposure to loud noise, bilateral high-frequency hearing

Occupational and Environmental Medicine

73, 34-41

DOI: [10.1136/oemed-2014-102778](https://doi.org/10.1136/oemed-2014-102778)

Citation Report

#	ARTICLE	IF	CITATIONS
1	Different types of housing and respiratory health outcomes. Preventive Medicine Reports, 2017, 7, 124-129.	0.8	14
2	Cardiovascular conditions, hearing difficulty, and occupational noise exposure within US industries and occupations. American Journal of Industrial Medicine, 2018, 61, 477-491.	1.0	105
3	Occupational noise exposure and hypertension: the Dongfeng-Tongji Cohort Study. Journal of the American Society of Hypertension, 2018, 12, 71-79.e5.	2.3	32
4	Occupational Noise Exposure, Bilateral High-Frequency Hearing Loss, and Blood Pressure. Journal of Occupational and Environmental Medicine, 2018, 60, 462-468.	0.9	17
5	Hearing loss is associated with increased CHD risk and unfavorable CHD-related biomarkers in the Dongfeng-Tongji cohort. Atherosclerosis, 2018, 271, 70-76.	0.4	16
6	The Influence of Occupational Noise Exposure on Cardiovascular and Hearing Conditions among Industrial Workers. Scientific Reports, 2019, 9, 11524.	1.6	30
7	Hearing loss is associated with increased stroke risk in the Dongfeng-Tongji Cohort. Atherosclerosis, 2019, 285, 10-16.	0.4	18
8	Occupational noise exposure: A review of its effects, epidemiology, and impact with recommendations for reducing its burden. Journal of the Acoustical Society of America, 2019, 146, 3879-3905.	0.5	147
9	Evaluation of noise risk level and its consequences on technical operators of tobacco processing equipment in a cigarette producing company in Nigeria. Scientific African, 2020, 8, e00344.	0.7	6
10	A risk model and nomogram for high-frequency hearing loss in noise-exposed workers. BMC Public Health, 2021, 21, 747.	1.2	2
11	Association of occupational noise exposure, bilateral hearing loss with atherosclerotic cardiovascular disease risk in Chinese adults. International Journal of Hygiene and Environmental Health, 2021, 235, 113776.	2.1	15
12	Ten-year risk of fatal cardiovascular disease and its association with metabolic risk factors among waste pickers in South Africa. BMC Cardiovascular Disorders, 2021, 21, 336.	0.7	2
13	Association of hearing loss with total and cause-specific mortality in US adults. Environmental Science and Pollution Research, 2022, 29, 5032-5042.	2.7	6
14	High frequency hearing impairment and cardiovascular disease in Canada: Results from the Canadian Health Measures Survey. Journal of the Acoustical Society of America, 2021, 150, 1001-1012.	0.5	1
15	Self-reported occupational noise exposure and cardiovascular disease in Canada: Results from the Canadian Health Measures Survey. Journal of the Acoustical Society of America, 2021, 150, 990-1000.	0.5	5
16	Association of occupational noise exposure, bilateral hearing loss with hypertension among Chinese workers. Journal of Hypertension, 2021, 39, 643-650.	0.3	11
17	Hearing loss is associated with increased risk of incident stroke but not coronary heart disease among middle-aged and older Chinese adults: the Dongfeng-Tongji cohort study. Environmental Science and Pollution Research, 2022, 29, 21198-21209.	2.7	4
18	Association of all Cause and Cause-Specific Mortality With Hearing Loss Among US Adults: A Secondary Analysis Study. International Journal of Public Health, 0, 67, .	1.0	1

#	ARTICLE	IF	CITATIONS
20	Association of noise exposure, plasma microRNAs with arterial stiffness among Chinese workers. Environmental Pollution, 2022, 311, 120002.	3.7	5
21	Economic Burden of Hearing Loss in Middle-Aged and Older Adults in China. Journal of Applied Gerontology, 0, , 073346482211241.	1.0	0
22	The Association between Duration of Noise Exposure in the Workplace and Glucose Metabolism Status: Evidence from the Korea National Health and Nutrition Examination Survey. Korean Journal of Family Medicine, 2022, 43, 396-402.	0.4	1