Comparative Value of Kentucky Bluegrass, Kentucky 31 Bromegrass as Pasture for Milk Cows. II. Effect of Kind Crude Protein Content and Digestibility and Intake of D

Journal of Dairy Science 39, 581-588

DOI: 10.3168/jds.s0022-0302(56)94789-0

Citation Report

#	Article	IF	CITATIONS
1	Methoxyl as an Indicator of the Nutritive Value of Forage. Journal of Dairy Science, 1958, 41, 1715-1722.	3.4	6
2	Growth, Physiological Responses, and Evidence of Toxicity in Yearling Dairy Cattle Grazing Different Grasses. Journal of Dairy Science, 1970, 53, 575-587.	3.4	52
3	Digestibility and Rate of Passage by Steers Fed Tall Fescue, Alfalfa and Orchardgrass Hay in 18 and 32 C Ambient Temperatures1. Journal of Animal Science, 1974, 39, 93-96.	0.5	82
4	Feed Intake in Hereford Calves Infused Intraperitoneally with Toxic Fescue Extract. Journal of Dairy Science, 1974, 57, 1385-1387.	3.4	11
5	Influence of Copper Supplementation on Blood and Liver Copper in Cattle Fed Tall Fescue or Quackgrass. Journal of Animal Science, 1986, 62, 263-271.	0.5	19
6	Fescue Endophyte: History and Impact on Animal Agriculture. Journal of Production Agriculture, 1988, 1, 39-44.	0.4	245
7	Nutritive Value. Agronomy, 0, , 157-201.	0.2	8
8	Cool-Season Grasses for Pasture. Agronomy, 2015, , 321-355.	0.2	4
9	A 100-Year Review: Metabolic modifiers in dairy cattle nutrition. Journal of Dairy Science, 2017, 100, 10113-10142.	3.4	34
10	History of Dairy Farming. , 2022, , 1-29.		O
11	Cow and Calf Performance on Tall Fescue- or Kentucky Bluegrass-Ladino Clover Forages. Journal of Animal Science, 1979, 49, 44-49.	0.5	4