CITATION REPORT List of articles citing

Current use of antihypertensive drugs in Japanese patients with hypertension: Analysis by age group

DOI: 10.1111/ggi.13276 Geriatrics and Gerontology International, 2018, 18, 899-906.

Source: https://exaly.com/paper-pdf/68783591/citation-report.pdf

Version: 2024-04-28

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

| # | Paper | IF | Citations |
|----|--|------|-----------|
| 12 | Treatment patterns of antihypertensive fixed-dose combinations according to age and number of agents prescribed: Retrospective analysis using a Japanese claims database. <i>Geriatrics and Gerontology International</i> , 2019 , 19, 1077-1083 | 2.9 | 2 |
| 11 | Effects of combination drugs on antihypertensive medication adherence in a real-world setting: a Korean Nationwide Study. <i>BMJ Open</i> , 2019 , 9, e029862 | 3 | 9 |
| 10 | Systematic Review with Network Meta-Analysis: Comparative Efficacy and Safety of Combination Therapy with Angiotensin II Receptor Blockers and Amlodipine in Asian Hypertensive Patients. <i>International Journal of Hypertension</i> , 2019 , 2019, 9516279 | 2.4 | 1 |
| 9 | Pharmacokinetic interactions of esaxerenone with amlodipine and digoxin in healthy Japanese subjects. <i>BMC Pharmacology & Document Subjects</i> , 21, 55 | 2.6 | 4 |
| 8 | An Analysis of Medication Prescriptions for Hypertension in Urban and Rural Residents in Tianjin. <i>Advances in Therapy</i> , 2020 , 37, 4414-4426 | 4.1 | 2 |
| 7 | Obesity predicts persistence of resistant hypertension after surgery in patients with primary aldosteronism. <i>Clinical Endocrinology</i> , 2020 , 93, 229-237 | 3.4 | 4 |
| 6 | Effective coverage of medical treatment for hypertension, diabetes and dyslipidaemia in Japan: An analysis of National Health and Nutrition Surveys 2003-2017. <i>Journal of Health Services Research and Policy</i> , 2021 , 26, 106-114 | 2.4 | 1 |
| 5 | Correlations Between SGLT-2 Inhibitors and Acute Renal Failure by Signal Detection Using FAERS: Stratified Analysis for Reporting Country and Concomitant Drugs. <i>Clinical Drug Investigation</i> , 2021 , 41, 235-243 | 3.2 | 2 |
| 4 | Effects of angiotensin II receptor blockers on serum potassium level and hyperkalemia risk: retrospective single-centre analysis. <i>European Journal of Hospital Pharmacy</i> , 2021 , | 1.6 | |
| 3 | Inverse association between hypertension treatment and COVID-19 prevalence in Japan. <i>International Journal of Infectious Diseases</i> , 2021 , 108, 517-521 | 10.5 | 2 |
| 2 | A Japan nationwide web-based survey of estimation on patients for renal denervation based on blood pressure level and the number of antihypertensives (J-NEEDs survey). <i>Journal of Clinical Hypertension</i> , 2021 , 23, 1684-1694 | 2.3 | 1 |
| 1 | Prevalence of hypertensive diseases and treated hypertensive patients in Japan: A nationwide administrative claims database study. <i>Hypertension Research</i> , 2022 , 45, 1123-1133 | 4.7 | 1 |