## CITATION REPORT List of articles citing

Remnant lipoproteins play an important role of in-stent restenosis in type 2 diabetes undergoing percutaneous coronary intervention: a single-centre observational cohort study

DOI: 10.1186/s12933-019-0819-z Cardiovascular Diabetology, 2019, 18, 11.

Source: https://exaly.com/paper-pdf/72844559/citation-report.pdf

Version: 2024-04-28

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
24	Inhibition of vascular neointima hyperplasia by FGF21 associated with FGFR1/Syk/NLRP3 inflammasome pathway in diabetic mice. <i>Atherosclerosis</i> , <b>2019</b> , 289, 132-142	3.1	22
23	Remnant Lipoprotein Cholesterol Independently Associates With In-Stent Restenosis After Drug-Eluting Stenting for Coronary Artery Disease. <i>Angiology</i> , <b>2019</b> , 70, 853-859	2.1	3
22	Comparison of remnant cholesterol levels estimated by calculated and measured LDL-C levels in Chinese patients with coronary heart disease. <i>Clinica Chimica Acta</i> , <b>2020</b> , 500, 75-80	6.2	10
21	Everolimus-eluting bioresorbable scaffolds and metallic stents in diabetic patients: a patient-level pooled analysis of the prospective ABSORB DM Benelux Study, TWENTE and DUTCH PEERS. <i>Cardiovascular Diabetology</i> , <b>2020</b> , 19, 165	8.7	
20	Prognostic impact of estimated remnant-like particle cholesterol in patients with differing glycometabolic status: an observational cohort study from China. <i>Lipids in Health and Disease</i> , <b>2020</b> , 19, 179	4.4	2
19	Visit-to-visit HbA variability is associated with in-stent restenosis in patients with type 2 diabetes after percutaneous coronary intervention. <i>Cardiovascular Diabetology</i> , <b>2020</b> , 19, 133	8.7	9
18	Could remnant-like particle cholesterol become a risk factor in diabetic menopausal women with coronary artery disease? A cross-sectional study of single academic center in China. <i>Lipids in Health and Disease</i> , <b>2020</b> , 19, 44	4.4	3
17	Long-term outcomes of medical therapy versus successful recanalisation for coronary chronic total occlusions in patients with and without type 2 diabetes mellitus. <i>Cardiovascular Diabetology</i> , <b>2020</b> , 19, 100	8.7	10
16	The atherogenic index of plasma plays an important role in predicting the prognosis of type 2 diabetic subjects undergoing percutaneous coronary intervention: results from an observational cohort study in China. <i>Cardiovascular Diabetology</i> , <b>2020</b> , 19, 23	8.7	16
15	Associations Between Bioavailable Vitamin D and Remnant Cholesterol in Patients With Type 2 Diabetes Mellitus. <i>Cureus</i> , <b>2021</b> , 13, e13248	1.2	2
14	Glycemic Control Status and Long-Term Clinical Outcomes in Diabetic Chronic Total Occlusion Patients: An Observational Study. <i>Journal of Interventional Cardiology</i> , <b>2021</b> , 2021, 5565987	1.8	
13	The Atherogenic Index of Plasma: A Powerful and Reliable Predictor for Coronary Artery Disease in Patients With Type 2 Diabetes. <i>Angiology</i> , <b>2021</b> , 72, 934-941	2.1	9
12	Evaluation of remnant cholesterol levels and Monocyte-to-HDL-cholesterol ratio in South Asian patients with acute coronary syndrome. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , <b>2021</b> , 31, 2144-2150	4.5	1
11	The Effects of Calculated Remnant-Like Particle Cholesterol on Incident Cardiovascular Disease: Insights from a General Chinese Population. <i>Journal of Clinical Medicine</i> , <b>2021</b> , 10,	5.1	0
10	The Effect of Untraditional Lipid Parameters in the Development of Coronary Artery Disease: Atherogenic Index of Plasma, Atherogenic Coefficient and Lipoprotein Combined Index. <i>Journal of the Saudi Heart Association</i> , <b>2021</b> , 33, 244-250	0.7	2
9	Atherogenic Index of Plasma and the Risk of In-Stent Restenosis in Patients with Acute Coronary Syndrome beyond the Traditional Risk Factors. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2021</b> ,	4	3
8	Atherogenic index of plasma is associated with major adverse cardiovascular events in patients with type 2 diabetes mellitus. <i>Cardiovascular Diabetology</i> , <b>2021</b> , 20, 201	8.7	4

## CITATION REPORT

7	Factors Influencing Stent Failure in Chronic Total Occlusion Coronary Intervention. <i>Interventional Cardiology Review</i> , <b>2021</b> , 16, e27	4.2	2
6	Association of Increased Remnant Cholesterol and the Risk of Coronary Artery Disease: A Retrospective Study. <i>Frontiers in Cardiovascular Medicine</i> , <b>2021</b> , 8, 740596	5.4	O
5	Elevated Remnant Cholesterol is Associated with Adverse Cardiovascular Outcomes in Patients with Acute Coronary Syndrome <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2022</b> ,	4	1
4	Diabetes and restenosis Cardiovascular Diabetology, <b>2022</b> , 21, 23	8.7	5
3	Diabetes and restenosis <i>Cardiovascular Diabetology</i> , <b>2022</b> , 21, 23  Plasma Proteome Profiling of Patients With In-stent Restenosis by Tandem Mass Tag-Based Quantitative Proteomics Approach <i>Frontiers in Cardiovascular Medicine</i> , <b>2022</b> , 9, 793405	8. <sub>7</sub>	5

Remnant Cholesterol and the Risk of Coronary Artery Disease in Patients With Type 2 Diabetes. 000331972211&10