

# Magnitude and determinants for place of postnatal care delivered at home in Ethiopia: a multinomial analysis from a health survey

Reproductive Health

16, 162

DOI: [10.1186/s12978-019-0818-2](https://doi.org/10.1186/s12978-019-0818-2)

Citation Report

#	ARTICLE	IF	CITATIONS
1	Determinants of postnatal checkup for newborns in Ethiopia: Further analysis of 2016 Ethiopia demographic and health survey. BMC Pregnancy and Childbirth, 2020, 20, 761.	2.4	4
2	Do mothers who delivered at health facilities return to health facilities for postnatal care follow-up? A multilevel analysis of the 2016 Ethiopian Demographic and Health Survey. PLoS ONE, 2021, 16, e0249793.	2.5	5
3	Coverage of RMNCH interventions: more analysis is needed. The Lancet Global Health, 2021, 9, e1037-e1038.	6.3	0
4	Barriers to Utilization of Postnatal Care: A Qualitative Study in Guinea. Open Journal of Obstetrics and Gynecology, 2021, 11, 391-402.	0.2	4
5	Postnatal Care Service Utilisation in Ethiopia: Reflecting on 20 Years of Demographic and Health Survey Data. International Journal of Environmental Research and Public Health, 2021, 18, 193.	2.6	9
6	Effectiveness of checklist-based box system intervention (CBBSI) versus routine care on improving postnatal care utilization in Northwest Ethiopia: a cluster randomized controlled trial. Reproductive Health, 2021, 18, 234.	3.1	1
7	Association between health insurance enrolment and maternal health care service utilization among women in Ethiopia. BMC Public Health, 2021, 21, 2329.	2.9	5
8	Knowledge, Practice, and Associated Factors of Essential Newborn Care among Sudanese Women in Eastern Sudan. Children, 2022, 9, 873.	1.5	0
9	Level of Postnatal Checkup in Ethiopia – Implications for Child Health Services. Frontiers in Pediatrics, 0, 10, .	1.9	0
10	Home based postpartum care and determinants in Ethiopia: A multilevel analysis. PLoS ONE, 2022, 17, e0272489.	2.5	2
11	Exploration of barriers to postnatal care service utilization in Debre Libanos District, Ethiopia: A descriptive qualitative study. Frontiers in Global Women S Health, 0, 3, .	2.3	2
12	Spatial distribution and determinants of newborns not receiving postnatal check-up within two days after birth in Ethiopia: a spatial and multilevel analysis of EDHS 2016. BMC Pediatrics, 2022, 22, .	1.7	3
13	Multilevel analysis of the predictors of completion of the continuum of maternity care in Ethiopia; using the recent 2019 Ethiopia mini demographic and health survey. BMC Pregnancy and Childbirth, 2022, 22, .	2.4	2
14	Association between frequency of mass media exposure and maternal health care service utilization among women in sub-Saharan Africa: Implications for tailored health communication and education. PLoS ONE, 2022, 17, e0275202.	2.5	11
16	Factors Associated with Post-Natal Care Utilization among the Women of Reproductive age Group: Evidence from Bangladesh Demographic and Health Survey 2017-18. Health Services Research and Managerial Epidemiology, 2022, 9, 233339282211363.	0.9	1
17	Postnatal Care Utilization Coverage and its Associated Factors Among Mothers in South Gondar Zone, Northwest Ethiopia: A Cross-Sectional Study. SAGE Open Nursing, 2022, 8, 237796082211403.	1.2	1
18	Effective coverage of newborn postnatal care in Ethiopia: Measuring inequality and spatial distribution of quality-adjusted coverage. PLoS ONE, 2023, 18, e0293520.	2.5	0
19	Spatial patterns and determinants of low utilization of delivery care service and postnatal check-up within 2 months following birth in Ethiopia: Bivariate analysis. PLoS ONE, 2024, 19, e0297038.	2.5	0

#	ARTICLE	IF	CITATIONS
21	Individual and community-level factors associated with women's utilization of postnatal care services in Uganda, 2016: a multilevel and spatial analysis. BMC Health Services Research, 2024, 24, .	2.2	0