

Clinical significance of the G8 screening tool in elderly patients with oral carcinoma

Clinical Oral Investigations

24, 1953-1961

DOI: [10.1007/s00784-019-03056-y](https://doi.org/10.1007/s00784-019-03056-y)

Citation Report

#	ARTICLE	IF	CITATIONS
1	Utility of the Geriatric 8 for the Prediction of Therapy-Related Toxicity in Older Adults with Diffuse Large B-Cell Lymphoma. <i>Oncologist</i> , 2021, 26, 215-223.	1.9	11
2	Clinical investigation of geriatric assessment for the prolongation of healthy life expectancy in older oral squamous cell carcinoma patients. <i>Journal of Japanese Society of Oral Oncology</i> , 2020, 32, 180-185.	0.0	0
3	Chemotherapy for older patients with head and neck cancer. <i>Journal of Japanese Society of Oral Oncology</i> , 2020, 32, 186-192.	0.0	0
4	Clinical usefulness of geriatric assessment in elderly patients with unresectable hepatocellular carcinoma receiving sorafenib or lenvatinib therapy. <i>Cancer Reports</i> , 2022, 5, e1613.	0.6	4
5	Examination of late stage elderly patients with oral cancer who were classified as prefrail and vulnerable. <i>Japanese Journal of Head and Neck Cancer</i> , 2022, 48, 28-33.	0.0	0