

# CITATION REPORT

List of articles citing

De-escalation of axillary surgery in breast cancer patients treated in the neoadjuvant setting: a Dutch population-based study

DOI: 10.1007/s10549-020-05589-3

Breast Cancer Research and Treatment, 2020, 180, 725-733.

**Source:** <https://exaly.com/paper-pdf/76909266/citation-report.pdf>

**Version:** 2024-04-27

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
16	Axillary surgery in breast cancer: An updated historical perspective. <i>Seminars in Oncology</i> , <b>2020</b> , 47, 341-352	3.52	15
15	Leveraging the increased rates of pathologic complete response after neoadjuvant treatment in breast cancer to de-escalate surgical treatments. <i>Journal of Surgical Oncology</i> , <b>2021</b> , 123, 71-79	2.8	3
14	Sentinel lymph node biopsy after neoadjuvant chemotherapy in patients with node-positive breast cancer: guiding a more selective axillary approach. <i>Breast Cancer Research and Treatment</i> , <b>2021</b> , 186, 527-534	4.4	8
13	Minimally Invasive Complete Response Assessment of the Breast After Neoadjuvant Systemic Therapy for Early Breast Cancer (MICRA trial): Interim Analysis of a Multicenter Observational Cohort Study. <i>Annals of Surgical Oncology</i> , <b>2021</b> , 28, 3243-3253	3.1	9
12	Targeted axillary biopsy and sentinel lymph node biopsy for axillary restaging after neoadjuvant chemotherapy. <i>Annals of Surgical Treatment and Research</i> , <b>2021</b> , 100, 305-312	2	1
11	Discrepancy of Breast and Axillary Pathologic Complete Response and Outcomes in Different Subtypes of Node-positive Breast Cancer after Neoadjuvant Chemotherapy. <i>Journal of Cancer</i> , <b>2021</b> , 12, 5365-5374	4.5	0
10	Axillary Pathologic Complete Response After Neoadjuvant Systemic Therapy by Breast Cancer Subtype in Patients With Initially Clinically Node-Positive Disease: A Systematic Review and Meta-analysis. <i>JAMA Surgery</i> , <b>2021</b> , 156, e210891	5.4	11
9	MINimal vs. MAXimal Invasive Axillary Staging and Treatment After Neoadjuvant Systemic Therapy in Node Positive Breast Cancer: Protocol of a Dutch Multicenter Registry Study (MINIMAX). <i>Clinical Breast Cancer</i> , <b>2021</b> ,	3	2
8	Factors Influencing Non-sentinel Lymph Node Involvement in Patients with Positive Sentinel Lymph Node(s) After Neoadjuvant Chemotherapy for Breast Cancer. <i>Annals of Surgical Oncology</i> ,	3.1	0
7	Is Nodal Clipping Beneficial for Node-Positive Breast Cancer Patients Receiving Neoadjuvant Chemotherapy?.		0
6	De-escalation of radiotherapy after primary chemotherapy in cT1N1 breast cancer (RAPCHEM; BOOG 201003): 5-year follow-up results of a Dutch, prospective, registry study. <b>2022</b> ,		2
5	Editorial for Different Imaging Modalities for the Diagnosis of Axillary Lymph Node Metastases in Breast Cancer: A Systematic Review and Network Meta-Analysis		
4	Evaluation of different imaging modalities for axillary lymph node staging in breast cancer patients to provide a personalized and optimized therapy algorithm.		0
3	Diagnostic Accuracy of Radioactive Iodine Seed Placement in the Axilla With Sentinel Lymph Node Biopsy After Neoadjuvant Chemotherapy in Node-Positive Breast Cancer.		5
2	Prediction of Primary Tumour and Axillary Lymph Node Response to Neoadjuvant Chemo(Targeted) Therapy with Dedicated Breast [18F]FDG PET/MRI in Breast Cancer. <b>2023</b> , 15, 401		0
1	Axillary staging and management of cN + breast cancer patients treated with neoadjuvant chemotherapy: results of a survey among breast cancer surgeons in Spain.		0