## Source apportionment and seasonal variation in particu Belgium

Environmental Science and Pollution Research 27, 14933-14943

DOI: 10.1007/s11356-020-07881-7

**Citation Report** 

#	Article	IF	CITATIONS
1	Receptor modelling and risk factors of polycyclic aromatic hydrocarbons (PAHs) in the atmospheric particulate matter at an IGP outflow location (island of the bay of Bengal—Bhola, Bangladesh). Air Quality, Atmosphere and Health, 2021, 14, 1417-1431.	3.3	10
2	Temporal Trend of Polycyclic Aromatic Hydrocarbons in Atmosphere: A Case Study within 24Âh after Snowfall. Bulletin of Environmental Contamination and Toxicology, 2022, 108, 652-657.	2.7	2
3	The Inherent Instability of Environmental Parameters Governing Indoor Air Quality on Board Ships and the Use of Temporal Trends to Identify Pollution Sources. Indoor Air, 2023, 2023, 1-19.	4.3	2
4	Concentrations of PAHs from urban and rural areas in southern nigeria using polyurethane foams (PUF) passive samplers: Health impact and cancer risk assessment. Scientific African, 2023, 22, e01976.	1.5	Ο
5	Source analysis and health risk assessment of polycyclic aromatic hydrocarbon (PAHs) in total suspended particulate matter (TSP) from Bengbu, China. Scientific Reports, 2024, 14, .	3.3	0