In situ assessment of health status and heavy metal bioaccumulation of adult Pelophylax ridibundus (Anura: Ranidae) individuals inhabiting polluted area in southern Bulgaria

DOI: 10.1016/j.ecolind.2020.106413
Ecological Indicators, 2020, 115, 106413.

#	Paper	IF	Citations
17	Exposure to common pesticides utilized in northern rice fields of Iran affects survival of non-target species, Pelophylax ridibundus (Amphibia: Ranidae). *Environmental Science and Pollution Research*, **2021**, 28, 33557	5.1	1
16	Determinations of Erythrocyte Sizes in Adult Pelophylax ridibundus (Amphibia: Anura: Ranidae) Inhabiting Industrial Area in Southern Bulgaria. *Water, Air, and Soil Pollution*, **2021**, 232, 1	2.6	1
15	Hematological parameters of a Neotropical wild frog population, with a phylogenetic perspective on blood cell composition in Anura.. *Environmental Epigenetics*, **2022**, 68, 361-369	2.4	1
14	In situ assessment of genetic and epigenetic alterations in frog Rana plancyi and Rana limnocharis inhabiting aquatic ecosystems associated with Pb/Zn/Cu mining. *Science of the Total Environment*, **2021**, 779, 146139	10.2	2
13	Vermiculite enriched by Fe(III) oxides as a novel adsorbent for toxic metals removal. *Journal of Environmental Chemical Engineering*, **2021**, 9, 106020	6.8	5
12	Utilisation of wastes for low-cost synthesis of chitosan composites with nanosized sodium aluminium silicate hydrate and geopolymer/zeolite A for the removal of Hg(II) and Pb(II) ions from aqueous media. *International Journal of Environmental Analytical Chemistry*, 1-19	1.8	2
11	Bioaccumulation of Trace Elements and Health Risk Predictions in Edible Tissues of the Marsh Frog. *Biological Trace Element Research*, **2021**, 1	4.5	1
10	Influence of nano and bulk copper on agile frog development.. *Ecotoxicology*, **2022**, 31, 357	2.9	
9	Leptodactylus macrosternum (Anura: Leptodactylidae) as a bioindicator of potentially toxic chemical elements in irrigated perimeters in northeastern Brazil. *Environmental Chemistry and Ecotoxicology*, **2022**, 4, 124-131	3.9	0
8	Are there correlations between the levels of fluctuating asymmetry in Pelophylax ridibundus (Anura: Ranidae) meristic morphological traits and morphological parameters used for assessing their physical fitness (health status)?. *Environmental Science and Pollution Research*, **2022**,	5.1	
7	Proximate Composition, Predictive Analysis and Allometric Relationships, of the Edible Water Frog (Pelophylax epeiroticus) in Lake Pamvotida (Northwest Greece). *Sustainability*, **2022**, 14, 3150	3.6	0
6	Bioaccumulation of Elemental Concentrations in Sediment and Frog (Pyxicephalus edulis) in Igbeebo River, Ondo State, Nigeria. *Chemistry Africa*,	2.2	0
5	Fat rather than health Ecotoxic responses of Bufo raddei to environmental heavy metal stress during the non-breeding season. **2022**, 244, 114040		0
4	Trace element bioaccumulation and health risk assessment derived from leg consumption of the marsh frog, Pelophylax ridibundus (Pallas, 1771). **2022**, 39, 182-190		0
3	First Report on the Presence of Toxic Metals and Metalloids in East Asian Bullfrog (Hoplobatrachus rugulosus) Legs. **2022**, 11, 3009		0
2	Neglected skin-associated microbial communities: a unique immune defense strategy of Bufo raddei under environmental heavy metal pollution.		0
1	Identification of cryptic forms of the hybridogenic complex of European water frogs (<i>Pelophylax esculentus complex</i>) in the conditions of transformed biotopes of the south of the Central Russian Upland based on DNA markers. **2022**, 20, 247-260		0