

CITATION REPORT

List of articles citing

Omission of Axillary Lymph Node Dissection is Associated with Inferior Survival in Breast Cancer Patients with Residual N1 Nodal Disease Following Neoadjuvant Chemotherapy

DOI: 10.1245/s10434-020-08928-2

Annals of Surgical Oncology, 2021, 28, 930-940.

Source: <https://exaly.com/paper-pdf/77771460/citation-report.pdf>

Version: 2024-04-27

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
21	ASO Author Reflections: Omission of Axillary Lymph Node Dissection for Breast Cancer Patients with Residual N1 Nodal Disease Following Neoadjuvant Chemotherapy: Not Ready for Primetime?. <i>Annals of Surgical Oncology</i> , 2020 , 27, 869-870	3.1	1
20	Discrepancy of Breast and Axillary Pathologic Complete Response and Outcomes in Different Subtypes of Node-positive Breast Cancer after Neoadjuvant Chemotherapy. <i>Journal of Cancer</i> , 2021 , 12, 5365-5374	4.5	0
19	Sentinel node biopsy alone for breast cancer patients with residual nodal disease after neoadjuvant chemotherapy. <i>Scientific Reports</i> , 2021 , 11, 9056	4.9	1
18	Axillary surgery in node-positive breast cancer. <i>Breast</i> , 2021 ,	3.6	1
17	De-Escalating Axillary Surgery in Node-Positive Breast Cancer Treated with Neoadjuvant Systemic Therapy.. <i>Breast Care</i> , 2021 , 16, 584-589	2.4	0
16	Customizing local and systemic therapies for women with early breast cancer: the St. Gallen International Consensus Guidelines for treatment of early breast cancer 2021. <i>Annals of Oncology</i> , 2021 , 32, 1216-1235	10.3	44
15	Tailored axillary surgery in patients with clinically node-positive breast cancer: Pre-planned feasibility substudy of TAXIS (OPBC-03, SAKK 23/16, IBCSG 57-18, ABCSG-53, GBG 101). <i>Breast</i> , 2021 , 60, 98-110	3.6	1
14	Survival and recurrence with or without axillary dissection in patients with invasive breast cancer and sentinel node metastasis. <i>Scientific Reports</i> , 2021 , 11, 19893	4.9	
13	De-escalation of axillary irradiation for early breast cancer - Has the time come?. <i>Cancer Treatment Reviews</i> , 2021 , 101, 102297	14.4	2
12	Breast and axillary surgery after neoadjuvant systemic treatment - A review of clinical routine recommendations and the latest clinical research.. <i>Breast</i> , 2022 ,	3.6	0
11	Postmastectomy Radiation Therapy in Patients With Minimally Involved Lymph Nodes: A Review of the Current Data and Future Directions.. <i>Journal of Breast Cancer</i> , 2022 , 25, 1-12	3	0
10	Management of the positive sentinel lymph node following neoadjuvant chemotherapy: results of a survey conducted with breast surgeons.. <i>Ecancermedicalscience</i> , 2022 , 16, 1357	2.7	
9	Impact of COVID-19 Disease in Early Breast Cancer Management: A Summary of the Current Evidence. <i>JCO Global Oncology</i> , 2022 ,	3.7	0
8	Consenso de la Sociedad Española de Senología y Patología Mamaria (SESPM) sobre la biopsia selectiva del ganglio centinela (BSGC) y el manejo axilar en el cáncer de mama (2022). 2022 , 35, 243-259		0
7	Omission of axillary lymph node dissection in patients with ypN+ breast cancer after neoadjuvant chemotherapy: A retrospective multicenter study (KROG 21-06). 2022 ,		0
6	Tailored axillary surgery [A novel concept for clinically node positive breast cancer. 2023 , 69, 281-289		0
5	Regional radiotherapy after primary systemic treatment for cN+ breast cancer patients. 2023 , 68, 181-188		0

- 4 Safety and Accuracy of Sentinel Lymph Node Biopsy Alone in Clinically Node-Positive Patients Undergoing Upfront Surgery for Invasive Breast Cancer: A Systematic Review. **2023**, 30, 3102-3110 ○
- 3 Non-sentinel node metastasis prediction during surgery in breast cancer patients with one to three positive sentinel node(s) following neoadjuvant chemotherapy. **2023**, 13, ○
- 2 Axillary response and outcome in breast cancer patients after neoadjuvant treatment: The role of radiotherapy in reducing recurrence in ypN0 patients with initially cN+ stage. 13, ○
- 1 Level I axillary dissection in patients with breast cancer and tumor-involved sentinel lymph node after NAC is not sufficient for adequate nodal staging. **2023**, 39, 1-6 ○