

CITATION REPORT

List of articles citing

Patient-reported outcomes at 1 and 2 years after total hip and knee arthroplasty: what is the minimum required follow-up?

DOI: 10.1007/s00402-021-03819-x

Archives of Orthopaedic and Trauma Surgery, 2021, , 1.

Source: <https://exaly.com/paper-pdf/79166890/citation-report.pdf>

Version: 2024-04-09

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
7	Are Minimum Two-Year PROMS Necessary for Accurate Assessment of Patient Outcomes after Primary TKA?. <i>Journal of Arthroplasty</i> , 2022 ,	4.4	1
6	No difference in patient-reported satisfaction after 12 months between customised individually made and off-the-shelf total knee arthroplasty.. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2022 , 1	5.5	1
5	Osteoarthritic Severity in Unresurfaced Patellae Does Not Adversely Affect Patient-reported Outcomes in Contemporary Primary TKA.. <i>Journal of the American Academy of Orthopaedic Surgeons Global Research and Reviews</i> , 2022 , 6,	1.2	0
4	Medial pivot versus (cam post) posterior stabilised total knee arthroplasty, systematic review and meta-analysis of 3837 knees.. <i>Acta Orthopaedica Belgica</i> , 2021 , 87, 665-680	1.3	0
3	An anatomic-functional implant positioning technique with robotic assistance for primary TKA allows the restoration of the native knee alignment and a natural functional ligament pattern, with a faster recovery at 6 months compared to an adjusted mechanical technique.. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2022 ,	5.5	0
2	Selective Patella Resurfacing in Contemporary Cruciate Retaining and Substituting Total Knee Arthroplasty: A Matched Cohort Analysis. 2022 ,		0
1	Trajectory of chronic and neuropathic pain, anxiety and depressive symptoms and pain catastrophizing after total knee replacement. Results of a prospective, single-center study at a mean follow-up of 7.5 years. 2023 , 103543		0