

Predication of oxygen requirement in COVID-19 patient inflammatory markers: CRP, hypertension, age, neutrop

Scientific Reports

11, 13026

DOI: [10.1038/s41598-021-92418-2](https://doi.org/10.1038/s41598-021-92418-2)

Citation Report

#	ARTICLE	IF	CITATIONS
1	HbA1c level may be a risk factor for oxygen therapy requirement in patients with coronavirus disease 2019. <i>Journal of Diabetes Investigation</i> , 2021, , .	1.1	4
2	Predicting the necessity of oxygen therapy in the early stage of COVID-19 using machine learning. <i>Medical and Biological Engineering and Computing</i> , 2022, 60, 957-968.	1.6	13
4	Novel Scoring Systems to Predict the Need for Oxygenation and ICU Care, and Mortality in Hospitalized COVID-19 Patients: A Risk Stratification Tool. <i>Cureus</i> , 2022, , .	0.2	1
5	Laboratory markers of severity across three COVID-19 outbreaks in Australia: has Omicron and vaccinations changed disease presentation?. <i>Internal and Emergency Medicine</i> , 2023, 18, 43-52.	1.0	8
6	In-hospital Mortality Rates in SARS-CoV-2 Patients Treated with Enoxaparin and Heparin. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2022, 28, 107602962211318.	0.7	1
8	Evaluation of emerging inflammatory markers for predicting oxygen support requirement in COVID-19 patients. <i>PLoS ONE</i> , 2022, 17, e0278145.	1.1	2