

Cardiac MRI shows an association of lower cardiorespiratory
myocardial mass and higher cardiac stiffness in the general population
Heart

Progress in Cardiovascular Diseases

68, 25-35

DOI: [10.1016/j.pcad.2021.09.003](https://doi.org/10.1016/j.pcad.2021.09.003)

Citation Report

#	ARTICLE	IF	CITATIONS
1	Low cardiopulmonary fitness is associated with higher liver fat content and higher γ -glutamyltransferase concentrations in the general population – The Sedentary's Liver. Liver International, 2022, 42, 585-594.	3.9	3
2	SHIP-MR and Radiology: 12 Years of Whole-Body Magnetic Resonance Imaging in a Single Center. Healthcare (Switzerland), 2022, 10, 33.	2.0	11
3	The Role of MR Assessments of Cardiac Morphology, Function, and Tissue Characteristics on Exercise Capacity in Well-Functioning Older Adults. Journal of Magnetic Resonance Imaging, 0, , .	3.4	0
4	Lower aldosterone concentrations are associated with a smaller and thinner heart in the general population. The Study of Health in Pomerania (SHIP). European Journal of Preventive Cardiology, 0, , .	1.8	0
5	Cardiac structural and functional abnormalities in epilepsy: A systematic review and meta-analysis. Epilepsia Open, 2023, 8, 46-59.	2.4	5
6	Too Little of a Good Thing. JACC: Cardiovascular Imaging, 2023, 16, 768-778.	5.3	9
7	Left ventricular hypertrophy as a risk factor for accelerated brain aging: Results from the Study of Health in Pomerania. Human Brain Mapping, 2024, 45, .	3.6	0
8	Lower muscular strength is associated with greater liver fat content and higher serum liver enzymes – The Sedentary's Liver. The Study of Health in Pomerania. European Journal of Sport Science, 0, , .	2.7	0