CITATION REPORT List of articles citing

An open, comparative, multicentre clinical study of combined oral therapy with sildenafil and doxazosin GITS for treating Chinese patients with erectile dysfunction and lower urinary tract symptoms secondary to benign prostatic hyperplasia

DOI: 10.1038/aja.2010.177 Asian Journal of Andrology, 2011, 13, 630-5.

Source: https://exaly.com/paper-pdf/84833475/citation-report.pdf

Version: 2024-04-09

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
16	Phosphodiesterase Type 5 Inhibitor and Erectile Dysfunction in Lower Urinary Tract Symptoms. <i>LUTS: Lower Urinary Tract Symptoms</i> , 2012 , 4 Suppl 1, 75-80	1.9	2
15	Systematic review of combination drug therapy for non-neurogenic male lower urinary tract symptoms. <i>European Urology</i> , 2013 , 64, 228-43	10.2	67
14	Phosphodiesterase-5 expression and function in the lower urinary tract: a critical review. <i>Urology</i> , 2013 , 81, 480-7	1.6	12
13	The Use of Phosphodiesterase Type 5 Inhibitors in the Treatment of Lower Urinary Tract Symptoms Due to Benign Prostatic Hyperplasia. 2014 , 113-128		
12	The efficacy of PDE5 inhibitors alone or in combination with alpha-blockers for the treatment of erectile dysfunction and lower urinary tract symptoms due to benign prostatic hyperplasia: a systematic review and meta-analysis. <i>Journal of Sexual Medicine</i> , 2014 , 11, 1539-45	1.1	34
11	Systematic review and meta-analysis on phosphodiesterase 5 inhibitors and Eadrenoceptor antagonists used alone or combined for treatment of LUTS due to BPH. <i>Asian Journal of Andrology</i> , 2015 , 17, 1022-32	2.8	15
10	Efficacy and safety of solifenacin plus tamsulosin oral controlled absorption system in men with lower urinary tract symptoms: a meta-analysis. <i>Asian Journal of Andrology</i> , 2015 , 17, 124-34	2.8	4
9	Useful Implications of Low-dose Long-term Use of PDE-5 Inhibitors. <i>Sexual Medicine Reviews</i> , 2016 , 4, 270-284	5.6	11
8	Sildenafil citrate in combination with tamsulosin versus tamsulosin monotherapy for management of male lower urinary tract symptoms due to benign prostatic hyperplasia: A randomised, double-blind, placebo-controlled trial. <i>Arab Journal of Urology Arab Association of Urology</i> , 2017 , 15, 53	1.7 3-59	12
7	Phosphodiesterase inhibitors for lower urinary tract symptoms consistent with benign prostatic hyperplasia. <i>The Cochrane Library</i> , 2018 , 11, CD010060	5.2	8
6	Medical management of benign prostatic hyperplasia. 2018 , 509-525		
5	Efficacy and safety of PDE5-Is and II blockers for treating lower ureteric stones or LUTS: a meta-analysis of RCTs. <i>BMC Urology</i> , 2018 , 18, 30	2.2	5
4	Assessment of Combination Therapies vs Monotherapy for Erectile Dysfunction: A Systematic Review and Meta-analysis. <i>JAMA Network Open</i> , 2021 , 4, e2036337	10.4	4
3	Systematic Review of Oral Combination Therapy for Erectile Dysfunction When Phosphodiesterase Type 5 Inhibitor Monotherapy Fails. <i>Sexual Medicine Reviews</i> , 2019 , 7, 430-441	5.6	13
2	Comparative effectiveness of oral drug therapies for lower urinary tract symptoms due to benign prostatic hyperplasia: a systematic review and network meta-analysis. <i>PLoS ONE</i> , 2014 , 9, e107593	3.7	33
1	Study of phosphodiesterase 5 inhibitors and Eddrenoceptor antagonists used alone or in combination for the treatment of lower urinary tract symptoms due to benign prostatic hyperplasia. <i>Minerva Urologica E Nefrologica = the Italian Journal of Urology and Nephrology</i> , 2020 , 72, 13-21	4.4	9