The Submerged Valley through Mackinac Straits

Journal of Geology 46, 966-974

DOI: 10.1086/624711

Citation Report

#	Article	IF	CITATIONS
1	Bedrock Topography and Pleistocene Glacial Lobes in Central United States. Journal of Geology, 1956, 64, 101-116.	1.4	30
2	Late Quaternary vertical movements in eastern North America: Quantitative evidence of glacioâ€isostatic rebound. Reviews of Geophysics, 1972, 10, 849-884.	23.0	228
3	Sedimentary processes in the Great Lakes. Reviews of Geophysics, 1981, 19, 635-648.	23.0	43
4	Oscillations of levels and cool phases of the Laurentian Great Lakes caused by inflows from glacial Lakes Agassiz and Barlow-Ojibway. Journal of Paleolimnology, 1989, 2, 99.	1.6	124
5	Stratigraphy and paleolimnologic record of lower Holocene sediments in northern Lake Huron and Georgian Bay. Canadian Journal of Earth Sciences, 1994, 31, 1586-1605.	1.3	55
6	Lakes of the Huron basin: their record of runoff from the laurentide ice sheet. Quaternary Science Reviews, 1994, 13, 891-922.	3.0	93
7	Holocene lake-level fluctuations and beach-ridge development along the northern shore of Lake Michigan, USA. Journal of Paleolimnology, 1996, 15, 147.	1.6	24
8	The history and architecture of lacustrine depositional systems in the northern Lake Michigan basin. Journal of Paleolimnology, 1999, 22, 475-496.	1.6	3
9	Title is missing!. Journal of Paleolimnology, 2003, 29, 221-234.	1.6	20
	Title 15 Mil55ing Journal of Fulcommittology, 2003, 25, 221 25 1.	1.6	20
10	Glacial Isostatic Adjustment of the Laurentian Great Lakes Basin: Using the Empirical Record of Strandline Deformation for Reconstruction of Early Holocene Paleo-Lakes and Discovery of a Hydrologically Closed Phase*. Géographie Physique Et Quaternaire, 2005, 59, 187-210.	0.2	48
	Glacial Isostatic Adjustment of the Laurentian Great Lakes Basin: Using the Empirical Record of Strandline Deformation for Reconstruction of Early Holocene Paleo-Lakes and Discovery of a		
10	Glacial Isostatic Adjustment of the Laurentian Great Lakes Basin: Using the Empirical Record of Strandline Deformation for Reconstruction of Early Holocene Paleo-Lakes and Discovery of a Hydrologically Closed Phase*. Géographie Physique Et Quaternaire, 2005, 59, 187-210. A multiproxy environmental investigation of Holocene wood from a submerged conifer forest in Lake	0.2	48
10	Glacial Isostatic Adjustment of the Laurentian Great Lakes Basin: Using the Empirical Record of Strandline Deformation for Reconstruction of Early Holocene Paleo-Lakes and Discovery of a Hydrologically Closed Phase*. Géographie Physique Et Quaternaire, 2005, 59, 187-210. A multiproxy environmental investigation of Holocene wood from a submerged conifer forest in Lake Huron, USA. Quaternary Research, 2006, 66, 67-77. The Stanley unconformity in Lake Huron basin: evidence for a climate-driven closed lowstand about 7900 14C BP, with similar implications for the Chippewa lowstand in Lake Michigan basin. Journal of	0.2	13
10 11 12	Glacial Isostatic Adjustment of the Laurentian Great Lakes Basin: Using the Empirical Record of Strandline Deformation for Reconstruction of Early Holocene Paleo-Lakes and Discovery of a Hydrologically Closed Phase*. GÃ@ographie Physique Et Quaternaire, 2005, 59, 187-210. A multiproxy environmental investigation of Holocene wood from a submerged conifer forest in Lake Huron, USA. Quaternary Research, 2006, 66, 67-77. The Stanley unconformity in Lake Huron basin: evidence for a climate-driven closed lowstand about 7900 14C BP, with similar implications for the Chippewa lowstand in Lake Michigan basin. Journal of Paleolimnology, 2007, 37, 435-452. Stratigraphy, age, and flora of the Southport forest bed, Southeastern Wisconsin. Journal of Great	0.2 1.7 1.6	48 13 54
10 11 12 13	Glacial Isostatic Adjustment of the Laurentian Great Lakes Basin: Using the Empirical Record of Strandline Deformation for Reconstruction of Early Holocene Paleo-Lakes and Discovery of a Hydrologically Closed Phase*. Géographie Physique Et Quaternaire, 2005, 59, 187-210. A multiproxy environmental investigation of Holocene wood from a submerged conifer forest in Lake Huron, USA. Quaternary Research, 2006, 66, 67-77. The Stanley unconformity in Lake Huron basin: evidence for a climate-driven closed lowstand about 7900 14C BP, with similar implications for the Chippewa lowstand in Lake Michigan basin. Journal of Paleolimnology, 2007, 37, 435-452. Stratigraphy, age, and flora of the Southport forest bed, Southeastern Wisconsin. Journal of Great Lakes Research, 2009, 35, 538-547.	1.7 1.6 1.9	48 13 54 2
10 11 12 13	Glacial Isostatic Adjustment of the Laurentian Great Lakes Basin: Using the Empirical Record of Strandline Deformation for Reconstruction of Early Holocene Paleo-Lakes and Discovery of a Hydrologically Closed Phase*. GÃ@ographie Physique Et Quaternaire, 2005, 59, 187-210. A multiproxy environmental investigation of Holocene wood from a submerged conifer forest in Lake Huron, USA. Quaternary Research, 2006, 66, 67-77. The Stanley unconformity in Lake Huron basin: evidence for a climate-driven closed lowstand about 7900 14C BP, with similar implications for the Chippewa lowstand in Lake Michigan basin. Journal of Paleolimnology, 2007, 37, 435-452. Stratigraphy, age, and flora of the Southport forest bed, Southeastern Wisconsin. Journal of Great Lakes Research, 2009, 35, 538-547. GEOLOGIC HISTORY OF GREAT LAKES BEACHES. Coastal Engineering Proceedings, 1953, , 5. Oscillations of levels and cool phases of the Laurentian Great Lakes caused by inflows from glacial	1.7 1.6 1.9	48 13 54 2